lieu of a Government-furnished headstone or marker.

(Authority: 38 U.S.C. 2306(d))

(h) The monetary allowance in lieu of a Government-furnished headstone or marker is not payable if death occurred on or after November 1, 1990.

(Authority: Pub. L. 101-508)

[44 FR 58711, Oct. 11, 1979, as amended at 47 FR 19131, May 4, 1982; 49 FR 19653, May 9, 1984; 51 FR 17629, May 14, 1986; 52 FR 34910, Sept. 16, 1987; 55 FR 50323, Dec. 6, 1990; 56 FR 25045, June 3, 1991; 56 FR 65851, Dec. 19, 1991; 61 FR 20727, May 8, 1996]

### PART 4—SCHEDULE FOR RATING DISABILITIES

### Subpart A—General Policy in Rating

Sec.

- 4.1 Essentials of evaluative rating.
- 4.2 Interpretation of examination reports.
- Resolution of reasonable doubt. 4.3
- Evaluation of evidence.
- Higher of two evaluations. 4.7
- Congenital or developmental defects.
- 4.10 Functional impairment.
- 4.13 Effect of change of diagnosis.
- 4.14 Avoidance of pyramiding.
- Total disability ratings. 4.15
- 4.16 Total disability ratings for compensation based on unemployability of the individual.
- 4.17 Total disability ratings for pension based on unemployability and age of the individual.
- 4.17a Misconduct etiology.
- 4.18 Unemployability.
- 4.19 Age in service-connected claims.
- Analogous ratings. 4.20
- Application of rating schedule.
- 4.22 Rating of disabilities aggravated by active service.
- 4.23 Attitude of rating officers.
- 4.24 Correspondence.
- Combined ratings table. 4.25
- 4.26 Bilateral factor.
- Use of diagnostic code numbers. 4 27
- 4.28 Prestabilization rating from date of discharge from service.
- 4.29 Ratings for service-connected disabilities requiring hospital treatment or observation.
- 4.30 Convalescent ratings.
- 4.31 Zero percent evaluations.

### Subpart B—Disability Ratings

THE MUSCULOSKELETAL SYSTEM

- 4.40 Functional loss.
- 4.41 History of injury.

- 4.42 Complete medical examination of injury cases.
  - Osteomyelitis.
- 4.44 The bones.
- 4.45 The joints.
- 4.46 Accurate measurement.
- 4.47—4.54 [Reserved]
- 4.55 Principles of combined ratings for muscle injuries.
- 4.56 Evaluation of muscle disabilities.
- 4.57 Static foot deformities.
- Arthritis due to strain.
- Painful motion.
- 4.60 [Reserved]
- 4.61 Examination Circulatory disturbances.
- 4.63 Loss of use of hand or foot.
- 4.64 Loss of use of both buttocks.
- 4.65[Reserved]
- 4.66 Sacroiliac joint.
- 4.67 Pelvic bones.
- 4.68 Amputation rule.
- 4 69 Dominant hand. 4.70
- Inadequate examinations.
- 4.71 Measurement of ankylosis and joint motion.
- 4.71a Schedule of ratings—musculoskeletal system. 4.72 [P
- [Reserved]
- 4.73 Schedule of ratings—muscle injuries.

### THE ORGANS OF SPECIAL SENSE

- Examination of visual acuity.
- Examination of field vision.
- 4.76a Computation of average concentric contraction of visual fields.
- 4.77 Examination of muscle function.
- 4.78 Computing aggravation.4.79 Loss of use of one eye, having only light perception.
- 4.80 Rating of one eye.
- 4.81—4.82 [Reserved]
- 4.83 Ratings at scheduled steps and distances.
- 4.83a Impairment of central visual acuity.
- 4.84 Differences between distant and near visual acuity
- 4.84a Schedule of ratings—eye.

### IMPAIRMENT OF AUDITORY ACUITY

- 4 85 Evaluation of hearing impairment.
- 4.86 Exceptional patterns of hearing impairment.
- 4.87 Schedule of ratings-ear.
- 4.87a Schedule of ratings-other sense or-

INFECTIOUS DISEASES, IMMUNE DISORDERS AND NUTRITIONAL DEFICIENCIES

- 4.88 [Reserved]
- Chronic fatigue syndrome.
- Schedule of ratings-infectious diseases, immune disorders and nutritional deficiencies.
- 4.88c Ratings for inactive nonpulmonary tuberculosis initially entitled after August 19, 1968.

### § 4.1

4.89 Ratings for inactive nonpulmonary tuberculosis in effect on August 19, 1968.

#### THE RESPIRATORY SYSTEM

- 4.96 Special provisions regarding evaluation of respiratory conditions.
- 4.97 Schedule of ratings—respiratory system.

#### THE CARDIOVASCULAR SYSTEM

- 4.100-4.103 [Reserved]
- 4.104 Schedule of ratings—cardiovascular system.

#### THE DIGESTIVE SYSTEM

- 4.110 Ulcers.
- 4.111 Postgastrectomy syndromes.
- 4.112 Weight loss
- 4.113 Coexisting abdominal conditions.
- 4.114 Schedule of ratings—digestive system.

#### THE GENITOURINARY SYSTEM

- 4.115 Nephritis.
- 4.115a Ratings of the genitourinary system—dysfunctions.
- 4.115b Ratings of the genitourinary system—diagnoses.

# GYNECOLOGICAL CONDITIONS AND DISORDERS OF THE BREAST

4.116 Schedule of ratings—gynecological conditions and disorders of the breast.

### THE HEMIC AND LYMPHATIC SYSTEMS

4.117 Schedule of ratings—hemic and lymphatic systems.

### THE SKIN

4.118 Schedule of ratings—skin.

### THE ENDOCRINE SYSTEM

4.119 Schedule of ratings—endocrine system.

# NEUROLOGICAL CONDITIONS AND CONVULSIVE DISORDERS

- 4.120 Evaluations by comparison.
- 4.121 Identification of epilepsy.
- 4.122 Psychomotor epilepsy.
- 4.123 Neuritis, cranial or peripheral.
- 4.124 Neuralgia, cranial or peripheral.
- 4.124a Schedule of ratings—neurological conditions and convulsive disorders.

### MENTAL DISORDERS

- 4.125 Diagnosis of mental disorders.
- 4.126 Evaluation of disability from mental disorders.
- 4.127 Mental retardation and personality disorders.
- ${\it 4.128} \ \ \, {\it Convalescence \ ratings \ following \ extended \ hospitalization}.$
- 4.129 Mental disorders due to traumatic stress.

4.130 Schedule of ratings—mental disorders.

DENTAL AND ORAL CONDITIONS

- 4.149 [Reserved]
- 4.150 Schedule of ratings—dental and oral conditions.
- APPENDIX A TO PART 4—TABLE OF AMEND-MENTS AND EFFECTIVE DATES SINCE 1946
- APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES
- APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES

AUTHORITY: 38 U.S.C. 1155, unless otherwise

SOURCE: 29 FR 6718, May 22, 1964, unless otherwise noted.

# Subpart A—General Policy in Rating

### §4.1 Essentials of evaluative rating.

This rating schedule is primarily a guide in the evaluation of disability resulting from all types of diseases and injuries encountered as a result of or incident to military service. The percentage ratings represent as far as can practicably be determined the average impairment in earning capacity resulting from such diseases and injuries and their residual conditions in civil occupations. Generally, the degrees of disability specified are considered adequate to compensate for considerable loss of working time from exacerbations or illnesses proportionate to the severity of the several grades of disability. For the application of this schedule, accurate and fully descriptive medical examinations are required, with emphasis upon the limitation of activity imposed by the disabling condition. Over a period of many years, a veteran's disability claim may require reratings in accordance with changes in laws, medical knowledge and his or her physical or mental condition. It is thus essential, both in the examination and in the evaluation of disability, that each disability be viewed in relation to its history.

[41 FR 11292, Mar. 18, 1976]

# § 4.2 Interpretation of examination reports.

Different examiners, at different times, will not describe the same disability in the same language. Features

of the disability which must have persisted unchanged may be overlooked or a change for the better or worse may not be accurately appreciated or described. It is the responsibility of the rating specialist to interpret reports of examination in the light of the whole recorded history, reconciling the various reports into a consistent picture so that the current rating may accurately reflect the elements of disability present. Each disability must be considered from the point of view of the veteran working or seeking work. If a diagnosis is not supported by the findings on the examination report or if the report does not contain sufficient detail, it is incumbent upon the rating board to return the report as inadequate for evaluation purposes.

[41 FR 11292, Mar. 18, 1976]

### §4.3 Resolution of reasonable doubt.

It is the defined and consistently applied policy of the Department of Veterans Affairs to administer the law under a broad interpretation, consistent, however, with the facts shown in every case. When after careful consideration of all procurable and assembled data, a reasonable doubt arises regarding the degree of disability such doubt will be resolved in favor of the claimant. See §3.102 of this chapter.

[40 FR 42535, Sept. 15, 1975]

### §4.6 Evaluation of evidence.

The element of the weight to be accorded the character of the veteran's service is but one factor entering into the considerations of the rating boards in arriving at determinations of the evaluation of disability. Every element in any way affecting the probative value to be assigned to the evidence in each individual claim must be thoroughly and conscientiously studied by each member of the rating board in the light of the established policies of the Department of Veterans Affairs to the end that decisions will be equitable and just as contemplated by the requirements of the law.

### § 4.7 Higher of two evaluations.

Where there is a question as to which of two evaluations shall be applied, the higher evaluation will be assigned if the disability picture more nearly approximates the criteria required for that rating. Otherwise, the lower rating will be assigned.

# § 4.9 Congenital or developmental defects.

Mere congenital or developmental defects, absent, displaced or supernumerary parts, refractive error of the eye, personality disorder and mental deficiency are not diseases or injuries in the meaning of applicable legislation for disability compensation purposes.

[41 FR 11292, Mar. 18, 1976]

#### §4.10 Functional impairment.

The basis of disability evaluations is the ability of the body as a whole, or of the psyche, or of a system or organ of the body to function under the ordinary conditions of daily life including employment. Whether the upper or lower extremities, the back or abdominal wall, the eyes or ears, or the cardiovascular, digestive, or other system, or psyche are affected, evaluations are based upon lack of usefulness, of these parts or systems, especially in self-support. This imposes upon the medical examiner the responsibility of furnishing, in addition to the etiological, anatomical, pathological, laboratory and prognostic data required for ordinary medical classification, full description of the effects of disability upon the person's ordinary activity. In this connection, it will be remembered that a person may be too disabled to engage in employment although he or she is up and about and fairly comfortable at home or upon limited activ-

[41 FR 11292, Mar. 18, 1976]

### §4.13 Effect of change of diagnosis.

The repercussion upon a current rating of service connection when change is made of a previously assigned diagnosis or etiology must be kept in mind. The aim should be the reconciliation and continuance of the diagnosis or etiology upon which service connection for the disability had been granted. The relevant principle enunciated in §4.125, entitled "Diagnosis of mental

disorders," should have careful attention in this connection. When any change in evaluation is to be made, the rating agency should assure itself that there has been an actual change in the conditions, for better or worse, and not merely a difference in thoroughness of the examination or in use of descriptive terms. This will not, of course, preclude the correction of erroneous ratings, nor will it preclude assignment of a rating in conformity with §4.7.

[29 FR 6718, May 22, 1964, as amended at 61 FR 52700, Oct. 8, 1996]

#### §4.14 Avoidance of pyramiding.

The evaluation of the same disability under various diagnoses is to be avoided. Disability from injuries to the muscles, nerves, and joints of an extremity may overlap to a great extent, so that special rules are included in the appropriate bodily system for their evaluation. Dyspnea, tachycardia, nervousness, fatigability, etc., may result from many causes; some may be service connected, others, not. Both the use of manifestations not resulting from service-connected disease or injury in establishing the service-connected evaluation, and the evaluation of the same manifestation under different diagnoses are to be avoided.

### §4.15 Total disability ratings.

The ability to overcome the handicap of disability varies widely among individuals. The rating, however, is based primarily upon the average impairment in earning capacity, that is, upon the economic or industrial handicap which must be overcome and not from individual success in overcoming it. However, full consideration must be given to unusual physical or mental effects in individual cases, to peculiar effects of occupational activities, to defects in physical or mental endowment preventing the usual amount of success in overcoming the handicap of disability and to the effect of combinations of disability. Total disability will be considered to exist when there is present any impairment of mind or body which is sufficient to render it impossible for the average person to follow a substantially gainful occupation; Provided, That permanent total disability shall be taken to exist when

the impairment is reasonably certain to continue throughout the life of the disabled person. The following will be considered to be permanent total disability: the permanent loss of the use of both hands, or of both feet, or of one hand and one foot, or of the sight of both eyes, or becoming permanently helpless or permanently bedridden. Other total disability ratings are scheduled in the various bodily systems of this schedule.

### §4.16 Total disability ratings for compensation based on unemployability of the individual.

(a) Total disability ratings for compensation may be assigned, where the schedular rating is less than total, when the disabled person is, in the judgment of the rating agency, unable to secure or follow a substantially gainful occupation as a result of service-connected disabilities: Provided That, if there is only one such disability, this disability shall be ratable at 60 percent or more, and that, if there are two or more disabilities, there shall be at least one disability ratable at 40 percent or more, and sufficient additional disability to bring the combined rating to 70 percent or more. For the above purpose of one 60 percent disability, or one 40 percent disability in combination, the following will be considered as one disability: (1) Disabilities of one or both upper extremities, or of one or both lower extremities, including the bilateral factor, if applicable, (2) disabilities resulting from common etiology or a single accident, (3) disabilities affecting a single body system, e.g. orthopedic, digestive, respiratory, cardiovascular-renal, neuropsychiatric, (4) multiple injuries incurred in action, or (5) multiple disabilities incurred as a prisoner of war. It is provided further that the existence or degree of nonservice-connected previous disabilities or unemployability status will be dis-regarded where the percentages referred to in this paragraph for the service-connected disability or disabilities are met and in the judgment of the rating agency such service-connected disabilities render the veteran unemployable. Marginal employment shall not

be considered substantially gainful employment. For purposes of this section, marginal employment generally shall be deemed to exist when a veteran's earned annual income does not exceed the amount established by the U.S. Department of Commerce, Bureau of the Census, as the poverty threshold for one person. Marginal employment may also be held to exist, on a facts found basis (includes but is not limited to employment in a protected environment such as a family business or sheltered workshop), when earned annual income exceeds the poverty threshold. Consideration shall be given in all claims to the nature of the employment and the reason for termination.

(Authority: 38 U.S.C. 501)

(b) It is the established policy of the Department of Veterans Affairs that all veterans who are unable to secure and follow a substantially gainful occupation by reason of service-connected disabilities shall be rated totally disabled. Therefore, rating boards should submit to the Director, Compensation and Pension Service, for extra-schedular consideration all cases of veterans who are unemployable by reason of service-connected disabilities, but who fail to meet the percentage standards set forth in paragraph (a) of this section. The rating board will include a full statement as to the veteran's service-connected disabilities, employment history, educational and vocational attainment and all other factors having a bearing on the issue.

[40 FR 42535, Sept. 15, 1975, as amended at 54 FR 4281, Jan. 30, 1989; 55 FR 31580, Aug. 3, 1990; 58 FR 39664, July 26, 1993; 61 FR 52700, Oct. 8, 1996]

# §4.17 Total disability ratings for pension based on unemployability and age of the individual.

All veterans who are basically eligible and who are unable to secure and follow a substantially gainful occupation by reason of disabilities which are likely to be permanent shall be rated as permanently and totally disabled. For the purpose of pension, the permanence of the percentage requirements of §4.16 is a requisite. When the percentage requirements are met, and the disabilities involved are of a permanent

nent nature, a rating of permanent and total disability will be assigned if the veteran is found to be unable to secure and follow substantially gainful employment by reason of such disability. Prior employment or unemployment status is immaterial if in the judgment of the rating board the veteran's disabilities render him or her unemployable. In making such determinations, the following guidelines will be used:

(a) Marginal employment, for example, as a self-employed farmer or other person, while employed in his or her own business, or at odd jobs or while employed at less than half the usual remuneration will not be considered incompatible with a determination of unemployability, if the restriction, as to securing or retaining better employment, is due to disability.

(b) Claims of all veterans who fail to meet the percentage standards but who meet the basic entitlement criteria and are unemployable, will be referred by the rating board to the Adjudication Officer under §3.321(b)(2) of this chapter

(Authority: 38 U.S.C. 1155; 38 U.S.C. 3102) [43 FR 45348, Oct. 2, 1978, as amended at 56 FR 57985, Nov. 15, 1991]

### §4.17a Misconduct etiology.

A permanent and total disability rating under the provisions of §§4.15, 4.16 and 4.17 will not be precluded by reason of the coexistence of misconduct disability when:

(a) A veteran, regardless of employment status, also has innocently acquired 100 percent disability, or

(b) Where unemployable, the veteran has other disabilities innocently acquired which meet the percentage requirements of §§4.16 and 4.17 and would render, in the judgment of the rating agency, the average person unable to secure or follow a substantially gainful occupation.

 $[40\ FR\ 42536,\ Sept.\ 15,\ 1975,\ as\ amended\ at\ 43\ FR\ 45349,\ Oct.\ 2,\ 1978]$ 

### §4.18 Unemployability.

A veteran may be considered as unemployable upon termination of employment which was provided on account of disability, or in which special consideration was given on account of

#### § 4.19

the same, when it is satisfactorily shown that he or she is unable to secure further employment. With amputations, sequelae of fractures and other residuals of traumatism shown to be of static character, a showing of continuous unemployability from date of incurrence, or the date the condition reached the stabilized level, is a general requirement in order to establish the fact that present unemployability is the result of the disability. However, consideration is to be given to the circumstances of employment in individual claims, and, if the employment was only occasional, intermittent, tryout or unsuccessful, or eventually terminated on account of the disability, present unemployability may be attributed to the static disability. Where unemployability for pension previously has been established on the basis of combined service-connected and nonservice-connected disabilities and the service-connected disability or disabilities have increased in severity, §4.16 is for consideration.

[40 FR 42536, Sept. 15, 1975, as amended at 43 FR 45349, Oct. 2, 1978]

### § 4.19 Age in service-connected claims.

Age may not be considered as a factor in evaluating service-connected disability; and unemployability, in service-connected claims, associated with advancing age or intercurrent disability, may not be used as a basis for a total disability rating. Age, as such, is a factor only in evaluations of disability not resulting from service, i.e., for the purposes of pension.

[29 FR 6718, May 22, 1964, as amended at 43 FR 45349, Oct. 2, 1978]

### § 4.20 Analogous ratings.

When an unlisted condition is encountered it will be permissible to rate under a closely related disease or injury in which not only the functions affected, but the anatomical localization and symptomatology are closely analogous. Conjectural analogies will be avoided, as will the use of analogous ratings for conditions of doubtful diagnosis, or for those not fully supported by clinical and laboratory findings. Nor will ratings assigned to organic dis-

eases and injuries be assigned by analogy to conditions of functional origin.

### §4.21 Application of rating schedule.

In view of the number of atypical instances it is not expected, especially with the more fully described grades of disabilities, that all cases will show all the findings specified. Findings sufficiently characteristic to identify the disease and the disability therefrom, and above all, coordination of rating with impairment of function will, however, be expected in all instances.

[41 FR 11293, Mar. 18, 1976]

# § 4.22 Rating of disabilities aggravated by active service.

In cases involving aggravation by active service, the rating will reflect only the degree of disability over and above the degree existing at the time of entrance into the active service, whether the particular condition was noted at the time of entrance into the active service, or it is determined upon the evidence of record to have existed at that time. It is necessary therefore, in all cases of this character to deduct from the present degree of disability the degree, if ascertainable, of the disability existing at the time of entrance into active service, in terms of the rating schedule, except that if the disability is total (100 percent) no deduction will be made. The resulting difference will be recorded on the rating sheet. If the degree of disability at the time of entrance into the service is not ascertainable in terms of the schedule, no deduction will be made.

### §4.23 Attitude of rating officers.

It is to be remembered that the majority of applicants are disabled persons who are seeking benefits of law to which they believe themselves entitled. In the exercise of his or her functions, rating officers must not allow their personal feelings to intrude; an antagonistic, critical, or even abusive attitude on the part of a claimant should not in any instance influence the officers in the handling of the case. Fairness and courtesy must at all times be shown to applicants by all employees

whose duties bring them in contact, directly or indirectly, with the Department's claimants.

[41 FR 11292, Mar. 18, 1976]

#### §4.24 Correspondence.

All correspondence relative to the interpretation of the schedule for rating disabilities, requests for advisory opinions, questions regarding lack of clarity or application to individual cases involving unusual difficulties, will be addressed to the Director, Compensation and Pension Service. A clear statement will be made of the point or points upon which information is desired, and the complete case file will be simultaneously forwarded to Central Office. Rating agencies will assure themselves that the recent report of physical examination presents an adequate picture of the claimant's condition. Claims in regard to which the schedule evaluations are considered inadequate or excessive, and errors in the schedule will be similarly brought to attention.

[41 FR 11292, Mar. 18, 1976]

### §4.25 Combined ratings table.

Table I, Combined Ratings Table, results from the consideration of the efficiency of the individual as affected first by the most disabling condition, then by the less disabling condition, then by other less disabling conditions, if any, in the order of severity. Thus, a person having a 60 percent disability is considered 40 percent efficient. Proceeding from this 40 percent efficiency, the effect of a further 30 percent disability is to leave only 70 percent of the efficiency remaining after consideration of the first disability, or 28 percent efficiency altogether. The individual is thus 72 percent disabled, as shown in table I opposite 60 percent and under 30 percent.

(a) To use table I, the disabilities will first be arranged in the exact order of their severity, beginning with the greatest disability and then combined with use of table I as hereinafter indicated. For example, if there are two disabilities, the degree of one disability will be read in the left column and the degree of the other in the top row, whichever is appropriate. The figures

appearing in the space where the column and row intersect will represent the combined value of the two. This combined value will then be converted to the nearest number divisible by 10, and combined values ending in 5 will be adjusted upward. Thus, with a 50 percent disability and a 30 percent disability, the combined value will be found to be 65 percent, but the 65 percent must be converted to 70 percent to represent the final degree of disability. Similarly, with a disability of 40 percent, and another disability of 20 percent, the combined value is found to be 52 percent, but the 52 percent must be converted to the nearest degree divisible by 10, which is 50 percent. If there are more than two disabilities, the disabilities will also be arranged in the exact order of their severity and the combined value for the first two will be found as previously described for two disabilities. The combined value, exactly as found in table I, will be combined with the degree of the third disability (in order of severity). The combined value for the three disabilities will be found in the space where the column and row intersect, and if there are only three disabilities will be converted to the nearest degree divisible by 10, adjusting final 5's upward. Thus, if there are three disabilities ratable at 60 percent, 40 percent, and 20 percent, respectively, the combined value for the first two will be found opposite 60 and under 40 and is 76 percent. This 76 will be combined with 20 and the combined value for the three is 81 percent. This combined value will be converted to the nearest degree divisible by 10 which is 80 percent. The same procedure will be employed when there are four or more disabilities. (See table I).

(b) Except as otherwise provided in this schedule, the disabilities arising from a single disease entity, e.g., arthritis, multiple sclerosis, cerebrovascular accident, etc., are to be rated separately as are all other disabiling conditions, if any. All disabilities are then to be combined as described in paragraph (a) of this section. The conversion to the nearest degree divisible by 10 will be done only once per rating decision, will follow the combining of

# § 4.25

all disabilities, and will be the last procedure in determining the combined degree of disability.

TABLE I—COMBINED RATINGS TABLE [10 combined with 10 is 19]

	ı.	TO COITIDI	nea with	10 15 19						
		10	20	30	40	50	60	70	80	90
19		27	35	43	51	60	68	76	84	92
		28	36	44	52	60	68	76	84	92
		29	37	45	53	61	68	76	84	92
		30	38 38	45 46	53 54	61 62	69	77 77	84 85	92
		31 32	39	46 47	54 54	62	69 70	77	85	92 92
		33	40	48	55	63	70	78	85	93
		33	41	48	56	63	70	78	85	93
		34	42	49	56	64	71	78	85	93
		35	42	50	57	64	71	78	86	93
		36	43	50	57	65	72	79	86	93
		37	44	51 52	58	65	72	79 79	86	93
		38 39	45 46	52 52	59 59	66 66	72 73	80	86 86	93 93
		40	46	53	60	67	73	80	87	93
		41	47	54	60	67	74	80	87	93
		42	48	55	61	68	74	81	87	94
		42	49	55	62	68	74	81	87	94
		43	50	56	62	69	75	81	87	94
		44	50	57	63	69	75	81	88	94
		45	51	57	63	70	76 76	82	88	94
		46 47	52 53	58 59	64 65	70 71	76 76	82 82	88 88	94 94
		48	54	59	65	71	77	83	88	94
		49	54	60	66	72	77	83	89	94
		50	55	61	66	72	78	83	89	94
		51	56	62	67	73	78	84	89	95
		51	57	62	68	73	78	84	89	95
		52	58	63	68	74	79	84	89	95
		53	58	64	69	74 75	79	84	90	95
		54 55	59 60	64 65	69 70	75 75	80 80	85 85	90 90	95 95
		56	61	66	71	76	80	85	90	95
		57	62	66	71	76	81	86	90	95
		58	62	67	72	77	81	86	91	95
54		59	63	68	72	77	82	86	91	95
		60	64	69	73	78	82	87	91	96
		60	65	69	74	78	82	87	91	96
		61	66	70	74	79 79	83	87	91	96
		62 63	66 67	71 71	75 75	79 80	83 84	87 88	92 92	96 96
		64	68	72	76	80	84	88	92	96
		65	69	73	77	81	84	88	92	96
		66	70	73	77	81	85	89	92	96
		67	70	74	78	82	85	89	93	96
		68	71	75	78	82	86	89	93	96
		69 69	72 73	76 76	79	83 83	86	90 90	93 93	97 97
		70	73 74	76	80 80	84	86 87	90	93	97
		71	74	78	81	84	87	90	94	97
		72	75	78	81	85	88	91	94	97
		73	76	79	82	85	88	91	94	97
		74	77	80	83	86	88	91	94	97
		75	78	80	83	86	89	92	94	97
		76 77	78 79	81 82	84 84	87 87	89 90	92 92	95 95	97 97
		78	79 80	83	85	88	90	92	95 95	98
		78	81	83	86	88	90	93	95	98
		79	82	84	86	89	91	93	95	98
		80	82	85	87	89	91	93	96	98
79		81	83	85	87	90	92	94	96	98
		82	84	86	88	90	92	94	96	98
		83	85	87	89	91	92	94	96	98
		84	86	87	89	91	93	95	96	98
		85 86	86 87	88 89	90 90	92 92	93 94	95 95	97 97	98 98
04		00	01	69	90 1	92	94	95	9/	96

TABLE I—COMBINED RATINGS TABLE—Continued
[10 combined with 10 is 19]

	10	20	30	40	50	60	70	80	90
85	87	88	90	91	93	94	96	97	99
86	87	89	90	92	93	94	96	97	99
87	88	90	91	92	94	95	96	97	99
88	89	90	92	93	94	95	96	98	99
89	90	91	92	93	95	96	87	38	99
90	91	92	93	94	95	96	97	98	99
91	92	93	94	95	96	96	97	98	99
92	93	94	94	95	96	97	98	98	99
93	94	94	95	96	97	97	98	99	99
94	95	95	96	96	97	98	98	99	99

(Authority: 38 U.S.C. 1155)

[41 FR 11293, Mar. 18, 1976, as amended at 54 FR 27161, June 28, 1989; 54 FR 36029, Aug. 31, 1989]

### §4.26 Bilateral factor.

When a partial disability results from disease or injury of both arms, or of both legs, or of paired skeletal muscles, the ratings for the disabilities of the right and left sides will be combined as usual, and 10 percent of this value will be added (i.e., not combined) before proceeding with further combinations, or converting to degree of disability. The bilateral factor will be applied to such bilateral disabilities before other combinations are carried out and the rating for such disabilities including the bilateral factor in this section will be treated as 1 disability for the purpose of arranging in order of severity and for all further combinations. For example, with disabilities evaluated at 60 percent, 20 percent, 10 percent and 10 percent (the two 10's representing bilateral disabilities), the order of severity would be 60, 21 and 20. The 60 and 21 combine to 68 percent and the 68 and 20 to 74 percent, converted to 70 percent as the final degree of disability.

(a) The use of the terms "arms" and "legs" is not intended to distinguish between the arm, forearm and hand, or the thigh, leg, and foot, but relates to the upper extremities and lower extremities as a whole. Thus with a compensable disability of the right thigh, for example, amputation, and one of the left foot, for example, pes planus, the bilateral factor applies, and similarly whenever there are compensable disabilities affecting use of paired ex-

tremities regardless of location or specified type of impairment.

(b) The correct procedure when applying the bilateral factor to disabilities affecting both upper extremities and both lower extremities is to combine the ratings of the disabilities affecting the 4 extremities in the order of their individual severity and apply the bilateral factor by adding, not combining, 10 percent of the combined value thus attained.

(c) The bilateral factor is not applicable unless there is partial disability of compensable degree in each of 2 paired extremities, or paired skeletal muscles.

### § 4.27 Use of diagnostic code numbers.

The diagnostic code numbers appearing opposite the listed ratable disabilities are arbitrary numbers for the purpose of showing the basis of the evaluation assigned and for statistical analysis in the Department of Veterans Affairs, and as will be observed, extend from 5000 to a possible 9999. Great care will be exercised in the selection of the applicable code number and in its citation on the rating sheet. No other numbers than these listed or hereafter furnished are to be employed for rating purposes, with an exception as described in this section, as to unlisted conditions. When an unlisted disease, injury, or residual condition is encountered, requiring rating by analogy, the diagnostic code number will be "builtup" as follows: The first 2 digits will be selected from that part of the schedule most closely identifying the part, or system, of the body involved; the last 2 digits will be "99" for all unlisted conditions. This procedure will facilitate a close check of new and unlisted conditions, rated by analogy. In the selection of code numbers, injuries will generally be represented by the number assigned to the residual condition on the basis of which the rating is determined. With diseases, preference is to be given to the number assigned to the disease itself; if the rating is determined on the basis of residual conditions, the number appropriate to the residual condition will be added, preceded by a hyphen. Thus, rheumatoid (atrophic) arthritis rated as ankylosis of the lumbar spine should be coded "5002-5289." In this way, the exact source of each rating can be easily identified. In the citation of disabilities on rating sheets, the diagnostic terminology will be that of the medical examiner, with no attempt to translate the terms into schedule nomenclature. Residuals of diseases or therapeutic procedures will not be cited without reference to the basic disease.

[41 FR 11293, Mar. 18, 1976]

# § 4.28 Prestabilization rating from date of discharge from service.

The following ratings may be assigned, in lieu of ratings prescribed elsewhere, under the conditions stated for disability from any disease or injury. The prestabilization rating is not to be assigned in any case in which a total rating is immediately assignable under the regular provisions of the schedule or on the basis of individual unemployability. The prestabilization 50-percent rating is not to be used in any case in which a rating of 50 percent or more is immediately assignable under the regular provisions.

	Rating
Unstabilized condition with severe disability— Substantially gainful employment is not fea- sible or advisable	100
Material impairment of employability likely	50

NOTE (1): Department of Veterans Affairs examination is not required prior to assignment of prestabilization ratings; however, the fact that examination was accomplished will not preclude assignment of these benefits. Prestabilization ratings are for assignment in the immediate postdischarge period. They will continue for a 12-month period following discharge from service. However,

prestabilization ratings may be changed to a regular schedular total rating or one authorizing a greater benefit at any time. In each prestabilization rating an examination will be requested to be accomplished not earlier than 6 months nor more than 12 months following discharge. In those prestabilization ratings in which following examination reduction in evaluation is found to be warranted, the higher evaluation will be continued to the end of the 12th month following discharge or to the end of the period provided under §3.105(e) of this chapter, whichever is later. Special monthly compensation should be assigned concurrently in these cases whenever records are adequate to establish entitlement.

NOTE (2): Diagnosis of disease, injury, or residuals will be cited, with diagnostic code number assigned from this rating schedule for conditions listed therein.

[35 FR 11906, July 24, 1970]

# § 4.29 Ratings for service-connected disabilities requiring hospital treatment or observation.

A total disability rating (100 percent) will be assigned without regard to other provisions of the rating schedule when it is established that a service-connected disability has required hospital treatment in a Department of Veterans Affairs or an approved hospital for a period in excess of 21 days or hospital observation at Department of Veterans Affairs expense for a service-connected disability for a period in excess of 21 days.

(a) Subject to the provisions of paragraphs (d), (e), and (f) of this section this increased rating will be effective the first day of continuous hospitalization and will be terminated effective the last day of the month of hospital discharge (regular discharge or release to non-bed care) or effective the last day of the month of termination of treatment or observation for the service-connected disability. A temporary release which is approved by an attending Department of Veterans Affairs physician as part of the treatment plan will not be considered an absence.

(1) An authorized absence in excess of 4 days which begins during the first 21 days of hospitalization will be regarded as the equivalent of hospital discharge effective the first day of such authorized absence. An authorized absence of 4 days or less which results in a total of more than 8 days of authorized absence

during the first 21 days of hospitalization will be regarded as the equivalent of hospital discharge effective the ninth day of authorized absence.

(2) Following a period of hospitalization in excess of 21 days, an authorized absence in excess of 14 days or a third consecutive authorized absence of 14 days will be regarded as the equivalent of hospital discharge and will interrupt hospitalization effective on the last day of the month in which either the authorized absence in excess of 14 days or the third 14 day period begins, except where there is a finding that convalescence is required as provided by paragraph (e) or (f) of this section. The termination of these total ratings will not be subject to §3.105(e) of this chapter.

(b) Notwithstanding that hospital admission was for disability not connected with service, if during such hospitalization, hospital treatment for a service-connected disability is instituted and continued for a period in excess of 21 days, the increase to a total rating will be granted from the first day of such treatment. If service connection for the disability under treatment is granted after hospital admission, the rating will be from the first day of hospitalization if otherwise in order.

(c) The assignment of a total disability rating on the basis of hospital treatment or observation will not preclude the assignment of a total disability rating otherwise in order under other provisions of the rating schedule, and consideration will be given to the propriety of such a rating in all instances and to the propriety of its continuance after discharge. Particular attention, with a view to proper rating under the rating schedule, is to be given to the claims of veterans discharged from hospital, regardless of length of hospitalization, with indications on the final summary of expected confinement to bed or house, or to inability to work with requirement of frequent care of physician or nurse at home.

(d) On these total ratings Department of Veterans Affairs regulations governing effective dates for increased benefits will control.

(e) The total hospital rating if convalescence is required may be continued for periods of 1, 2, or 3 months in addition to the period provided in paragraph (a) of this section.

(f) Extension of periods of 1, 2 or 3 months beyond the initial 3 months may be made upon approval of the Ad-

judication Officer.

(g) Meritorious claims of veterans who are discharged from the hospital with less than the required number of days but need post-hospital care and a prolonged period of convalescence will be referred to the Director, Compensation and Pension Service, under § 3.321(b)(1) of this chapter.

[29 FR 6718, May 22, 1964, as amended at 41 FR 11294, Mar. 18, 1976; 41 FR 34256, Aug. 13, 1976; 54 FR 4281, Jan. 30, 1989; 54 FR 34981, Aug. 23, 1989]

### §4.30 Convalescent ratings.

A total disability rating (100 percent) will be assigned without regard to other provisions of the rating schedule when it is established by report at hospital discharge (regular discharge or release to non-bed care) or outpatient release that entitlement is warranted under paragraph (a) (1), (2) or (3) of this section effective the date of hospital admission or outpatient treatment and continuing for a period of 1, 2, or 3 months from the first day of the month following such hospital discharge or outpatient release. The termination of these total ratings will not be subject to §3.105(e) of this chapter. Such total rating will be followed by appropriate schedular evaluations. When the evidence is inadequate to assign a schedular evaluation, a physical examination will be scheduled and considered prior to the termination of a total rating under this section.

(a) Total ratings will be assigned under this section if treatment of a service-connected disability resulted in:

(1) Surgery necessitating at least one month of convalescence (Effective as to outpatient surgery March 1, 1989.)

(2) Surgery with severe postoperative residuals such as incompletely healed surgical wounds, stumps of recent amputations, therapeutic immobilization of one major joint or more, application of a body cast, or the necessity for

house confinement, or the necessity for continued use of a wheelchair or crutches (regular weight-bearing prohibited). (Effective as to outpatient surgery March 1, 1989.)

(3) Immobilization by cast, without surgery, of one major joint or more. (Effective as to outpatient treatment March 10, 1976.)

A reduction in the total rating will not be subject to §3.105(e) of this chapter. The total rating will be followed by an open rating reflecting the appropriate schedular evaluation; where the evidence is inadequate to assign the schedular evaluation, a physcial examination will be scheduled prior to the end of the total rating period.

- (b) A total rating under this section will require full justification on the rating sheet and may be extended as follows:
- (1) Extensions of 1, 2 or 3 months beyond the initial 3 months may be made under paragraph (a) (1), (2) or (3) of this section.
- (2) Extensions of 1 or more months up to 6 months beyond the initial 6 months period may be made under paragraph (a) (2) or (3) of this section upon approval of the Adjudication Officer.

[41 FR 34256, Aug. 13, 1976, as amended at 54 FR 4281, Jan. 30, 1989]

### §4.31 Zero percent evaluations.

In every instance where the schedule does not provide a zero percent evaluation for a diagnostic code, a zero percent evaluation shall be assigned when the requirements for a compensable evaluation are not met.

[58 FR 52018, Oct. 6, 1993]

## Subpart B—Disability Ratings

THE MUSCULOSKELETAL SYSTEM

### §4.40 Functional loss.

Disability of the musculoskeletal system is primarily the inability, due to damage or infection in parts of the system, to perform the normal working movements of the body with normal excursion, strength, speed, coordination and endurance. It is essential that the examination on which ratings are based adequately portray the anatom-

ical damage, and the functional loss, with respect to all these elements. The functional loss may be due to absence of part, or all, of the necessary bones, joints and muscles, or associated structures, or to deformity, adhesions, defective innervation, or other pathology, or it may be due to pain, supported by adequate pathology and evidenced by the visible behavior of the claimant undertaking the motion. Weakness is as important as limitation of motion, and a part which becomes painful on use must be regarded as seriously disabled. A little used part of the musculoskeletal system may be expected to show evidence of disuse, either through atrophy, the condition of the skin, absence of normal callosity or the like.

#### §4.41 History of injury.

In considering the residuals of injury, it is essential to trace the medical-industrial history of the disabled person from the original injury, considering the nature of the injury and the attendant circumstances, and the requirements for, and the effect of, treatment over past periods, and the course of the recovery to date. The duration of the initial, and any subsequent, period of total incapacity, especially periods reflecting delayed union, inflammation, swelling, drainage, or operative intervention, should be given close attention. This consideration, or the absence of clear cut evidence of injury, may result in classifying the disability as not of traumatic origin, either reflecting congenital or developmental etiology, or the effects of healed disease.

# § 4.42 Complete medical examination of injury cases.

The importance of complete medical examination of injury cases at the time of first medical examination by the Department of Veterans Affairs cannot be overemphasized. When possible, this should include complete neurological and psychiatric examination, and other special examinations indicated by the physical condition, in addition to the required general and orthopedic or surgical examinations. When complete examinations are not conducted covering

all systems of the body affected by disease or injury, it is impossible to visualize the nature and extent of the service connected disability. Incomplete examination is a common cause of incorrect diagnosis, especially in the neurological and psychiatric fields, and frequently leaves the Department of Veterans Affairs in doubt as to the presence or absence of disabling conditions at the time of the examination.

#### §4.43 Osteomyelitis.

Chronic, or recurring, suppurative osteomyelitis, once clinically identified, including chronic inflammation of bone marrow, cortex, or periosteum, should be considered as a continuously disabling process, whether or not an actively discharging sinus or other obvious evidence of infection is manifest from time to time, and unless the focus is entirely removed by amputation will entitle to a permanent rating to be combined with other ratings for residual conditions, however, not exceeding amputation ratings at the site of election.

### §4.44 The bones.

The osseous abnormalities incident to trauma or disease, such as malunion with deformity throwing abnormal stress upon, and causing malalignment of joint surfaces, should be depicted from study and observation of all available data, beginning with inception of injury or disease, its nature, degree of prostration, treatment and duration of convalescence, and progress of recovery with development of permanent residuals. With shortening of a long bone, some degree of angulation is to be expected; the extent and direction should be brought out by X-ray and observation. The direction of angulation and extent of deformity should be carefully related to strain on the neighboring joints, especially those connected with weight-bearing.

### §4.45 The joints.

As regards the joints the factors of disability reside in reductions of their normal excursion of movements in different planes. Inquiry will be directed to these considerations:

(a) Less movement than normal (due to ankylosis, limitation or blocking,

adhesions, tendon-tie-up, contracted scars, etc.).

- (b) More movement than normal (from flail joint, resections, nonunion of fracture, relaxation of ligaments, etc.).
- (c) Weakened movement (due to muscle injury, disease or injury of peripheral nerves, divided or lengthened tendons, etc.).
  - (d) Excess fatigability.
- (e) Incoordination, impaired ability to execute skilled movements smoothly.
- (f) Pain on movement, swelling, deformity or atrophy of disuse. Instability of station, disturbance of locomotion, interference with sitting, standing and weight-bearing are related considerations. For the purpose of rating disability from arthritis, the shoulder, elbow, wrist, hip, knee, and ankle are considered major joints; multiple involvements of the interphalangeal, metacarpal and carpal joints of the upper extremities, the interphalangeal, metatarsal and tarsal joints of the lower extremities, the cervical vertebrae, the dorsal vertebrae, and the lumbar vertebrae, are considered groups of minor joints, ratable on a parity with major joints. lumbosacral articulation and both sacroiliac joints are considered to be a group of minor joints, ratable on disturbance of lumbar spine functions.

### § 4.46 Accurate measurement.

Accurate measurement of the length of stumps, excursion of joints, dimensions and location of scars with respect to landmarks, should be insisted on. The use of a goniometer in the measurement of limitation of motion is indispensable in examinations conducted within the Department of Veterans Affairs. Muscle atrophy must also be accurately measured and reported.

[41 FR 11294, Mar. 18, 1976]

### §§ 4.47—4.54 [Reserved]

# § 4.55 Principles of combined ratings for muscle injuries.

(a) A muscle injury rating will not be combined with a peripheral nerve paralysis rating of the same body part, unless the injuries affect entirely different functions.

### § 4.56

- (b) For rating purposes, the skeletal muscles of the body are divided into 23 muscle groups in 5 anatomical regions: 6 muscle groups for the shoulder girdle and arm (diagnostic codes 5301 through 5306); 3 muscle groups for the forearm and hand (diagnostic codes 5307 through 5309); 3 muscle groups for the foot and leg (diagnostic codes 5310 through 5312); 6 muscle groups for the pelvic girdle and thigh (diagnostic codes 5313 through 5318); and 5 muscle groups for the torso and neck (diagnostic codes 5319 through 5323).
- (c) There will be no rating assigned for muscle groups which act upon an ankylosed joint, with the following exceptions:
- (1) In the case of an ankylosed knee, if muscle group XIII is disabled, it will be rated, but at the next lower level than that which would otherwise be assigned.
- (2) In the case of an ankylosed shoulder, if muscle groups I and II are severely disabled, the evaluation of the shoulder joint under diagnostic code 5200 will be elevated to the level for unfavorable ankylosis, if not already assigned, but the muscle groups themselves will not be rated.
- (d) The combined evaluation of muscle groups acting upon a single unankylosed joint must be lower than the evaluation for unfavorable ankylosis of that joint, except in the case of muscle groups I and II acting upon the shoulder.
- (e) For compensable muscle group injuries which are in the same anatomical region but do not act on the same joint, the evaluation for the most severely injured muscle group will be increased by one level and used as the combined evaluation for the affected muscle groups.
- (f) For muscle group injuries in different anatomical regions which do not act upon ankylosed joints, each muscle group injury shall be separately rated and the ratings combined under the provisions of §4.25.

(Authority: 38 U.S.C. 1155) [62 FR 30237, June 3, 1997]

# § 4.56 Evaluation of muscle disabilities.

- (a) An open comminuted fracture with muscle or tendon damage will be rated as a severe injury of the muscle group involved unless, for locations such as in the wrist or over the tibia, evidence establishes that the muscle damage is minimal.
- (b) A through-and-through injury with muscle damage shall be evaluated as no less than a moderate injury for each group of muscles damaged.
- (c) For VA rating purposes, the cardinal signs and symptoms of muscle disability are loss of power, weakness, lowered threshold of fatigue, fatigue-pain, impairment of coordination and uncertainty of movement.
- (d) Under diagnostic codes 5301 through 5323, disabilities resulting from muscle injuries shall be classified as slight, moderate, moderately severe or severe as follows:
- (1) Slight disability of muscles—(i) Type of injury. Simple wound of muscle without debridement or infection.
- (ii) History and complaint. Service department record of superficial wound with brief treatment and return to duty. Healing with good functional results. No cardinal signs or symptoms of muscle disability as defined in paragraph (c) of this section.
- (iii) Objective findings. Minimal scar. No evidence of fascial defect, atrophy, or impaired tonus. No impairment of function or metallic fragments retained in muscle tissue.
- (2) Moderate disability of muscles—(i) Type of injury. Through and through or deep penetrating wound of short track from a single bullet, small shell or shrapnel fragment, without explosive effect of high velocity missile, residuals of debridement, or prolonged infection.
- (ii) History and complaint. Service department record or other evidence of in-service treatment for the wound. Record of consistent complaint of one or more of the cardinal signs and symptoms of muscle disability as defined in paragraph (c) of this section, particularly lowered threshold of fatigue after average use, affecting the particular

functions controlled by the injured muscles.

(iii) Objective findings. Entrance and (if present) exit scars, small or linear, indicating short track of missile through muscle tissue. Some loss of deep fascia or muscle substance or impairment of muscle tonus and loss of power or lowered threshold of fatigue when compared to the sound side.

(3) Moderately severe disability of muscles—(i) Type of injury. Through and through or deep penetrating wound by small high velocity missile or large low-velocity missile, with debridement, prolonged infection, or sloughing of soft parts, and intermuscular scarring.

- (ii) History and complaint. Service department record or other evidence showing hospitalization for a prolonged period for treatment of wound. Record of consistent complaint of cardinal signs and symptoms of muscle disability as defined in paragraph (c) of this section and, if present, evidence of inability to keep up with work requirements.
- (iii) Objective findings. Entrance and (if present) exit scars indicating track of missile through one or more muscle groups. Indications on palpation of loss of deep fascia, muscle substance, or normal firm resistance of muscles compared with sound side. Tests of strength and endurance compared with sound side demonstrate positive evidence of impairment.
- (4) Severe disability of muscles—(i) Type of injury. Through and through or deep penetrating wound due to high-velocity missile, or large or multiple low velocity missiles, or with shattering bone fracture or open comminuted fracture with extensive debridement, prolonged infection, or sloughing of soft parts, intermuscular binding and scarring.
- (ii) History and complaint. Service department record or other evidence showing hospitalization for a prolonged period for treatment of wound. Record of consistent complaint of cardinal signs and symptoms of muscle disability as defined in paragraph (c) of this section, worse than those shown for moderately severe muscle injuries, and, if present, evidence of inability to keep up with work requirements.
- (iii) Objective findings. Ragged, depressed and adherent scars indicating

wide damage to muscle groups in missile track. Palpation shows loss of deep fascia or muscle substance, or soft flabby muscles in wound area. Muscles swell and harden abnormally in contraction. Tests of strength, endurance, or coordinated movements compared with the corresponding muscles of the uninjured side indicate severe impairment of function. If present, the following are also signs of severe muscle disability:

- (A) X-ray evidence of minute multiple scattered foreign bodies indicating intermuscular trauma and explosive effect of the missile.
- (B) Adhesion of scar to one of the long bones, scapula, pelvic bones, sacrum or vertebrae, with epithelial sealing over the bone rather than true skin covering in an area where bone is normally protected by muscle.
- (C) Diminished muscle excitability to pulsed electrical current in electrodiagnostic tests.
- (D) Visible or measurable atrophy.(E) Adaptive contraction of an oppos-

(E) Adaptive contraction of an oppoing group of muscles.

(F) Atrophy of muscle groups not in the track of the missile, particularly of the trapezius and serratus in wounds of the shoulder girdle.

(G) Induration or atrophy of an entire muscle following simple piercing by a projectile.

(Authority: 38 U.S.C. 1155 [62 FR 30238, June 3, 1997]

### §4.57 Static foot deformities.

It is essential to make an initial distinction between bilateral flatfoot as a congenital or as an acquired condition. The congenital condition, with depression of the arch, but no evidence of abnormal callosities, areas of pressure, strain or demonstrable tenderness, is a congenital abnormality which is not compensable or pensionable. In the acquired condition, it is to be remembered that depression of the longitudinal arch, or the degree of depression, is not the essential feature. The attention should be given to anatomical changes, as compared to normal, in the relationship of the foot and leg, particularly to the inward rotation of the superior portion of the os calcis, medial deviation of the insertion of the

Achilles tendon, the medial tilting of the upper border of the astragalus. This is an unfavorable mechanical relationship of the parts. A plumb line dropped from the middle of the patella falls inside of the normal point. The forepart of the foot is abducted, and the foot everted. The plantar surface of the foot is painful and shows demonstrable tenderness, and manipulation of the foot produces spasm of the Achilles tendon, peroneal spasm due to adhesion about the peroneal sheaths, and other evidence of pain and limited motion. The symptoms should be apparent without regard to exercise. In severe cases there is gaping of bones on the inner border of the foot, and rigid valgus position with loss of the power of inversion and adduction. Exercise with undeveloped or unbalanced musculature, producing chronic irritation, can be an aggravating factor. In the absence of trauma or other definite evidence of aggravation, service connection is not in order for pes cavus which is a typically congenital or juvenile disease.

### §4.58 Arthritis due to strain.

With service incurred lower extremity amputation or shortening, a disabling arthritis, developing in the same extremity, or in both lower extremities, with indications of earlier, or more severe, arthritis in the injured extremity, including also arthritis of the lumbosacral joints and lumbar spine, if associated with the leg amputation or shortening, will be considered as service incurred, provided, however, that arthritis affecting joints not directly subject to strain as a result of the service incurred amputation will not be granted service connection. This will generally require separate evaluation of the arthritis in the joints directly subject to strain. Amputation, or injury to an upper extremity, is not considered as a causative factor with subsequently developing arthritis, except in joints subject to direct strain or actually injured.

## §4.59 Painful motion.

With any form of arthritis, painful motion is an important factor of disability, the facial expression, wincing, etc., on pressure or manipulation,

should be carefully noted and definitely related to affected joints. Muscle spasm will greatly assist the identification. Sciatic neuritis is not uncommonly caused by arthritis of the spine. The intent of the schedule is to recognize painful motion with joint or periarticular pathology as productive of disability. It is the intention to recognize actually painful, unstable, or malaligned joints, due to healed injury, as entitled to at least the minimum compensable rating for the joint. Crepitation either in the soft tissues such as the tendons or ligaments, or crepitation within the joint structures should be noted carefully as points of contact which are diseased. Flexion elicits such manifestations. The joints involved should be tested for pain on both active and passive motion, in weight-bearing and nonweight-bearing and, if possible, with the range of the opposite undamaged joint.

## §4.60 [Reserved]

### §4.61 Examination.

With any form of arthritis (except traumatic arthritis) it is essential that the examination for rating purposes cover all major joints, with especial reference to Heberden's or Haygarth's nodes.

### §4.62 Circulatory disturbances.

The circulatory disturbances, especially of the lower extremity following injury in the popliteal space, must not be overlooked, and require rating generally as phlebitis.

### §4.63 Loss of use of hand or foot.

Loss of use of a hand or a foot, for the purpose of special monthly compensation, will be held to exist when no effective function remains other than that which would be equally well served by an amputation stump at the site of election below elbow or knee with use of a suitable prosthetic appliance. The determination will be made on the basis of the actual remaining function of the hand or foot, whether the acts of grasping, manipulation, etc., in the case of the hand, or of balance and propulsion, etc., in the case of the foot, could be accomplished equally

well by an amputation stump with prosthesis.

(a) Extremely unfavorable complete ankylosis of the knee, or complete ankylosis of 2 major joints of an extremity, or shortening of the lower extremity of 3½ inches (8.9 cms.) or more, will be taken as loss of use of the hand or foot involved.

(b) Complete paralysis of the external popliteal nerve (common peroneal) and consequent, footdrop, accompanied by characteristic organic changes including trophic and circulatory disturbances and other concomitants confirmatory of complete paralysis of this nerve, will be taken as loss of use of the foot.

[29 FR 6718, May 22, 1964, as amended at 43 FR 45349, Oct. 2, 1978]

### §4.64 Loss of use of both buttocks.

Loss of use of both buttocks shall be deemed to exist when there is severe damage to muscle Group XVII, bilateral (diagnostic code number 5317) and additional disability rendering it impossible for the disabled person, without assistance, to rise from a seated position and from a stooped position (fingers to toes position) and to maintain postural stability (the pelvis upon head of femur). The assistance may be rendered by the person's own hands or arms, and, in the matter of postural stability, by a special appliance.

### §4.65 [Reserved]

### §4.66 Sacroiliac joint.

The common cause of disability in this region is arthritis, to be identified in the usual manner. The lumbosacral and sacroiliac joints should be considered as one anatomical segment for rating purposes. X-ray changes from arthritis in this location are decrease or obliteration of the joint space, with the appearance of increased bone density of the sacrum and ilium and sharpening of the margins of the joint. Disability is manifest from erector spinae spasm (not accounted for by other pathology), tenderness on deep palpation and percussion over these joints, loss of normal quickness of motion and resiliency, and postural defects often accompanied by limitation of flexion and extension of the hip. Traumatism is a

rare cause of disability in this connection, except when superimposed upon congenital defect or upon an existent arthritis; to permit assumption of pure traumatic origin, objective evidence of damage to the joint, and history of trauma sufficiently severe to injure this extremely strong and practically immovable joint is required. There should be careful consideration of lumbosacral sprain, and the various symptoms of pain and paralysis attributable to disease affecting the lumbar vertebrae and the intervertebral disc.

#### §4.67 Pelvic bones.

The variability of residuals following these fractures necessitates rating on specific residuals, faulty posture, limitation of motion, muscle injury, painful motion of the lumbar spine, manifest by muscle spasm, mild to moderate sciatic neuritis, peripheral nerve injury, or limitation of hip motion.

### §4.68 Amputation rule.

The combined rating for disabilities of an extremity shall not exceed the rating for the amputation at the elective level, were amputation to be performed. For example, the combined evaluations for disabilities below the knee shall not exceed the 40 percent evaluation, diagnostic code 5165. This 40 percent rating may be further combined with evaluation for disabilities above the knee but not to exceed the above the knee amputation elective level. Painful neuroma of a stump after amputation shall be assigned the evaluation for the elective site of reamputation.

### §4.69 Dominant hand.

Handedness for the purpose of a dominant rating will be determined by the evidence of record, or by testing on VA examination. Only one hand shall be considered dominant. The injured hand, or the most severely injured hand, of an ambidextrous individual will be considered the dominant hand for rating purposes.

(Authority: 38 U.S.C. 1155) [62 FR 30239, June 3, 1997]

### § 4.70

### §4.70 Inadequate examinations.

If the report of examination is inadequate as a basis for the required consideration of service connection and evaluation, the rating agency may request a supplementary report from the examiner giving further details as to the limitations of the disabled person's ordinary activity imposed by the disease, injury, or residual condition, the prognosis for return to, or continuance of, useful work. When the best interests of the service will be advanced by personal conference with the examiner, such conference may be arranged through channels.

# §4.71 Measurement of ankylosis and joint motion.

Plates I and II provide a standardized description of ankylosis and joint mo-

tion measurement. The anatomical position is considered as  $0^{\circ}$ , with two major exceptions: (a) Shoulder rotation-arm abducted to 90°, elbow flexed to 90° with the position of the forearm reflecting the midpoint 0° between internal and external rotation of the shoulder; and (b) supination and pronation—the arm next to the body, elbow flexed to 90°, and the forearm in midposition  $0^{\circ}$  between supination and pronation. Motion of the thumb and fingers should be described by appropriate reference to the joints (See Plate III) whose movement is limited, with a statement as to how near, in centimeters, the tip of the thumb can approximate the fingers, or how near the tips of the fingers can approximate the median transverse fold of the palm.




PLATE I




PLATE II

 $[29\;\mathrm{FR}\;6718,\,\mathrm{May}\;22,\,1964,\,\mathrm{as}\;\mathrm{amended}\;\mathrm{at}\;43\;\mathrm{FR}\;45349,\,\mathrm{Oct.}\;2,\,1978]$ 

# $\S4.71a$ Schedule of ratings—musculo-skeletal system.

ACUTE, SUBACUTE, OR CHRONIC DISEASES

	Rat- ing
5000 Osteomyelitis, acute, subacute, or chronic: Of the pelvis, vertebrae, or extending into major joints, or with multiple localization or with long history of intractability and debility, anemia, amyloid liver changes, or other continuous	
constitutional symptoms	100
Frequent episodes, with constitutional symptoms	60
With definite involucrum or sequestrum, with or without discharging sinus	30
tive infection within the past 5 years	20
Inactive, following repeated episodes, without evidence of active infection in past 5 years	10

ACUTE, SUBACUTE, OR CHRONIC DISEASES—Continued

	Rat- ing
Note (1): A rating of 10 percent, as an exception to the amputation rule, is to be assigned in any case of active osteomyelitis where the amputation rating for the affected part is no percent. This 10 percent rating and the other partial ratings of 30 percent or less are to be combined with ratings for ankylosis, limited motion, nonunion or malunion, shortening, etc., subject, of course, to the amputation rule. The 60 percent rating, as it is based on constitutional symptoms, is not subject to the amputation rule. A rating for osteomyelitis will not be applied following cure by removal or radical resection of the affected bone.	

## **Department of Veterans Affairs**

§ 4.71a

10

# ACUTE, SUBACUTE, OR CHRONIC DISEASES— Continued

Continued		7.00.2
	Rat- ing	-
NOTE (2): The 20 percent rating on the basis of		W
activity within the past 5 years is not assign- able following the initial infection of active os-		
teomyelitis with no subsequent reactivation.		
The prerequisite for this historical rating is an		W
established recurrent osteomyelitis. To qualify		
for the 10 percent rating, 2 or more episodes following the initial infection are required. This		Note
20 percent rating or the 10 percent rating,		on )
when applicable, will be assigned once only to		with
cover disability at all sites of previously active		Note
infection with a future ending date in the case of the 20 percent rating.		on >
5001 Bones and joints, tuberculosis of, active or in-		5013
active:		5004 Arth
Active	100	5005 Arth
Inactive: See §§ 4.88b and 4.895002 Arthritis rheumatoid (atrophic) As an active		5006 Arth
process:		5007 Arth 5008 Arth
With constitutional manifestations associated		5009 Arth
with active joint involvement, totally incapaci-	400	With th
tatingLess than criteria for 100% but with weight loss	100	thro
and anemia productive of severe impairment		arth
of health or severely incapacitating exacer-		5010 Arth ray findir
bations occurring 4 or more times a year or a		5011 Bon
lesser number over prolonged periods	60	cord inv
Symptom combinations productive of definite im- pairment of health objectively supported by ex-		severity
amination findings or incapacitating exacer-		5012 Bor Note:
bations occurring 3 or more times a year	40	for '
One or two exacerbations a year in a well-estab-	20	X-ra
lished diagnosis For chronic residuals:	20	ther
For residuals such as limitation of motion or an-		has the
kylosis, favorable or unfavorable, rate under		5013 Ost
the appropriate diagnostic codes for the spe-		5014 Ost
cific joints involved. Where, however, the limi- tation of motion of the specific joint or joints in-		5015 Bon
volved is noncompensable under the codes a		5016 Ost 5017 Gou
rating of 10 percent is for application for each		5017 Got 5018 Hyd
such major joint or group of minor joints af- fected by limitation of motion, to be combined,		5019 Bur
not added under diagnostic code 5002. Limita-		5020 Syn
tion of motion must be objectively confirmed		5021 Myd
by findings such as swelling, muscle spasm,		5022 Per 5023 Myo
or satisfactory evidence of painful motion.  Note: The ratings for the active process will not		5023 Myd 5024 Ten
be combined with the residual ratings for limi-		The o
tation of motion or ankylosis. Assign the high-		thro
er evaluation.		tion tive,
5003 Arthritis, degenerative (hypertrophic or osteo- arthritis):		agno
Degenerative arthritis established by X-ray find-		5025 Fibr
ings will be rated on the basis of limitation of		syndrom
motion under the appropriate diagnostic codes		With v
for the specific joint or joints involved (DC 5200 etc.). When however, the limitation of		der slee
motion of the specific joint or joints involved is		head
noncompensable under the appropriate diag-		sion
nostic codes, a rating of 10 pct is for applica-		Th
tion for each such major joint or group of minor joints affected by limitation of motion, to		TI
be combined, not added under diagnostic		"
code 5003. Limitation of motion must be ob-		
jectively confirmed by findings such as swell-		
ing, muscle spasm, or satisfactory evidence of painful motion. In the absence of limitation of		Th
motion, rate as below:		11
,		

# ACUTE, SUBACUTE, OR CHRONIC DISEASES—Continued

	Rat- ing
With X-ray evidence of involvement of 2 or	
more major joints or 2 or more minor joint	
groups, with occasional incapacitating ex-	00
acerbations With X-ray evidence of involvement of 2 or	20
more major joints or 2 or more minor joint	
groups	10
NOTE (1): The 20 pct and 10 pct ratings based	
on X-ray findings, above, will not be combined	
with ratings based on limitation of motion.  NOTE (2): The 20 pct and 10 pct ratings based	
on X-ray findings, above, will not be utilized in	
rating conditions listed under diagnostic codes	
5013 to 5024, inclusive.	
5004 Arthritis, gonorrheal.	
5005 Arthritis, pneumococcic. 5006 Arthritis, typhoid.	
5007 Arthritis, syphilitic.	
5008 Arthritis, streptococcic.	
5009 Arthritis, other types (specify).	
With the types of arthritis, diagnostic codes 5004	
through 5009, rate the disability as rheumatoid arthritis.	
5010 Arthritis, due to trauma, substantiated by X-	
ray findings: Rate as arthritis, degenerative.	
5011 Bones, caisson disease of: Rate as arthritis,	
cord involvement, or deafness, depending on the	
severity of disabling manifestations. 5012 Bones, new growths of, malignant	100
Note: The 100 percent rating will be continued	100
for 1 year following the cessation of surgical,	
X-ray, antineoplastic chemotherapy or other	
therapeutic procedure. At this point, if there has been no local recurrence or metastases,	
the rating will be made on residuals.	
5013 Osteoporosis, with joint manifestations.	
5014 Osteomalacia.	
5015 Bones, new growths of, benign.	
5016 Osteitis deformans. 5017 Gout.	
5017 Godt. 5018 Hydrarthrosis, intermittent.	
5019 Bursitis.	
5020 Synovitis.	
5021 Myositis.	
5022 Periostitis. 5023 Myositis ossificans.	
5024 Tenosynovitis.	
The diseases under diagnostic codes 5013	
through 5024 will be rated on limitation of mo-	
tion of affected parts, as arthritis, degenera-	
tive, except gout which will be rated under diagnostic code 5002.	
5025 Fibromyalgia (fibrositis, primary fibromyalgia	
syndrome)	
With widespread musculoskeletal pain and ten-	
der points, with or without associated fatigue, sleep disturbance, stiffness, paresthesias,	
headache, irritable bowel symptoms, depres-	
sion, anxiety, or Raynaud's-like symptoms:	
That are constant, or nearly so, and refrac-	
tory to therapy	40
That are episodic, with exacerbations often	
precipitated by environmental or emo- tional stress or by overexertion, but that	
are present more than one-third of the	
time	20
That require continuous medication for con-	10
not	

# § 4.71a

# ACUTE, SUBACUTE, OR CHRONIC DISEASES—Continued

	Rat- ing
Note: Widespread pain means pain in both the left and right sides of the body, that is both above and below the waist, and that affects both the axial skeleton (i.e., cervical spine, anterior chest, thoracic spine, or low back) and the extremities.	

## PROSTHETIC IMPLANTS

	Rati	ing
	Major	Minor
5051 Shoulder replacement (prosthesis).  Prosthetic replacement of the shoulder joint:		
For 1 year following implantation of prosthesis	100	100
ness in the affected extremity With intermediate degrees of residual weakness, pain or limitation of motion, rate by analogy to di-	60	50
agnostic codes 5200 and 5203.  Minimum rating	30	20
For 1 year following implantation of prosthesis	100	100
severe painful motion or weak- ness in the affected extremity With intermediate degrees of resid- ual weakness, pain or limitation of motion rate by analogy to di- agnostic codes 5205 through 5208.	50	40
Minimum evaluation	30	20
prosthesis	100	100
ness in the affected extremity With intermediate degrees of residual weakness, pain or limitation of motion, rate by analogy to diagnostic code 5214.	40	30
Minimum rating	20	20
prosthesis		100

# PROSTHETIC IMPLANTS—Continued

	Rat	ing
	Major	Mino
Following implantation of prosthesis		
with painful motion or weakness		
such as to require the use of		
crutches		1 90
Markedly severe residual weak-		
ness, pain or limitation of motion		
following implantation of pros-		
thesis		70
Moderately severe residuals of		
weakness, pain or limitation of motion		50
Minimum rating		30
5055 Knee replacement (prosthesis).		"
Prosthetic replacement of knee joint:		İ
For 1 year following implantation of		
prosthesis		100
With chronic residuals consisting of		
severe painful motion or weak-		
ness in the affected extremity		60
With intermediate degrees of resid-		
ual weakness, pain or limitation		
of motion rate by analogy to di-		
agnostic codes 5256, 5261, or		
5262.		
Minimum rating5056 Ankle replacement (prosthesis).		3
5056 Ankle replacement (prosthesis).  Prosthetic replacement of ankle joint:		
For 1 year following implantation of		
prosthesis		10
With chronic residuals consisting of		
severe painful motion or weak-		
ness		4
With intermediate degrees of resid-		
ual weakness, pain or limitation		
of motion rate by analogy to		
5270 or 5271.		
Minimum rating		20
NOTE (1): The 100 pct rating for 1 year		
following implantation of prosthesis		
will commence after initial grant of the 1-month total rating assigned under		
§ 4.30 following hospital discharge.		
NOTE (2): Special monthly compensa-		
tion is assignable during the 100 pct		
rating period the earliest date perma-		
nent use of crutches is established.		
COMBINATIONS OF DISABILITIES		
5104 Anatomical loss of one hand and loss		
of use of one foot		1100
5105 Anatomical loss of one foot and loss		'0'
of use of one hand		110
5106 Anatomical loss of both hands		1100
5107 Anatomical loss of both feet		110
5108 Anatomical loss of one hand and one		
foot		1100
5109 Loss of use of both hands		1100
		110
5110 Loss of use of both feet		
5110 Loss of use of both feet 5111 Loss of use of one hand and one foot		110

<sup>&</sup>lt;sup>1</sup> Also entitled to special monthly compensation.

TABLE II—RATINGS FOR MULTIPLE LOSSES OF EXTREMITIES WITH DICTATOR'S RATING CODE AND 38 CFR CITATION

			Impairment of other extremity	ther extremity		
Impairment of one extremity	Anatomical loss or loss of use below elbow	Anatomical loss or loss of use below knee	Anatomical loss or loss of use above elbow (preventing use of prosthesis)	Anatomical loss or loss of use above knee (preventing use of prosthesis)	Anatomical loss near shoulder (preventing use of prosthesis)	Anatomical loss near hip (preventing use of prosthesis)
Anatomical loss or loss of use below elbow.	M Codes M-1 a, b, or c, L Codes L-1 d, e, f, or 38 CFR 3.350 (c)(1)(i). g, 38 CFR 3.350(b).	L Codes L-1 d, e, f, or g, 38 CFR 3.350(b).	M½ Code M-5, 38 CFR   L½ Code L-2 c, 38 3.350 (f)(1)(x).   CFR 3.350 (f)(1)(v	L½ Code L-2 c, 38 CFR 3.350 (f)(1)(vi).	N Code N-3, 38 CFR 3.350 (f)(1)(xi).	M Code M-3 c, 38 CFR 3.350 (f)(1)(viii)
Anatomical loss or loss of use below knee.		_	L½ Code L–2 b, 38 CFR 3.350 (f)(1)(iii).	L½ Code L-2 a, 38 CFR 3.350 (f)(1)(i).	M Code M-3 b, 38 CFR 3.350 (f)(1)(iv).	M Code M-3 a, 38 CFR 3.350 (f)(1)(ii)
Anatomical loss or loss of use above elbow (preventing use of prosembles)			N Code N-1, 38 CFR 3.350 (d)(1).	M Code M-2 a, 38 CFR 3.350 (c)(1)(iii).	N½ Code N-4, 38 CFR 3.350 (f)(1)(ix).	M½ Code M-4 c, 38 CFR 3.350 (f)(1)(xi)
Anatomical loss or loss of use above knee (preventing use of pros-				M Code M-2 a, 38 CFR 3.350 (c)(1)(ii).	M½ Code M-4 b, 38 CFR 3.350 (f)(1)(vii).	M½ Code M–4 a, 38 CFR 3.350 (f)(1)(v)
Anatomical loss near shoulder (preventing use of prosthesis)					O Code O-1, 38 CFR 3.350 (e)(1)(i).	N Code N-2 b, 38 CFR 3.350 (d)(3)
Anatomical loss near hip (preventing use of prosthesis).						N Code N–2 a, 38 CFR 3.350 (d)(2)

NOTE.—Need for aid attendance or permanently bedridden qualifies for subpar. L. Code L–1 h, i (38 CFR 3.350(b)). Paraplegia with loss of use of both lower extremities and loss of anal and bladder sphincter control qualifies for subpar. O. Code O–2 (38 CFR 3.350(e)(2)). Where there are additional disabilities rated 50% or 100%, or anatomical or loss of use of a third extremity see 38 CFR 3.350(f) (3), (4) or (5).

(Authority: 38 U.S.C. 1115)

# § 4.71a

## AMPUTATIONS: UPPER EXTREMITY

		Rating	
		Major	Minor
А	rm, amputation of:		
5120	Disarticulation	190	190
5121	Above insertion of deltoid	190	180
5122	Below insertion of deltoid	180	170
F	orearm, amputation of:		
5123	Above insertion of pronator teres	180	170
5124	Below insertion of pronator teres	170	1 60
5125	Hand, loss of use of	170	1 60

	MULTIPLE FINGER AMPUTATIONS		
	Five digits of one hand, amputation	170	1 60
	our digits of one hand, amputation of:	. 70	. 00
5127	Thumb, index, middle and ring	170	1 60
5128	Thumb, index, middle and little	170	160
5128	Thumb, index, ring and little	170	1 60
5130	Thumb, middle, ring and little	170	1 60
5131	Index, middle, ring and little	60	50
	hree digits of one hand, amputation of:	00	30
5132	Thumb, index and middle	60	50
5133	Thumb, index and ring	60	50
5134	Thumb, index and little	60	50
5135	Thumb, middle and ring	60	50
5136	Thumb, middle and little	60	50
5137	Thumb, ring and little	60	50
5138	Index, middle and ring	50	40
5139	Index, middle and little	50	40
5140	Index, ring and little	50	40
5141	Middle, ring and little	40	30
	wo digits of one hand, amputation of:	10	00
5142	Thumb and index	50	40
5143	Thumb and middle	50	40
5144	Thumb and ring	50	40
5145	Thumb and little	50	40
5146	Index and middle	40	30
5147	Index and ring	40	30
5148	Index and little	40	30
5149	Middle and ring	30	20
5150	Middle and little	30	20
5151	Ring and little	30	20
(a	a) The ratings for multiple finger ampu-		
	tations apply to amputations at the		
	proximal interphalangeal joints or		
	through proximal phalanges		
(b	) Amputation through middle pha-		
	langes will be rated as prescribed for		
10	unfavorable ankylosis of the fingers c) Amputations at distal joints, or		
(0	through distal phalanges, other than		
	negligible losses, will be rated as pre-		
	scribed for favorable ankylosis of the		
	fingers		

## AMPUTATIONS: UPPER EXTREMITY—Continued

	Rating		
	Major	Minor	
(d) Amputation or resection of meta- carpal bones (more than one-half the bone lost) in multiple fingers injuries will require a rating of 10 percent added to (not combined with) the rat- ings, multiple finger amputations, sub- ject to the amputation rule applied to the forearm. (e) Combinations of finger amputations at various levels, or finger amputa- tions with ankylosis or limitation of motion of the fingers will be rated on the basis of the grade of disability; i.e., amputation, unfavorable anky- losis, most representative of the lev- els or combinations. With an even number of fingers involved, and adja- cent grades of disability, select the higher of the two grades. (f) Loss of use of the hand will be held to exist when no effective function re- mains other than that which would be equally well served by an amputation			
stump with a suitable prosthetic applicance.  SINGLE FINGER AMPUTATIONS			
5152 Thumb, amputation of: With metacarpal resection At metacarpophalangeal joint or through	40	30	
proximal phalanx  At distal joint or through distal phalanx 5153 Index finger, amputation of  With metacarpal resection (more than	30 20	20	
one-half the bone lost)	30	20	
thereto	20 10	10	
one-half the bone lost)	20	20	
thereto thereto finger, amputation of:  With metacarpal resection (more than one-half the bone lost)	10	20	
Without metacarpal resection, at proximal interphalangeal joint or proximal thereto	10	1(	
5156 Little finger, amputation of: With metacarpal resection (more than one-half the bone lost) Without metacarpal resection, at proxi-	20	20	
mal interphalangeal joint or proximal thereto	10	10	

<sup>&</sup>lt;sup>1</sup> Entitled to special monthly compensation.

### SINGLE FINGER AMPUTATIONS




PLATE III

# 38 CFR Ch. I (7-1-99 Edition)

# § 4.71a

# AMPUTATIONS: LOWER EXTREMITY

	Rat- ing
Thigh, amputation of:	
5160 Disarticulation, with loss of extrinsic pelvic gir-	
dle muscles	290
5161 Upper third, one-third of the distance from	
perineum to knee joint measured from perineum	<sup>2</sup> 80
5162 Middle or lower thirds	<sup>2</sup> 60
Leg, amputation of:	
5163 With defective stump, thigh amputation rec-	
ommended	<sup>2</sup> 60
5164 Amputation not improvable by prosthesis con-	
trolled by natural knee action	<sup>2</sup> 60
5165 At a lower level, permitting prosthesis	240
5166 Forefoot, amputation proximal to metatarsal	
bones (more than one-half of metatarsal loss)	240
5167 Foot, loss of use of	2 40

# AMPUTATIONS: LOWER EXTREMITY—Continued

	Rat- ing
5170 Toes, all, amputation of, without metatarsal	
loss	30
5171 Toe, great, amputation of:	
With removal of metatarsal head	30
Without metatarsal involvement	10
5172 Toes, other than great, amputation of, with re-	
moval of metatarsal head:	
One or two	20
Without metatarsal involvement	0
5173 Toes, three or four, amputation of, without	
metatarsal involvement:	
Including great toe	20
Not including great toe	10

<sup>&</sup>lt;sup>2</sup> Also entitled to special monthly compensation.

### AMPUTATIONS: LOWER EXTREMITY




PLATE IV

# § 4.71a

# THE SHOULDER AND ARM

# THE ELBOW AND FOREARM—Continued

	Rati	Rating	
	Major	Minor	
5200 Scapulohumeral articulation, anky-			
losis of:			
Note: The scapula and humerus move as one piece.			
Unfavorable, abduction limited to 25°			
from side	50	40	
Intermediate between favorable and un-			
favorable	40	30	
Favorable, abduction to 60°, can reach			
mouth and head	30	20	
5201 Arm, limitation of motion of:			
To 25° from side	40	30	
Midway between side and shoulder			
level	30	20	
At shoulder level	20	20	
5202 Humerus, other impairment of:			
Loss of head of (flail shoulder)	80	70	
Nonunion of (false flail joint)	60	50	
Fibrous union of	50	40	
Recurrent dislocation of at			
scapulohumeral joint.			
With frequent episodes and guard-			
ing of all arm movements	30	20	
With infrequent episodes, and			
guarding of movement only at			
shoulder level	20	20	
Malunion of:			
Marked deformity	30	20	
Moderate deformity	20	20	
5203 Clavicle or scapula, impairment of:			
Dislocation of	20	20	
Nonunion of:			
With loose movement	20	20	
Without loose movement	10	10	
Malunion of	10	10	
Or rate on impairment of function of			
contiguous joint.			

# THE ELBOW AND FOREARM

	Rating	
	Major	Minor
5205 Elbow, ankylosis of:		
Unfavorable, at an angle of less than		
50° or with complete loss of		
supination or pronation	60	50
Intermediate, at an angle of more than		
90°, or between 70° and 50°	50	40
Favorable, at an angle between 90° and		
70°	40	30
5206 Forearm, limitation of flexion of:		
Flexion limited to 45°	50	40
Flexion limited to 55°	40	30
Flexion limited to 70°	30	20
Flexion limited to 90°	20	20
Flexion limited to 100°	10	10
Flexion limited to 110°	0	0
5207 Forearm, limitation of extension of:		
Extension limited to 110°	50	40
Extension limited to 100°	40	30
Extension limited to 90°	30	20
Extension limited to 75°	20	20
Extension limited to 60°	10	10
Extension limited to 45°	10	10
5208 Forearm, flexion limited to 100° and		
extension to 45°	20	20
5209 Elbow, other impairment of Flail joint	60	50

	Rati	Rating	
	Major	Minor	
Joint fracture, with marked cubitus varus or cubitus valgus deformity or with ununited fracture of head of radius	20	20	
5210 Radius and ulna, nonunion of, with	50	40	
flail false joint	50	40	
Nonunion in upper half, with false movement:			
With loss of bone substance (1 inch (2.5 cms.) or more) and marked			
deformity Without loss of bone substance or	40	30	
deformity	30	20	
Nonunion in lower half	20	20	
Malunion of, with bad alignment 5212 Radius, impairment of:	10	10	
Nonunion in lower half, with false movement:			
With loss of bone substance (1 inch			
(2.5 cms.) or more) and marked			
deformity	40	30	
Without loss of bone substance or			
deformity	30	20	
Nonunion in upper half	20	20	
Malunion of, with bad alignment	10	10	
5213 Supination and pronation, impairment of:			
Loss of (bone fusion):			
The hand fixed in supination or			
hyperpronation	40	30	
The hand fixed in full pronation	30	20	
The hand fixed near the middle of			
the arc or moderate pronation	20	20	
Limitation of pronation:			
Motion lost beyond middle of arc	30	20	
Motion lost beyond last quarter of			
arc, the hand does not approach			
full pronation	20	20	
Limitation of supination:			
To 30° or less	10	10	
NOTE: In all the forearm and wrist inju-			
ries, codes 5205 through 5213, mul-			
tiple impaired finger movements due			
to tendon tie-up, muscle or nerve in-			
jury, are to be separately rated and			
combined not to exceed rating for			
loss of use of hand.			

# THE WRIST

	Rating	
	Major	Minor
5214 Wrist, ankylosis of:     Unfavorable, in any degree of palmar flexion, or with ulnar or radial deviation	50 40 30	40 30 20
5215 Wrist, limitation of motion of:  Dorsiflexion less than 15°  Palmar flexion limited in line with fore-	10	10
arm	10	10

## MULTIPLE FINGERS: UNFAVORABLE ANKYLOSIS

Rating Major Minor

In classifying the severity of ankylosis and limitation of motion of single dig- its and combinations of digits the fol-		
lowing rules will be observed:		
(1) Ankylosis of both the		
metacarpophalangeal and proxi-		
mal interphalangeal joints, with		
either joint in extension or in ex-		
treme flexion, will be rated as		
amputation.		
(2) Ankylosis of both the		
metacarpophalangeal and proxi-		
mal interphalangeal joints, even		
though each is individually in fa-		
vorable position, will be rated as		
unfavorable ankylosis.		
(3) With only one joint of a digit		
ankylosed or limited in its motion, the determination will be made		
on the basis of whether motion is		
possible to within 2 inches (5.1		
cms.) of the median transverse		
fold of the palm; when so pos-		
sible, the rating will be for favor-		
able ankylosis, otherwise unfa-		
vorable.		
(4) With the thumb, the		
carpometacarpal joint is to be re-		
garded as comparable to the metacarpophalangeal joint of		
other digits.		
5216 Five digits of one hand, unfavorable		
ankylosis of	60	50
5217 Four digits of one hand, unfavorable		
ankylosis of:		
Thumb, index, middle and ring	60	50
Thumb, index, middle and little	60	50
Thumb, index, ring and little	60	50
Thumb, middle, ring and little	60	50
Index, middle, ring and little	50	40
5218 Three digits of one hand, unfavorable		
ankylosis of:		
Thumb, index and middle	50	40
Thumb, index and ring	50	40
Thumb, index and little	50	40
Thumb, middle and ring	50	40
Thumb, middle and little	50	40
Thumb, ring and little	50	40
Index, middle and ring	40	30
Index, middle and little	40	30
Index, ring and little	40	30
Middle, ring and little	30	20
ankylosis of:		
Thumb and index	40	30
Thumb and middle	40	30

Thumb and middle .....

Thumb and ring .....

Middle and ring ..... Middle and little ..... Ring and little .....

Thumb and little Index and middle ..... Index and ring ..... Index and little .....

### MULTIPLE FINGERS: UNFAVORABLE ANKYLOSIS— Continued

	Rati	ing
	Major	Minor
<ul> <li>(a) Extremely unfavorable ankylosis of the fingers, all joints in extension or in extreme flexion, or with rotation and angulation of bones, will be rated as amputation.</li> <li>(b) The ratings for codes 5216 through 5219 apply to unfavorable ankylosis or limited motion preventing flexion of tips to within 2 inches (5.1 cms.) of median transverse fold of the palm.</li> <li>(c) Combinations of finger amputations at various levels, or of finger amputations with ankylosis or limitation of motion of the fingers will be rated on the basis of the grade of disability, i.e., amputation, unfavorable ankylosis, or favorable ankylosis, most representative of the levels or combinations. With an even number of fingers involved, and adjacent grades of disability, select the higher of the two grades.</li> </ul>		

## MULTIPLE FINGERS: FAVORABLE ANKYLOSIS

	Rating	
	Major	Minor
In classifying the severity of ankylosis and limitation of motion of single digits and combinations of digits the following rules will be observed:  (1) Ankylosis of both the metacarpophalangeal and proximal interphalangeal joints, with either joint in extension or in extreme flexion, will be rated as amputation.  (2) Ankylosis of both the metacarpophalangeal and proximal interphalangeal joints, even though each is individually in favorable position, will be rated as unfavorable ankylosis.  (3) With only one joint of a digit ankylosed or limited in its motion, the determination will be made on the basis of whether motion is possible to within 2 inches (5.1 cms.) of the median transverse fold of the palm; when so possible, the rating will be for favorable ankylosis, otherwise unfavorable.  (4) With the thumb, the carpometacarpal joint is to be regarded as comparable to the metacarpophalangeal joint of other digits.		
5220 Five digits of one hand, favorable ankylosis of	50	40
ankylosis of: Thumb, index, middle and ring	50	41
Thumb, index, middle and little	50	

40

30

# § 4.71a

# MULTIPLE FINGERS: FAVORABLE ANKYLOSIS—Continued

	Rating	
	Major	Minor
Index, middle, ring and little5222 Three digits of one hand, favorable ankylosis of:	40	30
Thumb, index and middle	40	30
Thumb, index and ring	40	30
Thumb, index and little	40	30
Thumb, middle and ring	40	30
Thumb, middle and little	40	30
Thumb, ring and little	40	30
Index, middle and ring	30	20
Index, middle and little	30	20
Index, ring and little	30	20
Middle, ring and little	20	20
5223 Two digits of one hand, favorable an-		
kylosis of:		
Thumb and index	30	20
Thumb and middle	30	20
Thumb and ring	30	20
Thumb and little	30	20
Index and middle	20	20
Index and ring	20	20
Index and little	20	20
Middle and ring	10	10
Middle and little	10	10
	10	10
Ring and little	10	10

### ANKYLOSIS OF INDIVIDUAL FINGERS

	Rati	ing
	Major	Minor
5224 Thumb, ankylosis of:		
Unfavorable	20	20
Favorable	10	10
5225 Index finger, ankylosis of:		
Unfavorable	10	10
Favorable	10	10
5226 Middle finger, ankylosis of:		
Unfavorable	10	10
Favorable	10	10
5227 Finger, any other, ankylosis of	0	0
NOTE: Extremely unfavorable ankylosis will be rated as amputation under di- agnostic codes 5152 through 5156.		

# THE HIP AND THIGH

	Rat- ing
5250 Hip, ankylosis of:	
Unfavorable, extremely unfavorable ankylosis,	
the foot not reaching ground, crutches neces-	
sitated	₃90
Intermediate	70
Favorable, in flexion at an angle between 20°	
and 40°, and slight adduction or abduction	60
5251 Thigh, limitation of extension of:	
Extension limited to 5°	10
5252 Thigh, limitation of flexion of:	
Flexion limited to 10°	40 30
Flexion limited to 20°	20
Flexion limited to 30°	10
5253 Thigh, impairment of:	10
Limitation of abduction of, motion lost beyond	
10°	20
Limitation of adduction of, cannot cross legs	10
Limitation of rotation of, cannot toe-out more	
than 15°, affected leg	10
5254 Hip, flail joint	80
5255 Femur, impairment of:	
Fracture of shaft or anatomical neck of:	
With nonunion, with loose motion (spiral or	
oblique fracture)	80
With nonunion, without loose motion,	
weightbearing preserved with aid of brace	60
Fracture of surgical neck of, with false joint	60
Malunion of:	
With marked knee or hip disability	30
With moderate knee or hip disability	20
With slight knee or hip disability	10

<sup>&</sup>lt;sup>3</sup> Entitled to special monthly compensation.

## THE KNEE AND LEG

	Rat- ing
5256 Knee, ankylosis of:	
Extremely unfavorable, in flexion at an angle of	
45° or more	60
In flexion between 20° and 45°	50
In flexion between 10° and 20°	40
Favorable angle in full extension, or in slight	
flexion between 0° and 10°	30
5257 Knee, other impairment of:	
Recurrent subluxation or lateral instability:	
Severe	30
Moderate	20
Slight	10
5258 Cartilage, semilunar, dislocated, with frequent	
episodes of "locking," pain, and effusion into the	
joint	20
5259 Cartilage, semilunar, removal of, symptomatic	10
5260 Leg, limitation of flexion of:	
Flexion limited to 15°	30
Flexion limited to 30°	20
Flexion limited to 45°	10
Flexion limited to 60°	0
5261 Leg, limitation of extension of:	
Extension limited to 45°	50
Extension limited to 30°	40
Extension limited to 20°	30
Extension limited to 15°	20
Extension limited to 10°	10
Extension limited to 5°	0
5262 Tibia and fibula, impairment of:	
Nonunion of, with loose motion, requiring brace	40
Malunion of:	
With marked knee or ankle disability	30

# **Department of Veterans Affairs**

# § 4.71a

# THE KNEE AND LEG—Continued

	Rat- ing
With moderate knee or ankle disability	20
With slight knee or ankle disability	10
tively demonstrated)	10

## THE ANKLE

	Rat- ing
5270 Ankle, ankylosis of:	
In plantar flexion at more than 40°, or in	
dorsiflexion at more than 10° or with abduc-	
tion, adduction, inversion or eversion deformity	40
In plantar flexion, between 30° and 40°, or in	
dorsiflexion, between 0° and 10°	30
In plantar flexion, less than 30°	20
5271 Ankle, limited motion of:	
Marked	20
Moderate	10
5272 Subastragalar or tarsal joint, ankylosis of:	
In poor weight-bearing position	20
In good weight-bearing position	10
5273 Os calcis or astragalus, malunion of:	
Marked deformity	20
Moderate deformity	10
5274 Astragalectomy	20

# SHORTENING OF THE LOWER EXTREMITY

	Rat- ing
5275 Bones, of the lower extremity, shortening of: Over 4 inches (10.2 cms.)	<sup>3</sup> 60 <sup>3</sup> 50 40 30 20 10

<sup>&</sup>lt;sup>3</sup> Also entitled to special monthly compensation.

### THE FOOT

	Rat- ing
5276 Flatfoot, acquired: Pronounced; marked pronation, extreme tenderness of plantar surfaces of the feet, marked inward displacement and severe spasm of the tendo achillis on manipulation, not improved by orthopedic shoes or appliances. Bilateral	50 30
Bilateral	30
Unilateral	20

# THE FOOT—Continued

	Rat- ing
Moderate; weight-bearing line over or medial to great toe, inward bowing of the tendo achillis, pain on manipulation and use of the feet, bilat-	
eral or unilateral	10
Mild; symptoms relieved by built-up shoe or arch support	(
5277 Weak foot, bilateral:  A symptomatic condition secondary to many constitutional conditions, characterized by atrophy of the musculature, disturbed circulation, and weakness:  Rate the underlying condition, minimum rat-	10
ing	10
Bilateral Unilateral All toes tending to dorsiflexion, limitation of dorsiflexion at ankle to right angle, shortened plantar fascia, and marked tenderness under metatarsal heads:	50 30
Bilateral  Unilateral  Great toe dorsiflexed, some limitation of dorsiflexion at ankle, definite tenderness under metatarsal heads:	30 20
Bilateral	10
Unilateral  Slight 5279 Metatarsalgia, anterior (Morton's disease),	10
unilateral, or bilateral	10
Operated with resection of metatarsal head Severe, if equivalent to amputation of great toe  5281 Hallux rigidus, unilateral, severe: Rate as hallux valgus, severe. Note: Not to be combined with claw foot ratinos.	1( 1)
5282 Hammer toe: All toes, unilateral without claw foot	10
nonunion of: Severe	3( 2) 1(
5284 Foot injuries, other: Severe Moderately severe Moderate NoTE: With actual loss of use of the foot, rate 40 percent.	30 20 10

## THE SPINE

	Rat- ing
5285 Vertebra, fracture of, residuals: With cord involvement, bedridden, or requiring long leg braces	100
Without cord involvement; abnormal mobility requiring neck brace (jury mast)	60

## 38 CFR Ch. I (7-1-99 Edition)

### THE SPINE—Continued

	Rat- ing
In other cases rate in accordance with definite limited motion or muscle spasm, adding 10 percent for demonstrable deformity of	
vertebral body.  NOTE: Both under ankylosis and limited motion,	
ratings should not be assigned for more than one segment by reason of involvement of only the first or last vertebrae of an adjacent seg- ment.	
5286 Spine, complete bony fixation (ankylosis) of: Unfavorable angle, with marked deformity and involvement of major joints (Marie-Strumpell	
type) or without other joint involvement (Bechterew type)	100 60
5287 Spine, ankylosis of, cervical: Unfavorable	40
Favorable	30
Favorable	20
Unfavorable Favorable	50 40
5290 Spine, limitation of motion of, cervical: Severe	30 20
Slight	10
Severe	10 10
Slight	0 40
ModerateSlight	20 10
5293 Intervertebral disc syndrome: Pronounced; with persistent symptoms compatible with sciatic neuropathy with characteristic pain and demonstrable muscle spasm, absent ankle jerk, or other neurological findings appropriate to site of diseased disc, little intermittent relief	60
Severe; recurring attacks, with intermittent relief Moderate; recurring attacks Mild Postoperative, cured	40 20 10 0
5294 Sacro-iliac injury and weakness: 5295 Lumbosacral strain: Severe; with listing of whole spine to opposite side, positive Goldthwaite's sign, marked limi- tation of forward bending in standing position, loss of lateral motion with osteo-arthritic changes, or narrowing or irregularity of joint	
space, or some of the above with abnormal mobility on forced motion	40
standing position	20 10
With characteristic pain on motion With slight subjective symptoms only	0

### THE SKULL

		Rat- ing
5206	Skull loss of part of both inner and outer ta-	

### THE SKULL—Continued

	Rat- ing
With brain hernia	80
Without brain hernia:	
Area larger than size of a 50-cent piece or	
1.140 in <sup>2</sup> (7.355 cm <sup>2</sup> )	50
Area intermediate	30
Area smaller than the size of a 25-cent piece or 0.716 in 2 (4.619 cm <sup>2</sup> )	10
Note: Rate separately for intracranial complications.	

### THE RIBS

	Rat- ing
5297 Ribs, removal of:	
More than six	50
Five or six	40
Three or four	30
Two	20
One or resection of two or more ribs without re-	
generation	10
NOTE (1): The rating for rib resection or removal is not to be applied with ratings for purrulent pleurisy, lobectomy, pneumonectomy or injuries of pleural cavity.	
NOTE (2): However, rib resection will be considered as rib removal in thoracoplasty performed for collapse therapy or to accomplish obliteration of space and will be combined with the rating for lung collapse, or with the rating for lobectomy, pneumonectomy or the graduated ratings for pulmonary tuberculosis.	

### THE COCCYX

	Rat- ing
5298 Coccyx, removal of:	
Partial or complete, with painful residuals	
Without painful residuals	0

[29 FR 6718, May 22, 1964, as amended at 34 FR 5062, Mar. 11, 1969; 40 FR 42536, Sept. 15, 1975; 41 FR 11294, Mar. 18, 1976; 43 FR 45350, Oct. 2, 1978; 51 FR 6411, Feb. 24, 1986; 61 FR 20439, May 7, 1996]

## §4.72 [Reserved]

# §4.73 Schedule of ratings—muscle injuries.

Note: When evaluating any claim involving muscle injuries resulting in loss of use of any extremity or loss of use of both buttocks (diagnostic code 5317, Muscle Group XVII), refer to §3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation.

# THE SHOULDER GIRDLE AND ARM

# THE FOREARM AND HAND

	Rating			Rat	
	Domi- nant	Non- domi- nant		Domi- nant	Non- domi- nant
5301 Group I. Function: Upward rotation of scapula; elevation of arm above shoulder level. Extrinsic muscles of shoulder girdle: (1) Trapezius; (2) levator scapulae; (3) serratus magnus.			5307 Group VII. Function: Flexion of wrist and fingers. Muscles arising from internal condyle of humerus: Flexors of the carpus and long flexors of fingers and thumb; pronator.		
Severe	40	30	Severe	40	30
Moderately Severe	30	20	Moderately Severe	30	20
Moderate	10	10	Moderate	10	10
Slight	0	0	Slight	0	(
5302 Group II. Function: Depression of arm from vertical overhead to hanging at side (1, 2); downward rotation of scapula (3, 4); 1 and 2 act with Group III in forward and backward swing of arm. Extrin-			5308 Group VIII. Function: Extension of wrist, fingers, and thumb; abduction of thumb. Muscles arising mainly from external condyle of humerus: Extensors of carpus, fingers, and thumb; supinator.		
sic muscles of shoulder girdle: (1) Pectoralis major II (costosternal); (2)			Severe	30	20
Pectoralis major II (costosternal); (2) latissimus dorsi and teres major (teres			Moderately Severe	20	20
major, although technically an intrinsic			Moderate Slight	10	10
muscle, is included with latissimus dorsi);			5309 Group IX. Function: The forearm	"	
(3) pectoralis minor; (4) rhomboid.			muscles act in strong grasping move-		
Severe	40	30	ments and are supplemented by the intrin-		
Moderately Severe	30	20	sic muscles in delicate manipulative		
Moderate	20	20	movements. <i>Intrinsic muscles of hand:</i> Thenar eminence; short flexor, opponens,		
Slight	0	0	abductor and adductor of thumb; hypothenar eminence; short flexor, opponens and abductor of little finger; 4 lumbricales; 4 dorsal and 3 palmar interossei.		
of shoulder girdle: (1) Pectoralis major I (clavicular); (2) deltoid. Severe	40	30	NOTE: The hand is so compact a structure that isolated muscle injuries are rare,		
Moderately Severe	30	20	being nearly always complicated with inju-		
Moderate	20	20	ries of bones, joints, tendons, etc. Rate on		
Slight	0	0	limitation of motion, minimum 10 percent.		
5304 Group IV. Function: Stabilization of shoulder against injury in strong move- ments, holding head of humerus in socket; abduction: outward rotation and inward ro-			THE FOOT AND LEG		
tation of arm. Intrinsic muscles of shoulder girdle: (1) Supraspinatus; (2) infraspinatus					Rat- ing
and teres minor; (3) subscapularis; (4) coracobrachialis.			5310 Group X. Function: Movements of and toes; propulsion thrust in walking.		
Severe	30	20	muscles of the foot: Plantar: (1) Flexor di		
Moderate	20 10	20 10	brevis; (2) abductor hallucis; (3) abductor		
Moderate Slight	0	0	minimi; (4) quadratus plantae; (5) lumbrica flexor hallucis brevis; (7) adductor hallucis;		
5305 Group V. Function: Elbow supination			or digiti minimi brevis; (9) dorsal and		
(1) (long head of biceps is stabilizer of			interossei. Other important plantar structure		
shoulder joint); flexion of elbow (1, 2, 3).			tar aponeurosis, long plantar	and	
Flexor muscles of elbow: (1) Biceps; (2)			calcaneonavicular ligament, tendons of p tibial, peroneus longus, and long flexors of	osterior	
brachialis; (3) brachioradialis.	40	30	and little toes.	Ji great	
Severe  Moderately Severe	30	20	Severe		30
Moderate	10	10	Moderately Severe		20
Slight	0	0	Moderate		10
5306 Group VI. Function: Extension of elbow (long head of triceps is stabilizer of shoulder joint). Extensor muscles of the elbow: (1) Triceps; (2) anconeus		-	Slight	extensor uctures:	C
Severe	40	30	dons of long extensors of toes and peron		
Moderately Severe	30	20	cles.		
Moderate	10	10	Severe		20
Slight	0	0	Moderately Severe		10
-	l	<u> </u>	Moderate Slight		10

# § 4.73

# THE FOOT AND LEG—Continued

	Rat- ing
NOTE: Minimum rating for through-and-through wounds of the foot—10.	
5311 Group XI. Function: Propulsion, plantar flexion of foot (1); stabilization of arch (2, 3); flexion of toes (4, 5); Flexion of knee (6). Posterior and lateral crural muscles, and muscles of the calf: (1) Triceps surae (gastrocnemius and soleus); (2) tibialis posterior; (3) peroneus longus; (4) peroneus brevis; (5) flexor hallucis longus; (6) flexor digitorum longus; (7) popliteus; (8) plantaris.  Severe	30 20 10 0
Severe	30
Moderately Severe	20
Moderate Slight	10 0

### THE PELVIC GIRDLE AND THIGH

THE PELVIC GIRDLE AND THIGH	
	Rat- ing
5313 Group XIII. Function: Extension of hip and flexion of knee; outward and inward rotation of flexed knee; acting with rectus femoris and sartorius (see XIV, 1, 2) synchronizing simultaneous flexion of hip and knee and extension of hip and knee by belt-over-pulley action at knee joint. Posterior thigh group, Hamstring complex of 2-joint muscles: (1) Biceps femoris; (2) semimembranosus; (3) semitendinosus.  Severe	40 30 10 0
Severe	40 30 10 0
Severe	30 20 10 0
Severe  Moderately Severe  Moderate  Slight	40 30 10 0

# THE PELVIC GIRDLE AND THIGH—Continued

	Rat- ing
5317 Group XVII. Function: Extension of hip (1); abduction of thigh; elevation of opposite side of pelvis (2, 3); tension of fascia lata and ilitotibial (Maissiat's) band, acting with XIV (6) in postural support of body steadying pelvis upon head of femur and condyles of femur on tibia (1). Pelvic girdle group 2: (1) Gluteus maximus; (2) gluteus medius; (3) gluteus minimus.  Severe Moderately Severe Moderately Severe Siight Siight 5318 Group XVIII. Function: Outward rotation of thigh and stabilization of hip joint. Pelvic girdle group 3: (1) Pyriformis; (2) gemellus (superior or inferior); (3) obturator (external or internal); (4) quadratus femoris.	*50 40 20 0
Severe	30
Moderately Severe	20
Moderate	10
Slight	0

<sup>\*</sup>If bilateral, see §3.350(a)(3) of this chapter to determine whether the veteran may be entitled to special monthly compensation.

### THE TORSO AND NECH

	Rat- ing
	3
5319 Group XIX. Function: Support and compression of abdominal wall and lower thorax; flexion	
and lateral motions of spine; synergists in strong	
downward movements of arm (1). Muscles of the	
abdominal wall: (1) Rectus abdominis; (2) external	
oblique; (3) internal oblique; (4) transversalis; (5)	
quadratus lumborum.	
Severe	5
Moderately Severe	3
Moderate	1
Slight	
5320 Group XX. Function: Postural support of body;	
extension and lateral movements of spine. Spinal	
muscles: Sacrospinalis (erector spinae and its pro-	
longations in thoracic and cervical regions).	
Cervical and thoracic region:.	
Severe	4
Moderately Severe	2
Moderate	1
Slight	
Lumbar region:.	
Severe	6 4
Moderate	2
Slight	
5321 Group XXI. Function: Respiration. Muscles of	
respiration: Thoracic muscle group.	
Severe or Moderately Severe	2
Moderate	1
Slight	
5322 Group XXII. Function: Rotary and forward	
movements of the head; respiration; deglutition.	
Muscles of the front of the neck: (Lateral, supra-,	
and infrahyoid group.) (1) Trapezius I (clavicular in-	
sertion); (2) sternocleidomastoid; (3) the "hyoid"	
muscles; (4) sternothyroid; (5) digastric.	
Severe	3
Moderately Severe	2
Moderate	1
Slight	,

THE TORSO AND NECK—Continued

	Rat- ing
5323 Group XXIII. Function: Movements of the head; fixation of shoulder movements. Muscles of the side and back of the neck: Suboccipital; lateral vertebral and anterior vertebral muscles.	
Severe	30
Moderately Severe	20
Moderate	10
Slight	0

#### MISCELLANEOUS

Rat- ing

- 5324 Diaphragm, rupture of, with herniation. Rate under diagnostic code 7346.
- 5325 Muscle injury, facial muscles. Evaluate functional impairment as seventh (facial) cranial nerve neuropathy (diagnostic code 8207), disfiguring scar (diagnostic code 7800), etc. Minimum, if interfering to any extent with mastication—10.
- 5326 Muscle hernia, extensive. Without other injury to the muscle—10.
- 5327 Muscle, neoplasm of, malignant (excluding soft tissue sarcoma)—100.

NOTE: A rating of 100 percent shall continue beyond the cessation of any surgery, radiation treatment, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residual impairment of function.

- 5328 Muscle, neoplasm of, benign, postoperative. Rate on impairment of function, i.e., limitation of motion, or scars, diagnostic code 7805, etc.
- 5329 Sarcoma, soft tissue (of muscle, fat, or fibrous connective tissue)—100.

NOTE: A rating of 100 percent shall continue beyond the cessation of any surgery, radiation treatment, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residual impairment of function.

(Authority: 38 U.S.C. 1155) [62 FR 30239, June 3, 1997]

THE ORGANS OF SPECIAL SENSE

### § 4.75 Examination of visual acuity.

Ratings on account of visual impairments considered for service connection are, when practicable, to be based only on examination by specialists. Such special examinations should include uncorrected and corrected cen-

tral visual acuity for distance and near, with record of the refraction. Snellen's test type or its equivalent will be used. Mydriatics should be routine, except when contraindicated. Funduscopic and ophthalmological findings must be recorded. The best distant vision obtainable after best correction by glasses will be the basis of rating, except in cases keratoconus in which contact lenses are medically required. Also, if there exists a difference of more than 4 diopters of spherical correction between the two eyes, the best possible visual acuity of the poorer eye without glasses, or with a lens of not more than 4 diopters difference from that used with the better eye will be taken as the visual acuity of the poorer eye. When such a difference exists, close attention will be given to the likelihood of congenital origin in mere refractive error.

[40 FR 42537, Sept. 15, 1975]

### §4.76 Examination of field vision.

Measurement of the visual field will be made when there is disease of the optic nerve or when otherwise indicated. The usual perimetric methods will be employed, using a standard perimeter and 3 mm. white test object. At least 16 meridians 221/2 degrees apart will be charted for each eye. (See Figure 1. For the 8 principal meridians, see table III.) The charts will be made a part of the report of examination. Not less than 2 recordings, and when possible, 3 will be made. The minimum limit for this function is established as a concentric central contraction of the visual field to 5°. This type of contraction of the visual field reduces the visual efficiency to zero. Where available the examination for form field should be supplemented, when indicated, by the use of tangent screen or campimeter. This last test is especially valuable in detection of scotoma.

[43 FR 45352, Oct. 2, 1978]

### §4.76a Computation of average concentric contraction of visual fields.

The extent of contraction of visual field in each eye is determined by recording the extent of the remaining visual fields in each of the eight 45 degree principal meridians. The number


## 38 CFR Ch. I (7-1-99 Edition)

### § 4.76a

of degrees lost is determined at each meridian by subtracting the remaining degrees from the normal visual fields given in table III. The degrees lost are then added together to determine total degrees lost. This is subtracted from 500. The difference represents the total remaining degrees of visual field. The difference divided by eight represents the average contraction for rating purposes.

TABLE III—NORMAL VISUAL FIELD EXTENT AT 8
PRINCIPAL MERIDIANS

Meridian	Normal de- grees
Temporally	85
Down temporally	85
Down	65
Down nasally	50
Nasally	60
Up nasally	55
Up	45
Up temporally	55
Total	500



TS-19 52a

Example of computation of concentric contraction under the schedule with abnormal findings taken from Figure 1.

Loss	Degrees
Temporally	55
Down temporally	55
Down	45

#### § 4.77

Loss	Degrees
Down nasally Nasally Up nasally Up temporally	30 40 35 25 35
Total loss	320

Remaining field 500° minus 320°=180°. 180°÷8=22½° average concentric contraction

[43 FR 45352, Oct. 2, 1978]

#### §4.77 Examination of muscle function.

The measurement of muscle function will be undertaken only when the history and findings reflect disease or injury of the extrinsic muscles of the eye, or of the motor nerves supplying these muscles. The measurement will

be performed using a Goldmann Perimeter Chart as in Figure 2 below. The chart identifies four major quadrants, (upward, downward, and two lateral) plus a central field (20 ° or less). The examiner will chart the areas in which diplopia exists, and such plotted chart will be made a part of the examination report. Muscle function is considered normal (20/40) when diplopia does not 40° in the lateral or exist within downward quadrants, or within  $30^{\circ}$  in the upward quadrant. Impairment of muscle function is to be supported in each instance by record of actual appropriate pathology. Diplopia which is only occasional or correctable is not considered a disability.




Figure 2. Goldmann Perimeter Chart

52c

[53 FR 30262, Aug. 11, 1988]

#### §4.78 Computing aggravation.

In determining the effect of aggravation of visual disability, even though the visual impairment of only one eye is service connected, evaluate the vision of both eyes, before and after suffering the aggravation, and subtract the former evaluation from the latter except when the bilateral vision amounts to total disability. In the event of subsequent increase in the disability of either eye, due to intercurrent disease or injury not associated with the service, the condition of the eyes before suffering the subsequent increase will be taken as the basis of compensation subject to the provisions of §3.383(a) of this chapter.

[29 FR 6718, May 22, 1964, as amended at 43 FR 45354, Oct. 2, 1978]

# §4.79 Loss of use of one eye, having only light perception.

Loss of use or blindness of one eye, having only light perception, will be held to exist when there is inability to recognize test letters at 1 foot (.30m.) and when further examination of the eyes reveals that perception of objects, hand movements or counting fingers cannot be accomplished at 3 feet (.91m.), lesser extents of visions, particularly perception of objects, hand movements, or counting fingers at distances less than 3 feet (.91 m.), being considered of negligible utility. With visual acuity 5/200 (1.5/60) or less or the visual field reduced to 5° concentric contraction, in either event in both eyes, the question of entitlement on account of regular aid and attendance will be determined on the facts in the individual case.

[43 FR 45354, Oct. 2, 1978]

## § 4.80 Rating of one eye.

Combined ratings for disabilities of the same eye should not exceed the amount for total loss of vision of that eye unless there is an enucleation or a serious cosmetic defect added to the total loss of vision.

#### §§ 4.81—4.82 [Reserved]

# §4.83 Ratings at scheduled steps and distances.

In applying the ratings for impairment of visual acuity, a person not having the ability to read at any one of the scheduled steps or distances, but reading at the next scheduled step or distance, is to be rated as reading at this latter step or distance. That is, a person who can read at 20/100 (6/30) but who cannot at 20/70 (6/21), should be rated as seeing at 20/100 (6/30).

[41 FR 34257, Aug. 13, 1976, as amended at 43 FR 45354, Oct. 2, 1978]

# $\S 4.83a$ Impairment of central visual acuity.

The percentage evaluation will be found from table V by intersecting the horizontal row appropriate for the Snellen index for one eye and the vertical column appropriate to the Snellen index of the other eye. For example, if one eye has a Snellen index of 5/200 (1.5/60) and the other eye has a Snellen index of 20/70 (6/21), the percentage evaluation is found in the third horizontal row from the bottom and the fourth vertical column from the left. The evaluation is 50 percent and the diagnostic code 6073.

[41 FR 11297, Mar. 18, 1976, as amended at 43 FR 45354, Oct. 2, 1978]

# §4.84 Differences between distant and near visual acuity.

Where there is a substantial difference between the near and distant corrected vision, the case should be referred to the Director, Compensation and Pension Service.

[40 FR 42537, Sept. 15, 1975]

## §4.84a Schedule of ratings—eye.

DISEASES OF THE EYE

		Rat- ing
6000	Uveitis	
6001	Keratitis	
6002	Scleritis	
6003	Iritis	
6004	Cyclitis	
6005	Choroiditis	
6006	Retinitis	
6007	Hemorrhage, intra-ocular, recent	

# § 4.84a

# DISEASES OF THE EYE—Continued

# DISEASES OF THE EYE—Continued

	Rat- ing		Rat- ing
6008 Retina, detachment of		6023 Eyebrows, loss of, complete, unilateral or bi-	
6009 Eye, injury of, unhealed:		lateral	10
The above disabilities, in chronic form, are to be		6024 Eyelashes, loss of, complete, unilateral or bi-	
rated from 10 percent to 100 percent for im-		lateral	10
pairment of visual acuity or field loss, pain,		6025 Epiphora (lacrymal duct, interference with,	
rest-requirements, or episodic incapacity, com-		from any cause):	
bining an additional rating of 10 percent during		Bilateral	20
continuance of active pathology. Minimum rat-		Unilateral	10
ing during active pathology	10	6026 Neuritis, optic:	
6010 Eye, tuberculosis of, active or inactive:	400	Rate underlying disease, and combine impair-	
Active	100	ment of visual acuity or field loss.	
Inactive: See §§ 4.88b and 4.89.		6027 Cataract, traumatic: Preoperative.	
6011 Retina, localized scars, atrophy, or irregular-		Rate on impairment of vision.	
ities of, centrally located, with irregular, duplicated enlarged or diminished image:		Postoperative.	
Unilateral or bilateral	10	Rate on impairment of vision and aphakia.	
6012 Glaucoma, congestive or inflammatory:	10	6028 Cataract, senile, and others:	
Frequent attacks of considerable duration; during		Preoperative.	
continuance of actual total disability	100	Rate on impairment of vision.	
Or, rate as iritis, diagnostic Code 6003.		Postoperative.	
6013 Glaucoma, simple, primary, noncongestive:		Rate on impairment of vision and aphakia.	
Rate on impairment of visual acuity or field loss.		6029 Aphakia:	
Minimum rating	10	Bilateral or unilateral	30
6014 New growths, malignant (eyeball only):		NOTE: The 30 percent rating prescribed for	
Pending completion of operation or other indi-		aphakia is a minimum rating to be applied to	
cated treatment	100	the unilateral or bilateral condition and is not	
Healed; rate on residuals.		to be combined with any other rating for im- paired vision. When only one eye is aphakic,	
6015 New growths, benign (eyeball and adnexa,		the eye having poorer corrected visual acuity	
other than superficial)	4.0	will be rated on the basis of its acuity without	
Rate on impaired vision, minimum	10	correction. When both eyes are aphakic, both	
Healed; rate on residuals. 6016 Nystagmus, central	10	will be rated on corrected vision. The cor-	
6017 Conjunctivitis, trachomatous, chronic:	10	rected vision of one or both aphakic eyes will	
Active; rate for impairment of visual acuity; min-		be taken one step worse than the ascertained	
imum rating while there is active pathology	30	value, however, not better than 20/70 (6/21).	
Healed; rate on residuals, if no residuals	0	Combined ratings for disabilities of the same	
6018 Conjunctivitis, other, chronic:		eye should not exceed the amount for total	
Active, with objective symptoms	10	loss of vision of that eye unless there is an	
Healed; rate on residuals, if no residuals	0	enucleation or a serious cosmetic defect	
6019 Ptosis, unilateral or bilateral:		added to the total loss of vision.	
Pupil wholly obscured.		6030 Accommodation, paralysis of	20
Rate equivalent to 5/200 (1.5/60).		Rate as epiphora.	
Pupile one-half or more obscured.		6032 Eyelids, loss of portion of:	
Rate equivalent to 20/100 (6/30).		Rate as disfigurement. (See diseases of the	
With less interference with vision.		skin.)	
Rate as disfigurement.		6033 Lens, crystalline, dislocation of:	
6020 Ectropion:		Rate as aphakia.	
Bilateral	20 10	6034 Pterygium:	
Unilateral	10	Rate for loss of vision, if any.	
Bilateral	20	6035 Keratoconus: To be evaluated on impairment	
Unilateral	10	of corrected visual acuity using contact lenses.	
6022 Lagophthalmos:	10	NOTE: When contact lenses are medically re-	
Bilateral	20	quired for keratoconus, either unilateral or bi-	
Unilateral	10	lateral, the minimum rating will be 30 percent.	

TABLE IV—TABLE FOR RATING BILATERAL BLINDNESS OR BLINDNESS COMBINED WITH HEARING LOSS WITH DICTATOR'S CODE AND 38 CFR CITATIONS

		Vision other eye			Plus ser	Plus service-connected Hearing loss	ng loss	
Vision one eye	5/200 (1.5/60) or less	Light perception only	No light percep- tion or anatomical loss	Total deafness one ear	10% or 20% at least one ear SC	30% at least one ear SC	40% at least one ear SC	60% or more at least one ear SC
5/200 (1.5/60) or less.	L¹ Code LB–1 38 CFR 3.350(b)(2).	L¹ Code LB–1 38	M Code MB-2 a or b 38 CFR 3.350(f)(2)(ii).	Add ½ step Code No additional PB–1 38 CFR SMC.	No additional SMC.	Add a full step Code PB–3 38 CFR	Add a full step Code PB-3 38 CFR	O Code OB-1 38 CFR 3.350(e)(1)(iii)
Light perception only.		M Code MB-1 a 38 CFR 3.350(c)(1)((iv).	M+½ Code MB-3 O Code OB-2 38 a or b 38 CFR CFR 3.350(f)(iii).	O Code OB-2 38 CFR 3.350(e)(1)(iv).	Add ½ step Code PB-2 38 CFR 3.350(f)(2)(v).	3.350(f)(2)(vi). Add a full step Code PB-3 38 CFR	3.350(f)(2)(vi). O Code OB-2 38 CFR 3.350(e)(1)(iv).	O Code OB-1 38 CFR 3.350(e)(1)(iii)
No light perception or anatomical loss.			N Code NB-1 a-b O Code OB-2 38 or c 38 CFR CFR 3.350(e)(4).	O Code OB-2 38 CFR 3.350(e)(1)(iv).	Add ½ step Code PB-2 38 CFR 3.350(f)(2)(v).	3.350(f)(2)(iv). Add full step Code PB-3 38 CFR 3.350(f)(2)(vi).	3.350(f)(2)(iv).  Add full step Code PB—3 8CPR CFR 3.350(f)(2)(vi). 3.350(f)(1)(iv).	O Code OB-1 38 CFR 3.350(e)(1)(iii)

1 With need for aid and attendance qualifies for Subpar. m. code MB-1, b; 38 CFR 3.350(c)(1)(v).

NOTE.—(1) Any of the additional SMC payable under Dictator's Codes PB-1, PB-2, or PB-3 is not to exceed the rate payable under Subpar. O. (2) If in addition to any of the above the veteran has the service-connected loss or loss of use of an extremity, additional SMC is payable, not to exceed the rate payable under Subpar. O. See Dictator's Codes PB-4, PB-5, PB-6, and 38 CFR 3.350(f)(2)(vii) (A), (B), (C).

(Authority: 38 U.S.C. 1115)

# § 4.84a

## IMPAIRMENT OF CENTRAL VISUAL ACUITY

		Rat- ing
6061	Anatomical loss both eyes	5 100
6062	Blindness in both eyes having only light per-	
	tion	5 100
Α	natomical loss of 1 eye:	
6063	In the other eye 5/200 (1.5/60)	5 100
6064	In the other eye 10/200 (3/60)	6 90
6064	In the other eye 15/200 (4.5/60)	680
6064	In the other eye 20/200 (6/60)	670
6065	In the other eye 20/100 (6/30)	6 60
6065	In the other eye 20/70 (6/21)	6 60
6065	In the other eye 20/50 (6/15)	<sup>6</sup> 50
6066	In the other eye 20/40 (6/12)	640
	lindness in 1 eye, having only light perception:	
6067	In the other eye 5/200 (1.5/60)	5 100
6068	In the other eye 10/200 (3/60)	5 90
6068	In the other eye 15/200 (4.5/60)	5 80
6068	In the other eye 20/200 (6/60)	5 70
6069	In the other eye 20/100 (6/30)	5 60
6069	In the other eye 20/70 (6/21)	5 50
6069	In the other eye 20/50 (6/15)	5 40
6070	In the other eye 20/40 (6/12)	5 30
	ision in 1 eye 5/200 (1.5/60):	
6071	In the other eye 5/200 (1.5/60)	5 100
6072	In the other eye 10/200 (3/60)	90
6072	In the other eye 15/200 (4.5/60)	80
6072	In the other eye 20/200 (6/60)	70
6073	In the other eye 20/100 (6/30)	60
6073	In the other eye 20/70 (6/21)	50
6073	In the other eye 20/50 (6/15)	40
6074	In the other eye 20/40 (6/12)	30
V	ision in 1 eye 10/200 (3/60):	
6075	In the other eye 10/200 (3/60)	90
6075	In the other eye 15/200 (4.5/60)	80
6075	In the other eye 20/200 (6/60)	70
6076	In the other eye 20/100 (6/30)	60

# IMPAIRMENT OF CENTRAL VISUAL ACUITY— Continued

		Rat- ing
6076	In the other eye 20/70 (6/21)	50
6076	In the other eye 20/50 (6/15)	40
6077	In the other eye 20/40 (6/12)	30
V	ision in 1 eye 15/200 (4.5/60):	
6075	In the other eye 15/200 (4.5/60)	80
6075	In the other eye 20/200 (6/60)	70
6076	In the other eye 20/100 (6/30)	60
6076	In the other eye 20/70 (6/21)	40
6076	In the other eye 20/50 (6/15)	30
6077	In the other eye 20/40 (6/12)	20
	ision in 1 eye 20/200 (6/60):	
6075	In the other eye 20/200 (6/60)	70
6076	In the other eye 20/100 (6/30)	60
6076	In the other eye 20/70 (6/21)	40
6076	In the other eye 20/50 (6/15)	30
6077	In the other eye 20/40 (6/12)	20
	ision in 1 eye 20/100 (6/30):	
6078	In the other eye 20/100 (6/30)	50
6078	In the other eye 20/70 (6/21)	30
6078	In the other eye 20/50 (6/15)	20
6079	In the other eye 20/40 (6/12)	10
	ision in 1 eye 20/70 (6/21):	
6078		30
6078	In the other eye 20/50 (6/15)	20
6079	In the other eye 20/40 (6/12)	10
078 6078	ision in 1 eye 20/50 (6/15):	10
		10
6079	In the other eye 20/40 (6/12)ision in 1 eye 20/40 (6/12):	10
	the other eye 20/40 (6/12):	0
	1 110 Othor Cyc 20170 (0/12)	

<sup>&</sup>lt;sup>5</sup> Also entitled to special monthly compensation. 
<sup>6</sup> Add 10% if artificial eye cannot be worn; also entitled to special monthly compensation.

# TABLE V—RATINGS FOR CENTRAL VISUAL ACUITY IMPAIRMENT [With Diagnostic Code]

						•			
				٧	ision in othe	r eye			
Vision in one eye	20/40 (6/ 12)	20/50 (6/ 15)	20/70 (6/ 21)	20/100 (6/30)	20/200 (6/60)	15/200 (4.5/60)	10/200 (3/60)	5/200 (1.5/60)	Light perception only/anatomical loss
20/40 (6/12)	0								
20/50 (6/15)	10 (6079)	10 (6078)							
20/70 (6/21)	10 (6079)	20 (6078)	30 (6078)						
20/100 (6/30)	10 (6079)	20 (6078)	30 (6078)	50 (6078)					
20/200 (6/60)	20 (6077)	30 (6076)	40 (6076)	60 (6076)	70 (6075)				
15/200 (4.5/60)	20 (6077)	30 (6076)	40 (6076)	60 (6076)	70 (6075)	80 (6075)			
10/200 (3/60)	30 (6077)	40 (6076)	50 (6076)	60 (6076)	70 (6075)	80 (6075)	90 (6075)		
5/200 (1.5/60)	30 (6074)	40 (6073)	50 (6073)	60 (6073)	70 (6072)	80 (6072)	90 (6072)	<sup>5</sup> 100 (6071)	
Light per- ception only	530	5 40	<sup>5</sup> 50	<sup>5</sup> 60	5 70	5 80	5 90	5 100	5 100

#### TABLE V—RATINGS FOR CENTRAL VISUAL ACUITY IMPAIRMENT—Continued [With Diagnostic Code]

				٧	ision in othe	r eye			
Vision in one eye	20/40 (6/ 12)	20/50 (6/ 15)	20/70 (6/ 21)	20/100 (6/30)	20/200 (6/60)	15/200 (4.5/60)	10/200 (3/60)	5/200 (1.5/60)	Light perception only/anatomical loss
	(6070)	(6069)	(6069)	(6069)	(6068)	(6068)	(6068)	(6067)	(6062)
Anatomical loss of one eye	640 (6066)	<sup>6</sup> 50 (6065)	<sup>6</sup> 60 (6065)	<sup>6</sup> 60 (6065)	<sup>6</sup> 70 (6064)	6 80 (6064)	<sup>6</sup> 90 (6064)	<sup>5</sup> 100 (6063)	<sup>5</sup> 100 (6061)

#### RATINGS FOR IMPAIRMENT OF FIELD VISION

	Rat- ing
6080 Field vision, impairment of:	
Homonymous hemianopsia	30
Field, visual, loss of temporal half:	
Bilateral	30
Unilateral	10
Or rate as 20/70 (6/21).	
Field, visual, loss of nasal half:	
Bilateral	20
Unilateral	10
Or rate as 20/50 (6/15).	
Field, visual, concentric contraction of:	
To 5°:	
Bilateral	100
Unilateral	30
Or rate as 5/200 (1.5/60).	
To 15° but not to 5°:	<b>-</b>
Bilateral	70
Unilateral	20
Or rate as 20/200 (6/60). To 30° but not to 15°:	
Bilateral	50 10
Unilateral	10
Or rate as 20/100 (6/30). To 45° but not to 30°:	
Bilateral	30
Unilateral	10
Or rate as 20/70 (6/21): To 60° but not to 45°:	
Bilateral	20
Unilateral	10
Or rate as 20/50 (6/15).	
NOTE (1): Correct diagnosis reflecting disease or	

injury should be cited. NOTE (2): Demonstrable pathology commensu-

rate with the functional loss will be required. The concentric contraction ratings require contraction within the stated degrees, temporally; the nasal contraction may be less. The alternative ratings are to be employed when there is ratable defect of visual acuity, or a different impairment of the visual field in the other eye. Concentric contraction resulting from demonstrable pathology to 5 degrees or less will be considered on a parity with reduction of central visual acuity to 5/200 (1.5/60) or less for all purposes including entitlement under \$3.350(b)(2) of this chapter; not however, for the purpose of §3.350(a) of this chapter. Entitlement on account of blindness requiring regular aid and attendance, §3.350(c) of this chapter, will continue to be determined on the facts in the individual case.

6081 Scotoma, pathological, unilateral:

#### RATINGS FOR IMPAIRMENT OF FIELD VISION-Continued

	Rat- ing
Large or centrally located, minimum	10

#### RATINGS FOR IMPAIRMENT OF MUSCLE FUNCTION [6090 Diplopia (double vision)]

Degree of diplopia	Equiva- lent visual acuity
(a) Central 20°	5/200
(b) 21° to 30°:	
(1) Down	15/200
(2) Lateral	20/100
(3) Up	20/70
(c) 31° to 40°:	
(1) Down	20/200
(2) Lateral	20/70
(3) Up	20/40

NOTES: (1) Correct diagnosis reflecting disease or injury should be cited.

- (2) The above ratings will be applied to only one eye. Ratings will not be applied for both diplopia and decreased visual acuity or field of vision in the same eye. When diplopia is present and there is also ratable impairment of visual acuity or field of vision of both eyes the above diplopia ratings will be applied to the poorer eye while the better eye is rated according to the best corrected visual acuity or visual field.
- (3) When the diplopia field extends beyond more than one quadrant or more than one range of degrees, the evaluation for diplopia will be based on the quadrant and degree range that provide the highest evaluation.
- (4) When diplopia exists in two individual and separate areas of the same eye, the equivalent visual acuity will be taken one step worse, but no worse than 5/200.

6091 Symblepharon.

 <sup>&</sup>lt;sup>5</sup> Also entitled to special monthly compensation.
 <sup>6</sup>Add 10 percent if artificial eye cannot be worn; also entitled to special monthly compensation.

Rate as limited muscle function, diagnostic code 6090.

6092 Diplopia, due to limited muscle function.

Rate as diagnostic code 6090.

[29 FR 6718, May 22, 1964, as amended at 34 FR 5062, Mar. 11, 1969; 40 FR 42537, Sept. 15, 1975; 41 FR 11297, Mar. 18, 1976; 43 FR 45354, Oct. 2, 1978; 51 FR 6411, Feb. 24, 1986; 53 FR 30264, Aug. 11, 1988; 53 FR 50955, Dec. 19, 1988; 57 FR 24364, June 9, 1992]

IMPAIRMENT OF AUDITORY ACUITY

# §4.85 Evaluation of hearing impairment.

- (a) An examination for hearing impairment for VA purposes must be conducted by a state-licensed audiologist and must include a controlled speech discrimination test (Maryland CNC) and a puretone audiometry test. Examinations will be conducted without the use of hearing aids.
- (b) Table VI, "Numeric Designation of Hearing Impairment Based on Puretone Threshold Average and Speech Discrimination," is used to determine a Roman numeral designation (I through XI) for hearing impairment based on a combination of the percent of speech discrimination (horizontal rows) and the puretone threshold average (vertical columns). The Roman numeral designation is located at the point where the percentage of speech discrimination and puretone threshold average intersect.
- (c) Table VIa, "Numeric Designation of Hearing Impairment Based Only on Puretone Threshold Average," is used to determine a Roman numeral designation (I through XI) for hearing im-

pairment based only on the puretone threshold average. Table VIa will be used when the examiner certifies that use of the speech discrimination test is not appropriate because of language difficulties, inconsistent speech discrimination scores, etc., or when indicated under the provisions of §4.86.

(d) "Puretone threshold average," as used in Tables VI and VIa, is the sum of the puretone thresholds at 1000, 2000, 3000 and 4000 Hertz, divided by four. This average is used in all cases (including those in §4.86) to determine the Roman numeral designation for hearing impairment from Table VI or VIa.

- (e) Table VII, "Percentage Evaluations for Hearing Impairment," is used to determine the percentage evaluation by combining the Roman numeral designations for hearing impairment of each ear. The horizontal rows represent the ear having the better hearing and the vertical columns the ear having the poorer hearing. The percentage evaluation is located at the point where the row and column intersect.
- (f) If impaired hearing is service-connected in only one ear, in order to determine the percentage evaluation from Table VII, the non-service-connected ear will be assigned a Roman Numeral designation for hearing impairment of I, subject to the provisions of §3.383 of this chapter.
- (g) When evaluating any claim for impaired hearing, refer to §3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation due either to deafness, or to deafness in combination with other specified disabilities.
  - (h) Numeric tables VI, VIA\*, and VII.

TABLE VI

# NUMERIC DESIGNATION OF HEARING IMPAIRMENT BASED ON PURETONE THRESHOLD AVERAGE AND SPEECH DISCRIMINATION

## **Puretone Threshold Average**

% of discrim- ination	0-41	42-49	50-57	58-65	66-73	74-81	82-89	90-97	98+
92-100	I	I	I	II	II	II	III	III	IV
84-90	11	II	II	III	III	III	IV	IV	IV
76-82	III	III	IV	IV	IV	V	V	V	V
68-74	IV	IV	V	V	VI	VI	VII	VII	VII
60-66	V	v	VI	VI	VII	VII	VIII	VIII	VIII
52-58	VI	VI	VII	VII	VIII	VIII	VIII	VIII	IX
44-50	VII	VII	VIII	VIII	VIII	IX	IX	IX	X
36-42	VIII	VIII	VIII	IX	IX	IX	X	X	X
0-34	IX	X	XI						

## TABLE VIA\*

# NUMERIC DESIGNATION OF HEARING IMPAIRMENT BASED ONLY ON PURETONE THRESHOLD AVERAGE

### **Puretone Threshold Average**

0-41	42-48	49-55	56-62	63-69	70-76	77-83	84-90	91-97	98-104	105+
I	II	III	IV	V	VI	VII	VIII	IX	X	XI

<sup>\*</sup> This table is for use only as specified in §§ 4.85 and 4.86.

### TABLE VII

# PERCENTAGE EVALUATION FOR HEARING IMPAIRMENT (DIAGNOSTIC CODE 6100)

#### **Poorer Ear**

	XI	100*										
	X	90	80									
	IX	80	70	60								
	VIII	70	60	50	50				And the second s			
ar	VII	60	60	50	40	40			The state of the s			
Better Ear	VI	50	50	40	40	30	30	MANA CHARACTER STREET, CO.				
Ř	V	40	40	40	30	30	20	20				
	IV	30	30	30	20	20	20	10	10			
	III	20	20	20	20	20	10	10	10	0		
	II	10	10	10	10	10	10	10	0 .	0	0	
-	I	10	10	0	0	0	0	0	0	. 0	0	0
i		XI	X	IX	VIII	VII	VI	V	IV	III	11	I

<sup>\*</sup> Review for entitlement to special monthly compensation under §3.350 of this chapter.

[64 FR 25206, May 11, 1999]

# § 4.86 Exceptional patterns of hearing impairment.

(a) When the puretone threshold at each of the four specified frequencies (1000, 2000, 3000, and 4000 Hertz) is 55 decibels or more, the rating specialist will determine the Roman numeral designation for hearing impairment from either Table VI or Table VIa, whichever results in the higher numeral. Each ear will be evaluated separately.

(b) When the puretone threshold is 30 decibels or less at 1000 Hertz, and 70 decibels or more at 2000 Hertz, the rating specialist will determine the Roman numeral designation for hearing impairment from either Table VI or Table VIa, whichever results in the higher numeral. That numeral will then be elevated to the next higher

#### **Department of Veterans Affairs**

§ 4.88a

Roman numeral. Each ear will be evaluated separately.

(Authority: 38 U.S.C. 1155) [64 FR 25209, May 11, 1999]

#### §4.87 Schedule of ratings—ear.

## DISEASES OF THE EAR

Rating

10

30 10

100

60

DISEASES OF THE EAR
6200 Chronic suppurative otitis media, mastoiditis, or cholesteatoma (or any combination):  During suppuration, or with aural polyps  NOTE: Evaluate hearing impairment, and complications such as labyrinthitis, tinnitus, facial nerve paralysis, or bone loss of skull, separately.
6201 Chronic nonsuppurative otitis media with effusion (serous otitis media): Rate hearing impairment
6202 Otosclerosis: Rate hearing impairment
6204 Peripheral vestibular disorders:  Dizziness and occasional staggering  Occasional dizziness  NOTE: Objective findings supporting the diagnosis of vestibular disequilibrium are required
before a compensable evaluation can be as- signed under this code. Hearing impairment or suppuration shall be separately rated and combined.
6205 Meniere's syndrome (endolymphatic hydrops): Hearing impairment with attacks of vertigo and cerebellar gait occurring more than once weekly, with or without tinnitus Hearing impairment with attacks of vertigo and cerebellar gait occurring from one to four
times a month, with or without tinnitus
hearing impairment, and tinnitus, whichever method results in a higher overall evaluation. But do not combine an evaluation for hearing impairment, tinnitus, or vertigo with an evaluation under diagnostic code 6205.  6207 Loss of auricle:
Complete loss of both  Complete loss of one  Deformity of one, with loss of one-third or more of the substance
6208 Malignant neoplasm of the ear (other than skin only)  NOTE: A rating of 100 percent shall continue beyond the cessation of any surgical, radiation treatment, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by
mandatory VA examination. Any change in

evaluation based on that or any subsequent examination shall be subject to the provisions

of §3.105(e) of this chapter. If there has been

no local recurrence or metastasis, rate on re-

6209 Benign neoplasms of the ear (other than skin

Rate on impairment of function.

6210 Chronic otitis externa:

DISEASES OF THE EAR—Continued

	Rat- ing
Swelling, dry and scaly or serous discharge, and itching requiring frequent and prolonged treatment	10 0 10

(Authority: 38 U.S.C. 1155) [64 FR 25210, May 11, 1999]

# § 4.87a Schedule of ratings—other sense organs.

	ing
6275 Sense of smell, complete loss	10
6276 Sense of taste, complete loss	10
NOTE: Evaluation will be assigned under diag- nostic codes 6275 or 6276 only if there is an anatomical or pathological basis for the condi- tion.	

(Authority: 38 U.S.C. 1155)

[64 FR 25210, May 11, 1999]

Infectious Diseases, Immune Disorders and Nutritional Deficiencies

#### §4.88 [Reserved]

## §4.88a Chronic fatigue syndrome.

- (a) For VA purposes, the diagnosis of chronic fatigue syndrome requires:
- (1) new onset of debilitating fatigue severe enough to reduce daily activity to less than 50 percent of the usual level for at least six months; and
- (2) the exclusion, by history, physical examination, and laboratory tests, of all other clinical conditions that may produce similar symptoms; and
  - (3) six or more of the following:
  - (i) acute onset of the condition,
  - (ii) low grade fever,
  - (iii) nonexudative pharyngitis,
- (iv) palpable or tender cervical or axillary lymph nodes,
- (v) generalized muscle aches or weakness,
- (vi) fatigue lasting 24 hours or longer after exercise,

# § 4.88b

(vii) headaches (of a type, severity, or pattern that is different from headaches in the pre-morbid state),

(viii) migratory joint pains,

(ix) neuropsychologic symptoms,(x) sleep disturbance.(b) [Reserved]

[59 FR 60902, Nov. 29, 1994]

#### §4.88b Schedule of ratings-infectious diseases, immune disorders and nutritional deficiencies.

	Rating
6300 Cholera, Asiatic:	
As active disease, and for 3 months convalescence	100
5301 Visceral Leishmaniasis:  During treatment for active disease	100
NOTE: A 100 percent evaluation shall continue beyond the cessation of treatment for active disease. Six m continuance of such treatment, the appropriate disability rating shall be determined by mandatory VA exchange in evaluation based upon that or any subsequent examination shall be subject to the provisions this chapter. Rate residuals such as liver damage or lymphadenopathy under the appropriate system.  6302 Leprosy (Hansen's Disease):	amination. Any
As active disease	100
NOTE: A 100 percent evaluation shall continue beyond the date that an examining physician has determine become inactive. Six months after the date of inactivity, the appropriate disability rating shall be determined tory VA examination. Any change in evaluation based upon that or any subsequent examination shall be provisions of § 3.105(e) of this chapter. Rate residuals such as skin lesions or peripheral neuropathy unpriate system.	ned by manda- e subject to the
6304 Malaria: As active disease	100
NOTE: The diagnosis of malaria depends on the identification of the malarial parasites in blood smears.	
served in an endemic area and presents signs and symptoms compatible with malaria, the diagnosis matchinical grounds alone. Relapses must be confirmed by the presence of malarial parasites in blood smears. Thereafter rate residuals such as liver or spleen damage under the appropriate system.	ay be based on
6305 Lymphatic Filariasis:  As active disease  Thereafter rate residuals such as epididymitis or lymphanditis under the appropriate system	100
6306 Bartonellosis: As active disease, and for 3 months convalescence	100
Thereafter rate residuals such as skin lesions under the appropriate system 6307 Plague:	1
As active disease	100
6308 Relapsing Fever:	1
As active disease Thereafter rate residuals such as liver or spleen damage or central nervous system involvement under the a tem	
6309 Rheumatic fever:	100
As active disease	100
6310 Syphilis, and other treponemal infections: Rate the complications of nervous system, vascular system, eyes or ears. (See DC 7004, syphilitic hea 8013, cerebrospinal syphilis, DC 8014, meningovascular syphilis, DC 8015, tabes dorsalis, and DC 9301, ciated with central nervous system syphilis)	
6311 Tuberculosis, miliary: As active disease	100
6313 Avitaminosis:  Marked mental changes, moist dermatitis, inability to retain adequate nourishment, exhaustion, and cachexis  With all of the symptoms listed below, plus mental symptoms and impaired bodily vigor	
With stomatitis, diarrhea, and symmetrical dermatitis	40
With stomatitis, or achlorhydria, or diarrhea  Confirmed diagnosis with nonspecific symptoms such as: decreased appetite, weight loss, abdominal discortor, weakness, inability to concentrate and irritability	n-
6314 Beriberi: As active disease:	
With congestive heart failure, anasarca, or Wernicke-Korsakoff syndrome	
With cardiomegaly, or; with peripheral neuropathy with footdrop or atrophy of thigh or calf muscles	
weakness, fatigue, anorexia, dizziness, heaviness and stiffness of legs, headache or sleep disturbance Thereafter rate residuals under the appropriate body system.	
6315 Pellagra:	

With stomatitis, diarrhea, and symmetrical dermatitis with stomatitis, or architorytria, or diarrhea Confirmed diagnosis with nonspecific symptoms such as: decreased appetite, weight loss, abdominal discomfort, weakness, inability to concentrate and irritability with stomatics, or architorytria, or diarrhea, and the state of the		Rating
With stomatitis, or achlorhydria, or diarrhea Confirmed diagnosis with nonspecific symptoms such as: decreased appetite, weight loss, abdominal discomfort, weakness, inability to concentrate and irritability  16 Brucellosis: As active disease Thereafter rate residuals such as liver or spleen damage or meningitis under the appropriate system  17 Typhus, scrub: As active disease, and for 3 months convalescence Thereafter rate residuals such as spleen damage or skin conditions under the appropriate system  18 Melioridosis: As active disease. Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system  19 Lyme Disease: As active disease Thereafter rate residuals such as arthritis under the appropriate system  20 Parasitic diseases otherwise not specified: As active disease Thereafter rate residuals such as a spleen or fiver damage under the appropriate system  210 Parasitic diseases otherwise not specified: As active disease Thereafter rate residuals such as a spleen or fiver damage under the appropriate system  210 Parasitic diseases otherwise not specified: As active disease Thereafter rate residuals such as a spleen or fiver damage under the appropriate system  210 Parasitic diseases otherwise not specified: As active disease Thereafter rate residuals such as a spleen or fiver damage under the appropriate system  210 Lupus erythematosus, systemic (disseminated): Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health Exacerbations lasting a week or more, 2 or 3 times per year  Not:: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 5350, whichever method results in a higher evaluation.  211 HV-Related Illness: AllDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions.  211 Refracto	With all of the symptoms listed below, plus mental symptoms and impaired bodily vigor	6
Confirmed diagnosis with nonspecific symptoms such as: decreased appetite, weight loss, abdominal discomfort, weakness, inability to concentrate and irritability  16 Brucellosis: As active disease Thereafter rate residuals such as liver or spleen damage or meningitis under the appropriate system  17 Typhus, scrub: As active disease, and for 3 months convalescence Thereafter rate residuals such as spleen damage or skin conditions under the appropriate system  18 Melioidosis: As active disease Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system  18 J. Lyme Disease: As active disease Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system  19 J. Lyme Disease: As active disease Thereafter rate residuals such as arthritis under the appropriate system  200 Parasitic diseases otherwise not specified: As active disease Thereafter rate residuals such as spleen or liver damage under the appropriate system  201 Lupus enythematosus, systemic (disseminated): Not be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health Exacerbations once or twice a year or symptomatic during the past 2 years  Notte: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  31 HIV-Related Illness: AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness: ARFarctory constitutional symptoms, intermittent diarrhea, and on approved medication(s), or; minimum rating following development of definite medical symptoms, intermittent diarrhea, and on approved medication(s), or; minimum rating with 17 cell count less than 200, or Haliy Cell Leukoplakia, or Oral Candidiasis  Following development of definite medical symptoms, intermittent diarrhea, and on approved medication(s), or; minimum rating with 17 cell count le		4
10. Brucellosis: As active disease. Thereafter rate residuals such as spleen damage or meningitis under the appropriate system 17. Typhus, scrub: As active disease, and for 3 months convalescence Thereafter rate residuals such as spleen damage or skin conditions under the appropriate system 18. Melioidosis: As active disease, and for 3 months convalescence Thereafter rate residuals such as spleen damage or skin conditions under the appropriate system 18. Melioidosis: As active disease. Thereafter tate residuals such as arthritis, lung lesions or meningitis under the appropriate system 19. Lynne Disease: As active disease. Thereafter rate residuals such as arthritis under the appropriate system 19. Parasitic diseases otherwise not specified: As active diseases otherwise not specified: As active diseases. Thereafter rate residuals such as aptent or liver damage under the appropriate system 10. Lynus erythematosus, systemic (disseminated): Not to be combined with ratings under DC 7803 Acute, with frequent exacerbations, producing severe impairment of health. Exacerbations once or twice a year or symptomate during the past 2 years. Nore: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 850s, whichever method results in a higher evaluation. 13.1 HIV-Related Illness: All DS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions. Refractory constitutional symptoms, distrate, and pathological weight loss, or minimum rating with T4 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidiasis Following development of definite medical symptoms, T4 cell of 200 or one and less than 500, and on approved medication(s), or, with evidence of depression or memory loss with employment limitations. Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or d		2
As active disease.  Thereafter rate residuals such as liver or spleen damage or meningitis under the appropriate system  17 Typhus, scrub:  As active disease, and for 3 months convalescence.  Thereafter rate residuals such as pleen damage or skin conditions under the appropriate system  18 Melioidosis:  As active disease.  Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system  18 Melioidosis:  As active disease.  Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system  19 Parasitic diseases otherwise not specified:  As active diseases of the state of the appropriate system  20 Parasitic diseases otherwise not specified:  As active diseases, and the state residuals such as appropriate system  20 Parasitic diseases otherwise not specified:  As active diseases, and the state residuals such as appropriate system  20 Parasitic diseases otherwise not specified:  As active diseases, and the state residuals such as appropriate system  20 Parasitic diseases otherwise not specified:  As active diseases, and the appropriate system  20 Parasitic diseases otherwise not specified:  As active diseases, and the appropriate system  20 Lipus eyrthematosus, systemic (disseminated):  Not to be combined with ratings under DC 75090 Acute, with frequent exacerbations, producing severe impairment of health.  Not to be combined with ratings under DC 75090 Acute, with frequent exacerbations, producing severe impairment of health.  Not E (12 Pataset this condition either by combining the evaluation or residuals under the appropriate system, or by evaluation of costs.  All Swith recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility appropriate system evaluation.  All V-Related illness with debility appropriate system evaluation or repolates		
As active disease  Thereafter rate residuals such as liver or spleen damage or meningitis under the appropriate system  17. Typhus, scrub.  As active disease, and for 3 months convalescence  Thereafter rate residuals such as spleen damage or skin conditions under the appropriate system  18. As active disease.  Thereafter rate residuals such as arthrifts under the appropriate system  180. Parasitic diseases otherwise not specified:  As active disease.  Thereafter rate residuals such as spleen or liver damage under the appropriate system  180. Lupus enythematosus, systemic (disseminated):  Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health  Exacerbations lasting a week or more, 2 or 3 times per year  Exacerbations once or twice a year or symptomatic during the past 2 years  NOTE: Evaluate this condition either by combining the evaluation.  11. HIV-Related Illness:  AlDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions.  Refractory constitutional symptoms, diarrhea, and pathological weight loss, or; minimum rating following development of definite medical symptoms, Ta cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations.  Following development of definite medical symptoms, Ta cell of 200 or more and less than 500, and on approved medication(s) or with evidence or depression or memory loss with employment limitations.  Following faiting exporting interior dispersions or memory loss with employment limitat		1
Thereafter rate residuals such as liver or spleen damage or meningitis under the appropriate system 317 Typhus, scrub:  As active disease, and for 3 months convalescence Thereafter rate residuals such as spleen damage or skin conditions under the appropriate system 318 Melioidosis:  As active disease.  Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system 318 Melioidosis:  As active disease.  Thereafter rate residuals such as arthritis under the appropriate system 320 Parasitic diseases otherwise not specified:  As active diseases.  Thereafter rate residuals such as arthritis under the appropriate system  500 Lupus erythematosus, systemic (diseaminated):  As active diseases.  Thereafter rate residuals such as arthritis under the appropriate system  510 Lupus erythematosus, systemic (diseaminated):  As active diseases.  As active		
As active disease, and for 3 months convalescence Thereafter rate residuals such as spleen damage or skin conditions under the appropriate system 318 Melioidosis: As active disease Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system 319 Lyme Disease: As active disease Thereafter rate residuals such as arthritis under the appropriate system 320 Parasitic diseases otherwise not specified: As active disease Thereafter rate residuals such as spleen or liver damage under the appropriate system 320 Lypus epithematious, systemic (disease) Thereafter rate residuals such as spleen or liver damage under the appropriate system 350 Lypus epithematious, systemic (disease) Thereafter rate residuals such as spleen or liver damage under the appropriate system 350 Lypus epithematious, systemic (disease) Thereafter rate residuals such as spleen or liver damage under the appropriate system 350 Lypus epithematious, systemic (disease) Thereafter rate residuals such as spleen or liver damage under the appropriate system 350 Lypus epithematious, systemic (disease) Thereafter rate residuals such as spleen or liver damage under the appropriate system 350 Lypus epithematious, systemic (disease) Thereafter late rate residuals such as spleen or liver damage under the appropriate system 350 Lypus epithematicus, systemic (disease) Thereafter rate residuals such as spleen or liver damage under the appropriate system 350 Lypus epithematicus, systemic (disease) Thereafter rate residuals such as spleen or liver damage under the appropriate system 351 Lypus epithematicus on systemic diseases afflicting multiple body systems; HIV-related illness: AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions Recurrent constitutional symptoms, diarrhea, and pathological weight loss, or, minimum rating of evolution of AIDS-related opportunistic infect		10
As active disease, and for 3 months convalescence Thereafter rate residuals such as spleen damage or skin conditions under the appropriate system  318 Melioidosis: As active disease Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system  319 Lyme Disease: As active disease Thereafter rate residuals such as arthritis under the appropriate system  320 Parasitic diseases otherwise not specified: As active diseases Thereafter rate residuals such as spleen or liver damage under the appropriate system  320 Parasitic diseases otherwise not specified: As active diseases Thereafter rate residuals such as spleen or liver damage under the appropriate system  350 Lupus erythematosus, systemic (disseminated): Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health Exacerbations lasting a week or more, 2 or 3 times per year Exacerbations once or twice a year or symptomatic during the past 2 years  Note: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  351 HIV-Related Illness: AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions  Refractory constitutional symptoms, diarrhea, and pathological weight loss, or, minimum rating following development of definite medical symptoms, and pathological weight loss, or, minimum rating following development of definite medical symptoms, and pathological weight loss, or, minimum rating following development of definite medical symptoms, resolution or neoplasm  Recurrent constitutional symptoms, intermittent diarrhea, and on approved medication(s), or, minimum rating following development of definite medical symptoms, resolution or neoplasm consuments of the pre-illness level, or, with resolution		
Thereafter rate residuals such as spleen damage or skin conditions under the appropriate system  318 Melioidosis:  As active disease Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system  319 Lyme Disease: As active disease Thereafter rate residuals such as arthritis under the appropriate system  320 Parasitic diseases otherwise not specified: As active disease Thereafter rate residuals such as spleen or liver damage under the appropriate system  330 Lupus erythematous, systemic (diseaseimiated): Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health Exacerbations once or twice a year or symptomatic during the past 2 years Exacerbations once or twice a year or symptomatic during the past 2 years MOTE: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  31 HIV-Related Illness: AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions Refractory constitutional symptoms, diarrhea, and pathological weight loss, or, minimum rating following development of AIDS-related opportunistic infection or neoplasm Recurrent constitutional symptoms, internal, and on approved medication(s), or, with evidence of depression or memory loss with employment limitations Asymptomalitic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count more approved medication of definite medical symptoms, increased and an accredited medical institution.  NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated sep arately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.		4.0
As active disease Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system 319 Lyme Disease: As active disease Thereafter rate residuals such as arthritis under the appropriate system 320 Parasitic diseases otherwise not specified: As active diseases Thereafter rate residuals such as spleen or liver damage under the appropriate system 350 Lupus erythematosus, systemic (disseminated): Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health Exacerbations lasting a week or more, 2 or 3 times per year Exacerbations lasting a week or more, 2 or 3 times per year Exacerbations lasting a week or more, 2 or 3 times per year Exacerbations lasting a week or more, 2 or 3 times per year Exacerbations lasting as week or more, 2 or 3 times per year Exacerbations lasting as week or more, 2 or 3 times per year Exacerbations lasting as week or more, 2 or 3 times per year Exacerbations lasting as week or more, 2 or 3 times per year Exacerbations lasting as week or more, 2 or 3 times per year Exacerbations lasting as week or more, 2 or 3 times per year Exacerbations lasting as week or more, 2 or 3 times per year Exacerbations lasting as week or more, 2 or 3 times per year Exacerbations lasting as week or more, 2 or 3 times per year Exacerbations lasting as week or more, 2 or 3 times per year Exacerbations lasting as week or more, 2 or 3 times per year Exacerbations lasting as week or more, 2 or 3 times per year  Exacerbations lasting as week or more, 2 or 3 times per year  Exacerbations once or twice a year or symptomatic during the past 2 years  NOTE: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluation.  Nore (2): Calculated liness with debility and progressive weight loss, or; minimum rating following development of definite medical symptoms, 74 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory		10
As active disease Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system 319 Lyme Disease: As active disease Thereafter rate residuals such as arthritis under the appropriate system 320 Parasitic diseases otherwise not specified: As active disease Thereafter rate residuals such as arthritis under the appropriate system 320 Lypus erythematosus, systemic (disseminated): Soft Lupus erythematosus, systemic (disseminated): Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health Exacerbations once or twice a year or symptomatic during the past 2 years MOTE: Evaluate this condition either by combining the evaluation for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation. 311 HIV-Related Illness: AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions		
Thereafter rate residuals such as arthritis, lung lesions or meningitis under the appropriate system  319 Lyme Disease:  As active disease  Thereafter rate residuals such as arthritis under the appropriate system  320 Parasitic diseases otherwise not specified:  As active disease  Thereafter rate residuals such as spleen or liver damage under the appropriate system  350 Lupus erythematosus, systemic (disseminated):  Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health  Exacerbations once or twice a year or symptomatic during the past 2 years  Notre: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  351 HIV-Related Illness:  AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions  Refractory constitutional symptoms, diarrhea, and pathological weight loss, or; minimum rating following development of AIDS-related opportunistic infection or neoplasm  Recurrent constitutional symptoms, intermittent diarrhea, and on approved medication(s), or; minimum rating with T4 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidiasis.  Following development of definite medical symptoms, T4 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations		10
As active disease Thereafter rate residuals such as arthritis under the appropriate system 320 Parasitic diseases otherwise not specified: As active diseases Thereafter rate residuals such as spleen or liver damage under the appropriate system 350 Lupus erythematosus, systemic (disseminated): Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health Exacerbations lasting a week or more, 2 or 3 times per year Exacerbations once or twice a year or symptomatic during the past 2 years NOTE: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation. 31 HIV-Related Illness: AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions.  Refractory constitutional symptoms, diarrhea, and pathological weight loss, or; minimum rating following development of AIDS-related opportunistic infection or neoplasm Recurrent constitutional symptoms, diarrhea, and on approved medication(s), or, with evidence of depression or memory loss with employment limitations.  Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count  NOTE (1): The term "approved medication(s)" includes medications prescribed as part of a research protocol at an accredited medical institution.  NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated sep arately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  34 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of the signs and symptoms.  Which are nearly constant and restric		10
As active disease Thereafter rate residuals such as arthritis under the appropriate system 320 Parasitic diseases otherwise not specified: As active diseases Thereafter rate residuals such as spleen or liver damage under the appropriate system 350 Lupus erythematosus, systemic (disseminated): Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health Exacerbations lasting a week or more, 2 or 3 times per year Exacerbations lasting a week or more, 2 or 3 times per year Exacerbations once or twice a year or symptomatic during the past 2 years Notre: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating 0.6 6350, whichever method results in a higher evaluation. 351 HIV-Related Illness: AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions		
Thereafter rate residuals such as arthritis under the appropriate system 320 Parasitic diseases otherwise not specified: As active disease  Thereafter rate residuals such as spleen or liver damage under the appropriate system 350. Lupus erythematosus, systemic (disseminated): Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health  Exacerbations lasting a week or more, 2 or 3 times per year  Exacerbations once or twice a year or symptomatic during the past 2 years  Notte: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation. 31 HIV-Related Illness: AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions.  Refractory constitutional symptoms, diarrhea, and pathological weight loss, or; minimum rating following development of AIDS-related opportunistic infection or neoplasm.  Recurrent constitutional symptoms, diarrhea, and pathological weight loss, or; minimum rating with 14 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidiasis.  Following development of definite medical symptoms, 14 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations.  Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count.  NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated sep arately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  354 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combin		10
As active diseases otherwise not specified: As active diseases Thereafter rate residuals such as spleen or liver damage under the appropriate system  350 Lupus erythematosus, systemic (disseminated): Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health Exacerbations lasting a week or more, 2 or 3 times per year Exacerbations once or twice a year or symptomatic during the past 2 years  Notre: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  351 HIV-Related Illness: AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions.  Refractory constitutional symptoms, intermittent diarrhea, and on approved medication(s), or; minimum rating with T4 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidiasis Following development of definite medical symptoms, T4 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations.  Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count  NOTE (1): The term "approved medication(s)" includes medications prescribed as part of a research protocol at an accredited medical institution.  NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated sep arately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  354 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of the real signs and symptoms:  Which are nearly constant and restrict routine dail		10
As active disease Thereafter rate residuals such as spleen or liver damage under the appropriate system  350 Lupus erythematosus, systemic (disseminated): Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health Exacerbations lasting a week or more, 2 or 3 times per year Exacerbations once or twice a year or symptomatic during the past 2 years  NOTE: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  351 HIV-Related Illness: AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions.  Refractory constitutional symptoms, diarrhea, and pathological weight loss, or; minimum rating with T4 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidiasis  Following development of definite medical symptoms, T4 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations.  Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count.  NOTE (1): The term "approved medication(s)" includes medications prescribed as part of a research protocol at an accredite medical institution.  NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated separately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  354 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level,		
Thereafter rate residuals such as spleen or liver damage under the appropriate system  100 Lupus erythematosus, systemic (disseminated):  101 Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health  102 Exacerbations lasting a week or more, 2 or 3 times per year  103 Exacerbations once or twice a year or symptomatic during the past 2 years  104 Note: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  105 HIV-Related Illness:  106 Allow with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions  103 Refractory constitutional symptoms, diarrhea, and pathological weight loss, or; minimum rating of llowing development of AIDS-related opportunistic infection or neoplasm  105 Recurrent constitutional symptoms, intermittent diarrhea, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations  107 Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count  108 Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count  109 Asymptomatic or central nervous system manifestations, opportunistic infections, and neoplasms may be rated sep arately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  108 Deblitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of the resigns and symptoms:  109 Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four b		10
Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health  Exacerbations lasting a week or more, 2 or 3 times per year  Exacerbations once or twice a year or symptomatic during the past 2 years  NOTE: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  31 HIV-Related Illness:  AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions.  Refractory constitutional symptoms, diarrhea, and pathological weight loss, or; minimum rating following development of AIDS-related opportunistic infection or neoplasm  Recurrent constitutional symptoms, intermittent diarrhea, and on approved medication(s), or; minimum rating with T4 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidasis  Following development of definite medical symptoms, T4 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations  Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count less than 200, or HIV infection, with or without lymphadenopathy or decreased T4 cell count less than 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations  Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count less than 500, or with evidence of depression or memory loss with employment limitations.  NOTE (1): The term "approved medication(s)" includes medications prescribed as part of a research protocol at an accredited medical institution.  NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infectio		10
Not to be combined with ratings under DC 7809 Acute, with frequent exacerbations, producing severe impairment of health  Exacerbations lasting a week or more, 2 or 3 times per year  Exacerbations once or twice a year or symptomatic during the past 2 years  NOTE: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  311 HIV-Related Illness:  AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions		
ment of health  Exacerbations lasting a week or more, 2 or 3 times per year  Exacerbations once or twice a year or symptomatic during the past 2 years  NOTE: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  351 HIV-Related Illness:  AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions.  Refractory constitutional symptoms, diarrhea, and pathological weight loss, or; minimum rating following development of AIDS-related opportunistic infection or neoplasm.  Recurrent constitutional symptoms, intermittent diarrhea, and on approved medication(s), or; minimum rating with T4 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidiasis  Following development of definite medical symptoms, T4 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations.  Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count  NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated sep arately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  34 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and os severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year or, which wax and wane, resulting in periods of incapacitation of at least two but		
Exacerbations lasting a week or more, 2 or 3 times per year  Exacerbations once or twice a year or symptomatic during the past 2 years  NOTE: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  351 HIV-Related Illness:  AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions		10
Exacerbations once or twice a year or symptomatic during the past 2 years  NOTE: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  351 HIV-Related Illness:  AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions  Refractory constitutional symptoms, diarrhea, and pathological weight loss, or; minimum rating following development of AIDS-related opportunistic infection or neoplasm		6
Note: Evaluate this condition either by combining the evaluations for residuals under the appropriate system, or by evaluating DC 6350, whichever method results in a higher evaluation.  31 HIV-Related Illness:  AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions		1
ating DC 6350, whichever method results in a higher evaluation.  351 HIV-Related Illness:  AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions	· · · · · · · · · · · · · · · · · · ·	
AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV-related illness with debility and progressive weight loss, without remission, or few or brief remissions		or by evalu-
AIDS with recurrent opportunistic infections or with secondary diseases afflicting multiple body systems; HIV- related illness with debility and progressive weight loss, without remission, or few or brief remissions		
related illness with debility and progressive weight loss, without remission, or few or brief remissions		
Refractory constitutional symptoms, diarrhea, and pathological weight loss, or; minimum rating following development of AIDS-related opportunistic infection or neopleam  Recurrent constitutional symptoms, intermittent diarrhea, and on approved medication(s), or; minimum rating with T4 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidiasis  Following development of definite medical symptoms, T4 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations  Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count  NOTE (1): The term "approved medication(s)" includes medications prescribed as part of a research protocol at an accredited medical institution.  NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated separately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  354 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, r		
velopment of AIDS-related opportunistic infection or neoplasm  Recurrent constitutional symptoms, intermittent diarrhea, and on approved medication(s), or; minimum rating with T4 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidiasis  Following development of definite medical symptoms, T4 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations		10
Recurrent constitutional symptoms, intermittent diarrhea, and on approved medication(s), or; minimum rating with T4 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidiasis  Following development of definite medical symptoms, T4 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations  Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count  NOTE (1): The term "approved medication(s)" includes medications prescribed as part of a research protocol at an accred ited medical institution.  NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated sep arately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  354 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 per		_
with T4 cell count less than 200, or Hairy Cell Leukoplakia, or Oral Candidiasis  Following development of definite medical symptoms, T4 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations		6
Following development of definite medical symptoms, T4 cell of 200 or more and less than 500, and on approved medication(s), or; with evidence of depression or memory loss with employment limitations		
proved medication(s), or; with evidence of depression or memory loss with employment limitations		3
Asymptomatic, following initial diagnosis of HIV infection, with or without lymphadenopathy or decreased T4 cell count  NOTE (1): The term "approved medication(s)" includes medications prescribed as part of a research protocol at an accredited medical institution.  NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated separately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  354 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane presulting in periods of incapacitation of at least two but less than four weeks total duration per year.		4
cell count  NOTE (1): The term "approved medication(s)" includes medications prescribed as part of a research protocol at an accredited medical institution.  NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated separately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  354 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care		'
Note (1): The term "approved medication(s)" includes medications prescribed as part of a research protocol at an accred ited medical institution.  Note (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated sep arately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  354 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year  Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which are nearly constant and restrict routine daily activities to less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year.  Which are nearly constant and restrict routine daily activities to less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year.		
ited medical institution.  NOTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated sep arately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  354 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year  Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year.  Which are nearly constant and restrict routine daily activities to so to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year, or; symptoms controlled by continuous medication.		
NoTE (2): Psychiatric or central nervous system manifestations, opportunistic infections, and neoplasms may be rated sep arately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  354 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year  Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which wax and wane but result in periods of incapacitation of at least one but less than two weeks total duration per year, or; symptoms controlled by continuous medication		an accred-
arately under appropriate codes if higher overall evaluation results, but not in combination with percentages otherwise assignable above.  354 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year  Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year, or; symptoms controlled by continuous medication		rated con
assignable above.  354 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year  Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which are nearly constant and restrict routine daily activities to less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which are nearly constant and restrict routine daily activities almost completely and which was and wane to restrict routine daily activities almost completely and which was and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year, or; symptoms controlled by continuous medication		
254 Chronic Fatigue Syndrome (CFS):  Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year  Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which wax and wane but result in periods of incapacitation of at least one but less than two weeks total duration per year, or; symptoms controlled by continuous medication		5 Offici wise
Debilitating fatigue, cognitive impairments (such as inability to concentrate, forgetfulness, confusion), or a combination of other signs and symptoms:  Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year  Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which wax and wane but result in periods of incapacitation of at least one but less than two weeks total duration per year, or; symptoms controlled by continuous medication		
other signs and symptoms:  Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year  Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which wax and wane but result in periods of incapacitation of at least two but less than two weeks total duration per year, or; symptoms controlled by continuous medication		nhination of
Which are nearly constant and so severe as to restrict routine daily activities almost completely and which may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year  Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which wax and wane but result in periods of incapacitation of at least one but less than two weeks total duration per year, or; symptoms controlled by continuous medication		iibiiialioii o
may occasionally preclude self-care  Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year  Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which wax and wane but result in periods of incapacitation of at least one but less than two weeks total duration per year, or; symptoms controlled by continuous medication		
Which are nearly constant and restrict routine daily activities to less than 50 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year.  Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year.  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year.  Which wax and wane but result in periods of incapacitation of at least two but less than two weeks total duration per year, or; symptoms controlled by continuous medication.		10
or; which wax and wane, resulting in periods of incapacitation of at least six weeks total duration per year Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year  Which wax and wane but result in periods of incapacitation of at least one but less than two weeks total duration per year, or; symptoms controlled by continuous medication		10
Which are nearly constant and restrict routine daily activities to 50 to 75 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year.  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year.  Which wax and wane but result in periods of incapacitation of at least one but less than two weeks total duration per year, or; symptoms controlled by continuous medication		6
which wax and wane, resulting in periods of incapacitation of at least four but less than six weeks total duration per year.  Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year.  Which wax and wane but result in periods of incapacitation of at least one but less than two weeks total duration per year, or; symptoms controlled by continuous medication.		· ·
ration per year		
Which are nearly constant and restrict routine daily activities by less than 25 percent of the pre-illness level, or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year.  Which wax and wane but result in periods of incapacitation of at least one but less than two weeks total duration per year, or; symptoms controlled by continuous medication		4
or; which wax and wane, resulting in periods of incapacitation of at least two but less than four weeks total duration per year		7
duration per year		
Which wax and wane but result in periods of incapacitation of at least one but less than two weeks total duration per year, or; symptoms controlled by continuous medication		2
ration per year, or, symptoms controlled by continuous medication		2
		1
NOTE: For the purpose of evaluating this disability, the condition will be considered incapacitating only while it requires be	Note: For the purpose of evaluating this disability, the condition will be considered incapacitating only while it re	

[61 FR 39875, July 31, 1996]

#### § 4.88c

#### §4.88c Ratings for inactive nonpulmonary tuberculosis initially entitled after August 19, 1968.

	Rat- ing
For 1 year after date of inactivity, following active tu- berculosis	100
Thereafter: Rate residuals under the specific body system or systems affected.	

Following the total rating for the 1 year period after date of inactivity, the schedular evaluation for residuals of nonpulmonary tuberculosis, i.e., ankylosis, surgical removal of a part, etc., will be assigned under the appropriate diagnostic code for the residual preceded by the diagnostic code for tuberculosis of the body part affected. For example, tuberculosis of the hip joint with residual ankylosis would be coded 5001–5250. Where there are existing residuals of pulmonary and nonpulmonary conditions, the evaluations for residual separate functional impairment may be combined.

functional impairment may be combined. Where there are existing pulmonary and nonpulmonary conditions, the total rating for the 1 year, after attainment of inactivity, may not be applied to both conditions during the same period. However, the total rating during the 1-year period for the pulmonary or for the nonpulmonary condition will be utilized, combined with evaluation for residuals of the condition not covered by the 1-year total evaluation, so as to allow any additional benefit provided during such period.

 $[34\ FR\ 5062,\ Mar.\ 11,\ 1969.\ Redesignated\ at\ 59\ FR\ 60902,\ Nov.\ 29,\ 1994]$ 

#### § 4.89 Ratings for inactive nonpulmonary tuberculosis in effect on August 19, 1968.

Public Law 90-493 repealed section 356 of title 38, United States Code which provided graduated ratings for inactive tuberculosis. The repealed section, however, still applies to the case of any veteran who on August 19, 1968, was receiving or entitled to receive compensation for tuberculosis. The use of the protective provisions of Pub. L. 90-493 should be mentioned in the discussion portion of all ratings in which these provisions are applied. For use in rating cases in which the protective provisions of Pub. L. 90-493 apply, the former evaluations are retained in this section.

	Rat- ing
For 2 years after date of inactivity, following active tuberculosis, which was clinically identified during	
service or subsequently	100
Thereafter, for 4 years, or in any event, to 6 years after date of inactivity	50
Thereafter, for 5 years, or to 11 years after date of inactivity	30
Thereafter, in the absence of a schedular compen-	
sable permanent residual	0

Following the total rating for the 2-year period after date of inactivity, the schedular evaluation for residuals of nonpulmonary tuberculosis, i.e., ankylosis, surgical removal of a part, etc., if in excess of 50 percent or 30 percent will be assigned under the appropriate diagnostic code for the specific residual preceded by the diagnostic code for tuberculosis of the body part affected. For example, tuberculosis of the hipjoint with residual ankylosis would be coded 5001–5250.

The graduated ratings for nonpulmonary tuberculosis will not be combined with residuals of nonpulmonary tuberculosis unless the graduated rating and the rating for residual disability cover separate functional losses, e.g., graduated ratings for tuberculosis of the kidney and residuals of tuberculosis of the spine. Where there are existing pulmonary and nonpulmonary conditions, the graduated evaluation for the pulmonary, or for the nonpulmonary, condition will be utilized, combined with evaluations for residuals of the condition not covered by the graduated evaluation utilized, so as to provide the higher evaluation over such period.

The ending dates of all graduated ratings of nonpulmonary tuberculosis will be controlled by the date of attainment of inactivity.

These ratings are applicable only to veterans with nonpulmonary tuberculosis active on or after October 10, 1949.

[29 FR 6718, May 22, 1964, as amended at 34 FR 5062, Mar. 11, 1969; 43 FR 45361, Oct. 2, 1978]

THE RESPIRATORY SYSTEM

# § 4.96 Special provisions regarding evaluation of respiratory conditions.

(a) Rating coexisting respiratory conditions. Ratings under diagnostic codes 6600 through 6817 and 6822 through 6847 will not be combined with each other. Where there is lung or pleural involvement, ratings under diagnostic codes 6819 and 6820 will not be combined with each other or with diagnostic codes 6600 through 6817 or 6822 through 6847. A single rating will be assigned under the diagnostic code which reflects the predominant disability with elevation to the next higher evaluation where the severity of the overall disability warrants such elevation. However, in cases protected by the provisions of Pub. L. 90-493, the graduated ratings of 50 and 30 percent for inactive tuberculosis will not be elevated.

(b) Rating "protected" tuberculosis cases. Public Law 90–493 repealed section 356 of title 38, United States Code which had provided graduated ratings for inactive tuberculosis. The repealed

section, however, still applies to the case of any veteran who on August 19, 1968, was receiving or entitled to receive compensation for tuberculosis. The use of the protective provisions of Pub. L. 90–493 should be mentioned in the discussion portion of all ratings in which these provisions are applied. For application in rating cases in which the protective provisions of Pub. L. 90–493 apply the former evaluations pertaining to pulmonary tuberculosis are retained in §4.97.

(c) Special monthly compensation. When evaluating any claim involving complete organic aphonia, refer to \$3.350 of this chapter to determine

whether the veteran may be entitled to special monthly compensation. Footnotes in the schedule indicate conditions which potentially establish entitlement to special monthly compensation; however, there are other conditions in this section which under certain circumstances also establish entitlement to special monthly compensation.

(Authority: 38 U.S.C. 1155)

[34 FR 5062, Mar. 11, 1969, as amended at 61 FR 46727, Sept. 5, 1996]

# §4.97 Schedule of ratings—respiratory system.

	Rating
DISEASES OF THE NOSE AND THROAT	
6502 Septum, nasal, deviation of:	
Traumatic only, With 50-percent obstruction of the nasal passage on both sides or complete obstruction on one side	10
6504 Nose, loss of part of, or scars:	0.0
Exposing both nasal passages	30
Note: Or evaluate as DC 7800, scars, disfiguring, head, face, or neck.	
6510 Sinusitis, pansinusitis, chronic. 6511 Sinusitis, ethmoid, chronic. 6512 Sinusitis, frontal, chronic. 6513 Sinusitis, maxillary, chronic. 6514 Sinusitis, sphenoid, chronic. General Rating Formula for Sinusitis (DC's 6510 through 6514):	
Following radical surgery with chronic osteomyelitis, or; near constant sinusitis characterized by headaches, pain and tenderness of affected sinus, and purulent discharge or crusting after repeated surgeries	50
biotic treatment, or; more than six non-incapacitating episodes per year of sinusitis characterized by head- aches, pain, and purulent discharge or crusting	30
One or two incapacitating episodes per year of sinusitis requiring prolonged (lasting four to six weeks) anti- biotic treatment, or; three to six non-incapacitating episodes per year of sinusitis characterized by head-	_
aches, pain, and purulent discharge or crusting	10
Note: An incapacitating episode of sinusitis means one that requires bed rest and treatment by a physician.	
6515 Laryngitis, tuberculous, active or inactive.  Rate under §§ 4.88c or 4.89, whichever is appropriate. 6516 Laryngitis, chronic:	
Hoarseness, with thickening or nodules of cords, polyps, submucous infiltration, or pre-malignant changes on bi-	
opsy Hoarseness, with inflammation of cords or mucous membrane	30
6518 Laryngectomy, total	110
Rate the residuals of partial laryngectomy as laryngitis (DC 6516), aphonia (DC 6519), or stenosis of larynx (DC 6520).	
6519 Aphonia, complete organic:  Constant inability to communicate by speech	1100
Constant inability to speak above a whisper	6
Note: Evaluate incomplete aphonia as laryngitis, chronic (DC 6516).	
6520 Larynx, stenosis of, including residuals of laryngeal trauma (unilateral or bilateral):  Forced expiratory volume in one second (FEV–1) less than 40 percent of predicted value, with Flow-Volume Loop compatible with upper airway obstruction, or; permanent tracheostomy  FEV–1 of 40- to 55-percent predicted, with Flow-Volume Loop compatible with upper airway obstruction  FEV–1 of 56- to 70-percent predicted, with Flow-Volume Loop compatible with upper airway obstruction  FEV–1 of 71- to 80-percent predicted, with Flow-Volume Loop compatible with upper airway obstruction	10 6 3
Note: Or evaluate as aphonia (DC 6519).	
6521 Pharynx, injuries to: Stricture or obstruction of pharynx or nasopharynx, or; absence of soft palate secondary to trauma, chemical burn, or granulomatous disease, or; paralysis of soft palate with swallowing difficulty (nasal regurgitation) and speech	5
impairment	

NECO Allumina and a district	Ratin
S522 Allergic or vasomotor rhinitis:  With polyps	;
Without polyps, but with greater than 50-percent obstruction of nasal passage on both sides or complete obstruction on one side	
S523 Bacterial rhinitis:	
Rhinoscleroma  With permanent hypertrophy of turbinates and with greater than 50-percent obstruction of nasal passage on both sides or complete obstruction on one side	
524 Granulomatous rhinitis: Wegener's granulomatosis, lethal midline granuloma Other types of granulomatous infection	10
DISEASES OF THE TRACHEA AND BRONCHI	
6600 Bronchitis, chronic:	
FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy	10
FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)	:
FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent predicted	
601 Bronchiectasis: With incapacitating episodes of infection of at least six weeks total duration per year	1
usage almost continuously  With incapacitating episodes of infection of two to four weeks total duration per year, or; daily productive cough with sputum that is at times purulent or blood-tinged and that requires prolonged (lasting four to six weeks) antibiotic usage more than twice a year	
Intermittent productive cough with acute infection requiring a course of antibiotics at least twice a year  Or rate according to pulmonary impairment as for chronic bronchitis (DC 6600).	
Note: An incapacitating episode is one that requires bedrest and treatment by a physician.	
602 Asthma, bronchial:  FEV-1 less than 40-percent predicted, or; FEV-1/FVC less than 40 percent, or; more than one attack per week with episodes of respiratory failure, or; requires daily use of systemic (oral or parenteral) high dose corticosteroids or immuno-suppressive medications	1
FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; at least monthly visits to a physician for required care of exacerbations, or; intermittent (at least three per year) courses of systemic (oral or parenteral) corticosteroids	
FEV-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; daily inhalational or oral broncho- dilator therapy, or; inhalational anti-inflammatory medication	
FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; intermittent inhalational or oral bron- chodilator therapy	
<b>Note:</b> In the absence of clinical findings of asthma at time of examination, a verified history of asthmatic attacks must be of record.	
Emphysema, pulmonary:  FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy.  FEV-1/FVC of 40 to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent	1
predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)	
FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent predicted	
604 Chronic obstructive pulmonary disease:  FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; episode(s) of acute respiratory failure, or; requires outpatient oxygen therapy.	1
FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)	

	Rating
FEV-1 of 56- to 70-percent predicted, or; FEV-1/FVC of 56 to 70 percent, or; DLCO (SB) 56- to 65-percent pre-	
dicted	30
dicted	10
DISEASES OF THE LUNGS AND PLEURA—TUBERCULOSIS Ratings for Pulmonary Tuberculosis Entitled on August 19, 1968	
6701 Tuberculosis, pulmonary, chronic, far advanced, active	100 100
6703 Tuberculosis, pulmonary, chronic, minimal, active	100
6704 Tuberculosis, pulmonary, chronic, active, advancement unspecified	100
6722 Tuberculosis, pulmonary, chronic, nar advanced, inactive.	
6723 Tuberculosis, pulmonary, chronic, minimal, inactive.	
6724 Tuberculosis, pulmonary, chronic, inactive, advancement unspecified.  General Rating Formula for Inactive Pulmonary Tuberculosis: For two years after date of inactivity, following active	
tuberculosis, which was clinically identified during service or subsequently	100
Thereafter for four years, or in any event, to six years after date of inactivity	50 30
Following far advanced lesions diagnosed at any time while the disease process was active, minimum	30
Following moderately advanced lesions, provided there is continued disability, emphysema, dyspnea on exertion,	00
impairment of health, etc	20 0
Note (1): The 100-percent rating under codes 6701 through 6724 is not subject to a requirement of precedent hospital treatment. It will be reduced to 50 percent for failure to submit to examination or to follow prescribed treatment upon	Ü
report to that effect from the medical authorities. When a veteran is placed on the 100-percent rating for inactive tu- berculosis, the medical authorities will be appropriately notified of the fact, and of the necessity, as given in footnote	
1 to 38 U.S.C. 1156 (and formerly in 38 U.S.C. 356, which has been repealed by Public Law 90-493), to notify the	
Adjudication Division in the event of failure to submit to examination or to follow treatment.  Note (2): The graduated 50-percent and 30-percent ratings and the permanent 30 percent and 20 percent ratings for	
inactive pulmonary tuberculosis are not to be combined with ratings for other respiratory disabilities. Following	
thoracoplasty the rating will be for removal of ribs combined with the rating for collapsed lung. Resection of the ribs	
incident to thoracoplasty will be rated as removal.	
Ratings for Pulmonary Tuberculosis Initially Evaluated After August 19, 1968	
6730 Tuberculosis, pulmonary, chronic, active	100
<b>Note:</b> Active pulmonary tuberculosis will be considered permanently and totally disabling for non-service-connected pension purposes in the following circumstances:	
(a) Associated with active tuberculosis involving other than the respiratory system.	
(b) With severe associated symptoms or with extensive cavity formation.	
(c) Reactivated cases, generally.     (d) With advancement of lesions on successive examinations or while under treatment.	
(e) Without retrogression of lesions or other evidence of material improvement at the end of six months hos-	
pitalization or without change of diagnosis from "active" at the end of 12 months hospitalization. Material improvement means lessening or absence of clinical symptoms, and X-ray findings of a stationary or retro-	
gressive lesion.	
6731 Tuberculosis, pulmonary, chronic, inactive:	
Depending on the specific findings, rate residuals as interstitial lung disease, restrictive lung disease, or, when ob-	
structive lung disease is the major residual, as chronic bronchitis (DC 6600). Rate thoracoplasty as removal of ribs under DC 5297.	
Note: A mandatory examination will be requested immediately following notification that active tuberculosis evaluated under DC 6730 has become inactive. Any change in evaluation will be carried out under the provisions of	
§ 3.105(e).  6732 Pleurisy, tuberculous, active or inactive: Rate under §§ 4.88c or 4.89, whichever is appropriate.	
NONTUBERCULOUS DISEASES	
6817 Pulmonary Vascular Disease:	
Primary pulmonary hypertension, or; chronic pulmonary thromboembolism with evidence of pulmonary hypertension, right ventricular hypertrophy, or cor pulmonale, or; pulmonary hypertension secondary to other obstructive disease of pulmonary arteries or veins with evidence of right ventricular hypertrophy or cor pulmonale	100
Chronic pulmonary thromboembolism requiring anticoagulant therapy, or; following inferior vena cava surgery with-	100
out evidence of pulmonary hypertension or right ventricular dysfunction	60
Symptomatic, following resolution of acute pulmonary embolism	30 0
Note: Evaluate other residuals following pulmonary embolism under the most appropriate diagnostic code, such	J
as chronic bronchitis (DC 6600) or chronic pleural effusion or fibrosis (DC 6844), but do not combine that eval- uation with any of the above evaluations.	
6819 Neoplasms, malignant, any specified part of respiratory system exclusive of skin growths	100

-		Rating
N	<b>lote:</b> A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals.	
6820	Neoplasms, benign, any specified part of respiratory system. Evaluate using an appropriate respiratory analogy.	
	Bacterial Infections of the Lung	
6822	Actinomycosis.	
	Nocardiosis.	
	Chronic lung abscess.	
	ieneral Rating Formula for Bacterial Infections of the Lung (diagnostic codes 6822 through 6824):  Active infection with systemic symptoms such as fever, night sweats, weight loss, or hemoptysisepending on the specific findings, rate residuals as interstitial lung disease, restrictive lung disease, or, when obstructive lung disease is the major residual, as chronic bronchitis (DC 6600).	100
	Interstitial Lung Disease	
6825	Diffuse interstitial fibrosis (interstitial pneumonitis, fibrosing alveolitis).	
	Desquamative interstitial pneumonitis.	
	Pulmonary alveolar proteinosis.	
	Eosinophilic granuloma of lung.	
	Drug-induced pulmonary pneumonitis and fibrosis.	
6831	Radiation-induced pulmonary pneumonitis and fibrosis.  Hypersensitivity pneumonitis (extrinsic allergic alveolitis).	
	Pneumoconiosis (silicosis, anthracosis, etc.).	
	Asbestosis.	
G	eneral Rating Formula for Interstitial Lung Disease (diagnostic codes 6825 through 6833):	
	Forced Vital Capacity (FVC) less than 50-percent predicted, or; Diffusion Capacity of the Lung for Carbon	
	Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise	
	capacity less than 15 ml/kg/min oxygen consumption with cardiorespiratory limitation, or; cor pulmonale or pulmonary hypertension, or; requires outpatient oxygen therapy	100
	FVC of 50- to 64-percent predicted, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum exercise ca-	100
	pacity of 15 to 20 ml/kg/min oxygen consumption with cardiorespiratory limitation	60
	FVC of 65- to 74-percent predicted, or; DLCO (SB) of 56- to 65-percent predicted	
	,	30
	FVC of 75- to 80-percent predicted, or; DLCO (SB) of 66- to 80-percent predicted	30 10
	FVC of 75- to 80-percent predicted, or; DLCO (SB) of 66- to 80-percent predicted	
6835	FVC of 75- to 80-percent predicted, or; DLCO (SB) of 66- to 80-percent predicted	
6835 6836	FVC of 75- to 80-percent predicted, or; DLCO (SB) of 66- to 80-percent predicted	
6835 6836 6837	FVC of 75- to 80-percent predicted, or; DLCO (SB) of 66- to 80-percent predicted	
6835 6836 6837 6838	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis.	
6835 6836 6837 6838 6839	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis.	
6835 6836 6837 6838 6839	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis.	10
6835 6836 6837 6838 6839	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis. Hucormycosis. Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis Chronic pulmonary mycosis requiring suppressive therapy with no more than minimal symptoms such as oc-	10
6835 6836 6837 6838 6839	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis. Hucormycosis. Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis	100
6835 6836 6837 6838 6839	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis. Hung Disease (diagnostic codes 6834 through 6839): Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis Chronic pulmonary mycosis requiring suppressive therapy with no more than minimal symptoms such as occasional minor hemoptysis or productive Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive	100
6835 6836 6837 6838 6839	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis. Mucormycosis. Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis casional minor hemoptysis or productive cough  Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough  Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough  Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough	100 50 30
6835 6836 6837 6838 6839 G	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis. Mucormycosis. Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis casional minor hemoptysis or productive cough Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough Healed and inactive mycotic lesions, asymptomatic	100 50 30
6835 6836 6837 6838 6839	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis. Identify a fating Formula for Mycotic Lung Disease (diagnostic codes 6834 through 6839): Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis. Chronic pulmonary mycosis requiring suppressive therapy with no more than minimal symptoms such as occasional minor hemoptysis or productive cough. Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough. Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough Healed and inactive mycotic lesions, asymptomatic	100
6835 6836 6837 6838 6839 G	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis. Eineral Rating Formula for Mycotic Lung Disease (diagnostic codes 6834 through 6839): Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis Chronic pulmonary mycosis requiring suppressive therapy with no more than minimal symptoms such as occasional minor hemoptysis or productive cough Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough Healed and inactive mycotic lesions, asymptomatic	100 50 30
6835 6836 6837 6838 6839 G	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis. Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis	100 50 30
6835 6836 6837 6838 6839 G	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis. Seneral Rating Formula for Mycotic Lung Disease (diagnostic codes 6834 through 6839): Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis	100 50 30
6835 6836 6837 6838 6839 G	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis. Beneral Rating Formula for Mycotic Lung Disease (diagnostic codes 6834 through 6839): Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis Chronic pulmonary mycosis requiring suppressive therapy with no more than minimal symptoms such as occasional minor hemoptysis or productive cough Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough Healed and inactive mycotic lesions, asymptomatic lote: Coccidioidomycosis has an incubation period up to 21 days, and the disseminated phase is ordinarily manifest within six months of the primary phase. However, there are instances of dissemination delayed up to many years after the initial infection which may have been unrecognized. Accordingly, when service connection is under consideration in the absence of record or other evidence of the disease in service, service in southwestern United States where the disease is endemic and absence of prolonged residence in this locality before or after service will be the deciding factor.  Restrictive Lung Disease  Diaphragm paralysis or paresis. Spinal cord injury with respiratory insufficiency.	100 50 30
6835 6836 6837 6838 6839 G N N	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis Ideneral Rating Formula for Mycotic Lung Disease (diagnostic codes 6834 through 6839): Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis. Chronic pulmonary mycosis requiring suppressive therapy with no more than minimal symptoms such as occasional minor hemoptysis or productive cough. Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough. Healed and inactive mycotic lesions, asymptomatic.  Iote: Coccidioidomycosis has an incubation period up to 21 days, and the disseminated phase is ordinarily manifest within six months of the primary phase. However, there are instances of dissemination delayed up to many years after the initial infection which may have been unrecognized. Accordingly, when service connection is under consideration in the absence of record or other evidence of the disease in service, service in southwestern United States where the disease is endemic and absence of prolonged residence in this locality before or after service will be the deciding factor.  Restrictive Lung Disease  Diaphragm paralysis or paresis. Spinal cord injury with respiratory insufficiency. Kyphoscoliosis, pectus excavatum, pectus carinatum.	100 50 30
6836 6837 6838 6839 G	Mycotic Lung Disease  Histoplasmosis of lung. Coccidioidomycosis. Blastomycosis. Cryptococcosis. Aspergillosis. Mucormycosis. Beneral Rating Formula for Mycotic Lung Disease (diagnostic codes 6834 through 6839): Chronic pulmonary mycosis with persistent fever, weight loss, night sweats, or massive hemoptysis Chronic pulmonary mycosis requiring suppressive therapy with no more than minimal symptoms such as occasional minor hemoptysis or productive cough Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough Chronic pulmonary mycosis with minimal symptoms such as occasional minor hemoptysis or productive cough	100 50 30

	Rating
General Rating Formula for Restrictive Lung Disease (diagnostic codes 6840 through 6845):  FEV-1 less than 40 percent of predicted value, or; the ratio of Forced Expiratory Volume in one second to Forced Vital Capacity (FEV-1/FVC) less than 40 percent, or; Diffusion Capacity of the Lung for Carbon Monoxide by the Single Breath Method (DLCO (SB)) less than 40-percent predicted, or; maximum exercise capacity less than 15 ml/kg/min oxygen consumption (with cardiac or respiratory limitation), or; cor pulmonale (right heart failure), or; right ventricular hypertrophy, or; pulmonary hypertension (shown by Echo or cardiac catheterization), or; epoded(s) of acute respiratory failure, or; requires outpatient oxygen therapy FEV-1 of 40- to 55-percent predicted, or; FEV-1/FVC of 40 to 55 percent, or; DLCO (SB) of 40- to 55-percent predicted, or; maximum oxygen consumption of 15 to 20 ml/kg/min (with cardiorespiratory limit)	100 60 30
FEV-1 of 71- to 80-percent predicted, or; FEV-1/FVC of 71 to 80 percent, or; DLCO (SB) 66- to 80-percent predicted	10
Note (1): A 100-percent rating shall be assigned for pleurisy with empyema, with or without pleurocutaneous fistula, until resolved.	
Note (2): Following episodes of total spontaneous pneumothorax, a rating of 100 percent shall be assigned as of the date of hospital admission and shall continue for three months from the first day of the month after hospital discharge.	
Note (3): Gunshot wounds of the pleural cavity with bullet or missile retained in lung, pain or discomfort on exertion, or with scattered rales or some limitation of excursion of diaphragm or of lower chest expansion shall be rated at least 20-percent disabiling. Disabiling injuries of shoulder girdle muscles (Groups I to IV) shall be separately rated and combined with ratings for respiratory involvement. Involvement of Muscle Group XXI (DC 5321), however, will not be separately rated.	
6846 Sarcoidosis:  Cor pulmonale, or; cardiac involvement with congestive heart failure, or; progressive pulmonary disease with fever,	
night sweats, and weight loss despite treatment	
corticosteroids  Chronic hilar adenopathy or stable lung infiltrates without symptoms or physiologic impairment	30
6847 Sleep Apnea Syndromes (Obstructive, Central, Mixed): Chronic respiratory failure with carbon dioxide retention or cor pulmonale, or; requires tracheostomy	10 5 3

<sup>&</sup>lt;sup>1</sup> Review for entitlement to special monthly compensation under §3.350 of this chapter.

[61 FR 46728, Sept. 5, 1996]

## THE CARDIOVASCULAR SYSTEM

# §§ 4.100—4.103 [Reserved]

#### §4.104 Schedule of ratings—cardiovascular system.

DISEASES OF THE HEART

	Rat- ing
NOTE (1): Evaluate car pulmonals, which is a form of	

NOTE (1): Evaluate cor pulmonale, which is a form of secondary heart disease, as part of the pulmonary condition that causes it.

	Rat- ing
NOTE (2): One MET (metabolic equivalent) is the energy cost of standing quietly at rest and represents an oxygen uptake of 3.5 milliliters per kilogram of body weight per minute. When the level of METs at which dyspnea, fatigue, angina, dizziness, or syncope develops is required for evaluation, and a laboratory determination of METs by exercise testing cannot be done for medical reasons, an estimation by a medical examiner of the level of activity (expressed in METs and supported by specific examples, such as slow stair climbing or shoveling snow) that results in dyspnea, fatigue, angina, dizziness, or syncope may be used.	
heart disease):	
During active infection with valvular heart damage and for three months following cessation of therapy for the active infection	100

# § 4.104

# DISEASES OF THE HEART—Continued

# DISEASES OF THE HEART—Continued

		Rat- ing	Rat- ing
	Thereafter, with valvular heart disease (doc- umented by findings on physical examina- tion and either echocardiogram, Doppler echocardiogram, or cardiac catheteriza- tion) resulting in:		More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ven-
	Chronic congestive heart failure, or; work- load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	100	tricular dysfunction with an ejection frac- tion of 30 to 50 percent
	More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ven-		evidence of cardiac hypertrophy or dilatation on electro-cardiogram, echocardiogram, or X-ray
	tricular dysfunction with an ejection fraction of 30 to 50 percent	60	fatigue, angina, dizziness, or syncope, or; continuous medication required
	Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or, evidence of cardiac hypertrophy or dilatation on electro-cardiogram, echocardiogram, or X-ray	30	7003 Pericardial adhesions: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent
	Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required	10	More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue,
7001	Endocarditis:  For three months following cessation of therapy for active infection with cardiac in-	100	angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent
	volvement	100	Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electro-cardiogram, echocardio-
	sulting in:  Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or;		gram, or X-ray
	left ventricular dysfunction with an ejection fraction of less than 30 percent	100	continuous medication required
	than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ven- tricular dysfunction with an ejection frac-		left ventricular dysfunction with an ejection fraction of less than 30 percent
	tion of 30 to 50 percent	60	heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection frac-
	tion on electrocardiogram, echocardio- gram, or X-ray	30	tion of 30 to 50 percent
7002	fatigue, angina, dizziness, or syncope, or; continuous medication required	10	evidence of cardiac hypertrophy or dilata- tion on electrocardiogram, echocardio- gram, or X-ray
	therapy for active infection with cardiac in- volvement	100	greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required
	sulting in:  Chronic congestive heart failure, or; work-load of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent.	100	NOTE: Evaluate syphilitic aortic aneurysms under DC 7110 (aortic aneurysm).

# **Department of Veterans Affairs**

# § 4.104

# DISEASES OF THE HEART—Continued

	Rat- ing			Rat- ing
7005 Arteriosclerotic heart disease (Coronary artery disease):  With documented coronary artery disease resulting in:  Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea,			More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection frac-	
fatigue, angina, dizziness, or syncope, or, left ventricular dysfunction with an ejection fraction of less than 30 percent	100		tion of 30 to 50 percent	60
of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent	60		gram, or X-ray	30
Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray  Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;	30	7008	continuous medication required	10
NOTE: If nonservice-connected arteriosclerotic heart disease is superimposed on service-connected valvular or other non-arteriosclerotic heart disease, request a medical opinion as to which condition is causing the current signs and symptoms.	10	7010	Supraventricular arrhythmias:  Paroxysmal atrial fibrillation or other supraventricular tachycardia, with more than four episodes per year documented by ECG or Holter monitor  Permanent atrial fibrillation (lone atrial fibril-	30
7006 Myocardial infarction:  During and for three months following myocardial infarction, documented by laboratory tests  Thereafter:	100	7011	lation), or; one to four episodes per year of paroxysmal atrial fibrillation or other supraventricular tachycardia documented by ECG or Holter monitor	10
With history of documented myocardial infarction, resulting in: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection			For indefinite period from date of hospital admission for initial evaluation and medical therapy for a sustained ventricular arrhythmia, or; for indefinite period from date of hospital admission for ventricular aneurysmectomy, or; with an automatic implantable Cardioverter-Defibrillator	
fraction of less than 30 percent	100		(AICD) in place	100
tricular dysfunction with an ejection fraction of 30 to 50 percent	60		More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent	60
tion on electrocardiogram, echocardiogram, or X-ray	30		Workload of greater than 5 METs but not greater than 7 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or, evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardio-	00
continuous medication required	10		gram, or X-ray  Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or;	30
left ventricular dysfunction with an ejection fraction of less than 30 percent	100		continuous medication required	10

# § 4.104

# DISEASES OF THE HEART—Continued

	Rat- ing		Rat- ing
Note: A rating of 100 percent shall be assigned from the date of hospital admission for initial evaluation and medical therapy for a sustained ventricular arrhythmia or for ventricular aneurysmectomy. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of § 3.105(e) of this chapter.  7015 Atrioventricular block:  Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	100 60 30	Note: A rating of 100 percent shall be assigned as of the date of hospital admission for valve replacement. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.  7017 Coronary bypass surgery:  For three months following hospital admission for surgery.  Thereafter:  Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	1000 1000 600 1000
in the absence of other evidence of cardiac disease, is not a disability.  7016 Heart valve replacement (prosthesis):		7011), or atrioventricular block (DC 7015).  Minimum	10
For indefinite period following date of hospital admission for valve replacement	100	(AICD's) under DC 7011. 7019 Cardiac transplantation:	
Thereafter: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection		For an indefinite period from date of hospital admission for cardiac transplantation Thereafter: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea,	100
fraction of less than 30 percent	100	fatigue, angina, dizziness, or syncope, or, left ventricular dysfunction with an ejection fraction of less than 30 percent	100
tricular dysfunction with an ejection frac- tion of 30 to 50 percent	60	than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of 30 to 50 percent	60 30
evidence of cardiac hypertrophy or dilatation on electrocardiogram, echocardiogram, or X-ray  Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatique, angina, dizziness, or syncope, or;	30	NOTE: A rating of 100 percent shall be assigned as of the date of hospital admission for cardiac trans- plantation. One year following discharge, the ap- propriate disability rating shall be determined by mandatory VA examination. Any change in evalua- tion based upon that or any subsequent examina-	

# DISEASES OF THE HEART—Continued

	D-4		·
	Rat- ing		Rat- ing
7020 Cardiomyopathy: Chronic congestive heart failure, or; workload of 3 METs or less results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ventricular dysfunction with an ejection fraction of less than 30 percent	100	If symptomatic, or; for indefinite period from date of hospital admission for surgical correction	100
More than one episode of acute congestive heart failure in the past year, or; workload of greater than 3 METs but not greater than 5 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; left ven-	100	ischemic ulcers or ankle/brachial index of 0.4 or less	100
tricular dysfunction with an ejection frac- tion of 30 to 50 percent	60	brachial index of 0.5 or less	60
tion on electrocardiogram, echocardiogram, or X-ray	30	Claudication on walking more than 100 yards, and; diminished peripheral pulses	40
Workload of greater than 7 METs but not greater than 10 METs results in dyspnea, fatigue, angina, dizziness, or syncope, or; continuous medication required	10	or ankle/brachial index of 0.9 or less  NOTE (1): The ankle/brachial index is the ratio of the systolic blood pressure at the ankle (determined by Doppler study) divided by the simultaneous brachial artery systolic blood pressure. The normal index is 1.0 or greater.	20
Diastolic pressure predominantly 130 or more Diastolic pressure predominantly 120 or more Diastolic pressure predominantly 110 or more, or; systolic pressure predominantly 200 or more Diastolic pressure predominantly 100 or more, or; systolic pressure predominantly 100 or more, or; systolic pressure predominantly 160 or more, or; minimum evaluation for	60 40 20	NOTE (2): These evaluations are for involvement of a single extremity. If more than one extremity is affected, evaluate each extremity separately and combine (under § 4.25), using the bilateral factor, if applicable.  NOTE (3): A rating of 100 percent shall be assigned as of the date of hospital admission for surgical correction. Six months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examina-	
an individual with a history of diastolic pressure predominantly 100 or more who requires continuous medication for control NOTE (1): Hypertension or isolated systolic hyper-	10	tion shall be subject to the provisions of §3.105(e) of this chapter.  7112 Aneurysm, any small artery:	
tension must be confirmed by readings taken two or more times on at least three different days. For purposes of this section, the term hypertension means that the diastolic blood pressure is predominantly 90mm. or greater, and isolated systolic hy-		Asymptomatic	(
pertension means that the systolic blood pressure is predominantly 160mm. or greater with a diastolic blood pressure of less than 90mm.  NOTE (2): Evaluate hypertension due to aortic insuffi-		7113 Arteriovenous fistula, traumatic:  With high output heart failure  Without heart failure but with enlarged heart, wide pulse pressure, and tachycardia  Without cardiac involvement but with	10
ciency or hyperthyroidism, which is usually the iso- lated systolic type, as part of the condition causing it rather than by a separate evaluation.		edema, stasis dermatitis, and either ulceration or cellulitis:	_
110 Aortic aneurysm:  If five centimeters or larger in diameter, or; if		Lower extremity  Upper extremity  With edema or stasis dermatitis:	5 4
symptomatic, or, for indefinite period from date of hospital admission for surgical correction (including any type of graft insertion)  Precluding exertion  Evaluate residuals of surgical correction according to the property of the period o	100 60	Lower extremity	3 2
Evaluate residuals of surgical correction according to organ systems affected.  IOTE: A rating of 100 percent shall be assigned as of the date of admission for surgical correction. Six		0.4 or less	10
months following discharge, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.  111 Aneurysm, any large artery:		or ankle/brachial index of 0.5 or less  Claudication on walking between 25 and 100 yards on a level grade at 2 miles per hour, and; trophic changes (thin skin, absence of hair, dystrophic nails) or ankle/ brachial index of 0.7 or less	6

# § 4.104

# DISEASES OF THE HEART—Continued

	Rat- ing		Rat- ing
Claudication on walking more than 100 yards, and; diminished peripheral pulses or ankle/brachial index of 0.9 or less  NOTE (1): The ankle/brachial index is the ratio of the systolic blood pressure at the ankle (determined by Doppler study) divided by the simultaneous brachial artery systolic blood pressure. The normal index is 1.0 or greater.  NOTE (2): Evaluate residuals of aortic and large arterial bypass surgery or arterial graft as arteriosclerosis obliterans.  NOTE (3): These evaluations are for involvement of a single extremity. If more than one extremity is affected, evaluate each extremity separately and combine (under § 4.25), using the bilateral factor (§ 4.26), if applicable.  7115 Thrombo-angiitis obliterans (Buerger's Disease):	20	Attacks without laryngeal involvement lasting one to seven days and occurring five to eight times a year, or; attacks with laryngeal involvement of any duration occurring once or twice a year	20 10
Ischemic limb pain at rest, and; either deep ischemic ulcers or ankle/brachial index of 0.4 or less	100	Characteristic attacks that occur daily or more often but that respond to treatment Characteristic attacks that occur less than daily but at least three times a week and	30
either persistent coldness of the extremity or ankle/brachial index of 0.5 or less	60	that respond to treatment	10
yards, and; diminished peripheral pulses or ankle/brachial index of 0.9 or less  NOTE (1): The ankle/brachial index is the ratio of the systolic blood pressure at the ankle (determined by Doppler study) divided by the simultaneous brachial artery systolic blood pressure. The normal index is 1.0 or greater.  NOTE (2): These evaluations are for involvement of a single extremity. If more than one extremity is affected, evaluate each extremity separately and combine (under §4.25), using the bilateral factor (§4.26), if applicable.	20	7120 Varicose veins:  With the following findings attributed to the effects of varicose veins: Massive board-like edema with constant pain at rest  Persistent edema or subcutaneous induration, stasis pigmentation or eczema, and persistent ulceration  Persistent edema and stasis pigmentation or eczema, with or without intermittent ulceration  Persistent edema, incompletely relieved by elevation of extremity, with or without beginning stasis pigmentation or eczema	100 60 40
With two or more digital ulcers plus autoamputation of one or more digits and history of characteristic attacks	100 60 40 20 10	Intermittent edema of extremity or aching and fatigue in leg after prolonged standing or walking, with symptoms relieved by elevation of extremity or compression hosiery	10
tacks consist of sequential color changes of the digits of one or more extremities lasting minutes to hours, sometimes with pain and paresthesias, and precipitated by exposure to cold or by emotional upsets. These evaluations are for the disease as a whole, regardless of the number of extremities involved or whether the nose and ears are involved.  7118 Angioneurotic edema:		(§ 4.26), if applicable.  7121 Post-phlebitic syndrome of any etiology: With the following findings attributed to venous disease:  Massive board-like edema with constant pain at rest  Persistent edema or subcutaneous induration, stasis pigmentation or ecrama, and persistent ulcara.	100
Attacks without laryngeal involvement last- ing one to seven days or longer and oc- curring more than eight times a year, or; attacks with laryngeal involvement of any duration occurring more than twice a year	40	eczema, and persistent ulcera- tion	60 40

#### DISEASES OF THE HEART—Continued

Rat-Persistent edema, incompletely relieved by elevation of extremity, with or without beginning stasis pigmentation or eczema ..... Intermittent edema of extremity or aching and fatigue in leg after prolonged standing or walking, with symptoms relieved by elevation of extremity or compression hosiery Asymptomatic palpable or visible varicose veins NOTE: These evaluations are for involvement of a

single extremity. If more than one extremity is involved, evaluate each extremity separately and combine (under § 4.25), using the bilateral factor (§ 4.26), if applicable.

#### 7122 Cold injury residuals:

With the following in affected parts:

Arthralgia or other pain, numbness or cold sensitivity plus two or more of the following: tissue loss, nail abnormalities, color changes, impaired hyperhidrosis, X-ray abnormalities (osteoporosis, subarticular punched out lesions, or osteoarthritis)

Arthralgia or other pain, numbness or cold sensitivity plus tissue loss, nail abnormalities, color changes, locally impaired sensation, hyperhidrosis, or X-ray abnormalities (osteoporosis, articular punched out lesions, or osteoarthritis) ...

Arthralgia or other pain, numbness, or cold sensitivity

NOTE (1): Separately evaluate amputations of fingers or toes, and complications such as squamous cell carcinoma at the site of a cold injury scar or peripheral neuropathy, under other diagnostic codes. Separately evaluate other disabilities that have been diagnosed as the residual effects of cold injury, such as Raynaud's phenomenon, muscle atrophy, etc., unless they are used to support an evaluation under diagnostic code 7122

NOTE (2): Evaluate each affected part (e.g., hand, foot, ear, nose) separately and combine the ratings in accordance with §§ 4.25 and 4.26.

7123 Soft tissue sarcoma (of vascular origin) ...

NOTE: A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals.

(Authority: 38 U.S.C. 1155)

[62 FR 65219, Dec. 11, 1997, as amended at 63 FR 37779, July 14, 1998]

THE DIGESTIVE SYSTEM

#### §4.110 Ulcers.

ing

Experience has shown that the term "peptic ulcer" is not sufficiently specific for rating purposes. Manifest differences in ulcers of the stomach or duodenum in comparison with those at an anastomotic stoma are sufficiently recognized as to warrant two separate graduated descriptions. In evaluating the ulcer, care should be taken that the findings adequately identify the particular location.

#### §4.111 Postgastrectomy syndromes.

There are various postgastrectomy symptoms which may occur following anastomotic operations of the stomach. When present, those occurring during or immediately after eating and known as the "dumping syndrome" are characterized by gastrointestinal complaints and generalized symptoms simulating hypoglycemia; those occurring from 1 to 3 hours after eating usually present definite manifestations of hypoglycemia.

#### §4.112 Weight loss.

Minor weight loss or greater losses of weight for periods of brief duration are not considered of importance in rating. Rather, weight loss becomes of importance where there is appreciable loss which is sustained over a period of time. In evaluating weight loss generally, consideration will be given not only to standard age, height, and weight tables, but also to the parpredominant individual's ticular weight pattern as reflected by the records. The use of the term "inability to gain weight" indicates that there has been a significant weight loss with inability to regain it despite appropriate therapy.

#### §4.113 Coexisting abdominal conditions.

There are diseases of the digestive system, particularly within the abdomen, which, while differing in the site of pathology, produce a common disability picture characterized in the main by varying degrees of abdominal distress or pain, anemia and disturbances in nutrition. Consequently, certain coexisting diseases in this area, as

#### § 4.114

indicated in the instruction under the title "Diseases of the Digestive System," do not lend themselves to distinct and separate disability evaluations without violating the fundamental principle relating pyramiding as outlined in  $\S4.14$ .

#### §4.114 Schedule of ratings—digestive system.

Ratings under diagnostic codes 7301 to 7329, inclusive, 7331, 7342, and 7345 to 7348 inclusive will not be combined with each other. A single evaluation will be assigned under the diagnostic code which reflects the predominant disability picture, with elevation to the next higher evaluation where the severity of the overall disability warrants such elevation.

	Rat- ing
7200 Mouth, injuries of.	
Rate as for disfigurement and impairment of	
function of mastication.	
7201 Lips, injuries of.	
Rate as for disfigurement of face.	
7202 Tongue, loss of whole or part:	400
With inability to communicate by speech	100
One-half or more	30
·	30
7203 Esophagus, stricture of: Permitting passage of liquids only, with marked	
impairment of general health	80
Severe, permitting liquids only	50
Moderate	30
7204 Esophagus, spasm of (cardiospasm).	
If not amenable to dilation, rate as for the de-	
gree of obstruction (stricture).	
7205 Esophagus, diverticulum of, acquired.	
Rate as for obstruction (stricture).	
7301 Peritoneum, adhesions of:	
Severe; definite partial obstruction shown by X-	
ray, with frequent and prolonged episodes of	
severe colic distension, nausea or vomiting,	
following severe peritonitis, ruptured appendix,	
perforated ulcer, or operation with drainage	50
Moderately severe; partial obstruction mani- fested by delayed motility of barium meal and	
less frequent and less prolonged episodes of	
pain	30
Moderate; pulling pain on attempting work or ag-	
gravated by movements of the body, or occa-	
sional episodes of colic pain, nausea, con-	
stipation (perhaps alternating with diarrhea) or	
abdominal distension	10
Mild	0
NOTE: Ratings for adhesions will be considered	
when there is history of operative or other traumatic or infectious (intraabdominal) proc-	
ess, and at least two of the following: disturb-	
ance of motility, actual partial obstruction, re-	
flex disturbances, presence of pain.	
7304 Ulcer, gastric.	
7305 Ulcer, duodenal:	

`	•
	Rat- ing
Severe; pain only partially relieved by standard ulcer therapy, periodic vomiting, recurrent	
hematemesis or melena, with manifestations	
of anemia and weight loss productive of defi-	
nite impairment of health	60
pairment of health manifested by anemia and	
weight loss; or recurrent incapacitating epi-	
sodes averaging 10 days or more in duration	40
at least four or more times a year	40
Moderate; recurring episodes of severe symptoms two or three times a year averaging 10	
days in duration; or with continuous moderate	
manifestations	20
Mild; with recurring symptoms once or twice yearly	10
7306 Ulcer, marginal (gastrojejunal):	
Pronounced; periodic or continuous pain	
unrelieved by standard ulcer therapy with peri-	
odic vomiting, recurring melena or hematemesis, and weight loss. Totally inca-	
pacitating	100
Severe; same as pronounced with less pro-	
nounced and less continuous symptoms with definite impairment of health	60
Moderately severe; intercurrent episodes of ab-	
dominal pain at least once a month partially or	
completely relieved by ulcer therapy, mild and	40
transient episodes of vomiting or melena  Moderate; with episodes of recurring symptoms	40
several times a year	20
Mild; with brief episodes of recurring symptoms	4.0
once or twice yearly	10
scope):	
Chronic; with severe hemorrhages, or large ul-	00
cerated or eroded areas	60
areas, and symptoms	30
Chronic; with small nodular lesions, and symp-	
tomsGastritis, atrophic.	10
A complication of a number of diseases, includ-	
ing pernicious anemia.	
Rate the underlying condition. 7308 Postgastrectomy syndromes:	
Severe; associated with nausea, sweating, cir-	
culatory disturbance after meals, diarrhea,	
hypoglycemic symptoms, and weight loss with	00
malnutrition and anemia	60
disorders with characteristic mild circulatory	
symptoms after meals but with diarrhea and	
weight loss Mild; infrequent episodes of epigastric distress	40
with characteristic mild circulatory symptoms	
or continuous mild manifestations	20
7309 Stomach, stenosis of.	
Rate as for gastric ulcer. 7310 Stomach, injury of, residuals.	
Rate as peritoneal adhesions.	
7311 Liver, injury of.	
With residual disability, rate as peritoneal adhesions.	
Healed, no residuals	C
7312 Liver, cirrhosis of:	
Pronounced; aggravation of the symptoms for	
moderate and severe, necessitating frequent tapping	100
Severe; ascites requiring infrequent tapping, or	100
recurrent hemorrhage from esophageal	
varices, aggravated symptoms and impaired health	70
HEGINI	/()

70

100 60 30

> 80 30

> 50

30

10

	Rat- ing		Ra ing
Moderately severe; liver definitely enlarged with abdominal distention due to early ascites and with muscle wasting and loss of strength	50	With marked interference with absorption and nutrition, manifested by severe impairment of health objectively supported by examination	
Moderate; with dilation of superficial abdominal veins, chronic dyspepsia, slight loss of weight	00	findings including material weight loss	
or impairment of health	30	nutrition, manifested by impairment of health	
313 Liver, abscess of, residuals: With severe symptoms	30	objectively supported by examination findings including definite weight loss	
With moderate symptoms	20	Symptomatic with diarrhea, anemia and inability	
Cholecystitis, chronic:	00	to gain weight	
Severe; frequent attacks of gall bladder colic  Moderate; gall bladder dyspepsia, confirmed by X-ray technique, and with infrequent attacks	30	predominant disability, rate under diagnostic code 7301.	
(not over two or three a year) of gall bladder	4.0	7329 Intestine, large, resection of: With severe symptoms, objectively supported by	
colic, with or without jaundice	10 0	examination findings	
15 Cholelithiasis, chronic.	O	With moderate symptoms	
Rate as for chronic cholecystitis.		With slight symptoms	
16 Cholangitis, chronic.		NOTE: Where residual adhesions constitute the predominant disability, rate under diagnostic	
Rate as for chronic cholecystitis.  17 Gall bladder, injury of.		code 7301.	
Rate as for peritoneal adhesions.		7330 Intestine, fistula of, persistent, or after attempt	
18 Gall bladder, removal of:		at operative closure:  Copious and frequent, fecal discharge	1
With severe symptoms	30	Constant or frequent, fecal discharge	
With mild symptoms  Nonsymptomatic	10 0	Slight infrequent, fecal discharge	
Spleen, disease or injury of.	Ü	Healed; rate for peritoneal adhesions.	
See Hemic and Lymphatic Systems.		7331 Peritonitis, tuberculous, active or inactive:	1
19 Irritable colon syndrome (spastic colitis, mu-		Inactive: See §§ 4.88b and 4.89.	
cous colitis, etc.):  Severe; diarrhea, or alternating diarrhea and		7332 Rectum and anus, impairment of sphincter	
constipation, with more or less constant ab-		control:  Complete loss of sphincter control	
dominal distress	30	Extensive leakage and fairly frequent involuntary	
Moderate; frequent episodes of bowel disturb- ance with abdominal distress	10	bowel movements	
Mild; disturbances of bowel function with occa-	10	Occasional involuntary bowel movements, ne-	
sional episodes of abdominal distress	0	cessitating wearing of pad  Constant slight, or occasional moderate leakage	
21 Amebiasis:		Healed or slight, without leakage	
Mild gastrointestinal disturbances, lower abdominal cramps, nausea, gaseous distention,		7333 Rectum and anus, stricture of:	
chronic constipation interrupted by diarrhea	10	Requiring colostomy  Great reduction of lumen, or extensive leakage	1
Asymptomatic	0	Moderate reduction of lumen, or moderate constant leakage	
tis and should be rated on the scale provided		7334 Rectum, prolapse of: Severe (or complete), persistent	
for the latter. Similarly, lung abscess due to amebiasis will be rated under the respiratory		Moderate, persistent or frequently recurring Mild with constant slight or occasional moderate	
system schedule, diagnostic code 6809.  22 Dysentery, bacillary.		leakage	
Rate as for ulcerative colitis		Rate as for impairment of sphincter control.	
23 Colitis, ulcerative:  Pronounced; resulting in marked malnutrition,		7336 Hemorrhoids, external or internal:	
anemia, and general debility, or with serious		With persistent bleeding and with secondary anemia, or with fissures	
complication as liver abscess	100	Large or thrombotic, irreducible, with excessive	
Severe; with numerous attacks a year and mal- nutrition, the health only fair during remissions	60	redundant tissue, evidencing frequent	
Moderately severe; with frequent exacerbations	30	recurrences	
Moderate; with infrequent exacerbations	10	7337 Pruritus ani.	
24 Distomiasis, intestinal or hepatic:		Rate for the underlying condition.	
Severe symptoms	30 10	7338 Hernia, inguinal:	
Mild or no symptoms	0	Large, postoperative, recurrent, not well sup- ported under ordinary conditions and not read-	
25 Enteritis, chronic.		ily reducible, when considered inoperable	
Rate as for irritable colon syndrome.		Small, postoperative recurrent, or unoperated ir-	
26 Enterocolitis, chronic.  Rate as for irritable colon syndrome.		remediable, not well supported by truss, or not readily reducible	
27 Diverticulitis.		Postoperative recurrent, readily reducible and	
Rate as for irritable colon syndrome, peritoneal		well supported by truss or belt	
adhesions, or colitis, ulcerative, depending upon the predominant disability picture.		Not operated, but remediable	
328 Intestine, small, resection of:		sion	

	Rat- ing
Note: Add 10 percent for bilateral involvement, provided the second hernia is compensable. This means that the more severely disabling hernia is to be evaluated, and 10 percent, only, added for the second hernia, if the latter is of compensable degree.	
7339 Hernia, ventral, postoperative:  Massive, persistent, severe diastasis of recti muscles or extensive diffuse destruction or weakening of muscular and fascial support of abdominal wall so as to be inoperable	10
Large, not well supported by belt under ordinary conditions	4
Small, not well supported by belt under ordinary conditions, or healed ventral hernia or post-operative wounds with weakening of abdominal	
wall and indication for a supporting belt Wounds, postoperative, healed, no disability, belt not indicated	2
7340 Hernia, femoral.	
Rate as for inguinal hernia.  7342 Visceroptosis, symptomatic, marked	1
7343 New growths, malignant, exclusive of skin	
growths	10
NOTE: The rating under diagnostic code 7343 will be continued for 1 year following the cessation of surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. At this point, if there has been no local recurrence or metastases, the rating will be made on residuals.	
7344 New growths, benign, any specified part of di-	
gestive system, exclusive of skin growths.	
The rating will be based on interference with di- gestion, using any applicable digestive anal- ogy.	
7345 Hepatitis, infectious:	
With marked liver damage manifest by liver func- tion test and marked gastrointestinal symp- toms, or with episodes of several weeks dura- tion aggregating three or more a year and ac- companied by disabling symptoms requiring	
rest therapy	10
With moderate liver damage and disabling recurrent episodes of gastrointestinal disturbance,	
fatigue, and mental depression	6
Minimal liver damage with associated fatigue, anxiety, and gastrointestinal disturbance of lesser degree and frequency but necessitating dietary restriction or other therapeutic meas-	
Demonstrable liver damage with mild gastro-	3
intestinal disturbance	1
Healed, nonsymptomatic	
7346 Hernia hiatal:	
Symptoms of pain, vomiting, material weight loss and hematemesis or melena with moderate anemia; or other symptom combinations pro- ductive of severe impairment of health	6
Persistently recurrent epigastric distress with dysphagia, pyrosis, and regurgitation, accom- panied by substernal or arm or shoulder pain,	
productive of considerable impairment of health	3
With two or more of the symptoms for the 30 percent evaluation of less severity	1
7347 Pancreatitis:	

With frequently recurrent disabling attacks of abdominal pain with few pain free intermissions

and severe malnutrition .

and with steatorrhea, malabsorption, diarrhea

	Rat- ing
With frequent attacks of abdominal pain, loss of normal body weight and other findings showing continuing pancreatic insufficiency be-	
tween acute attacks	6
mission between attacks	3
vere abdominal pain in the past year NOTE 1: Abdominal pain in this condition must	1
be confirmed as resulting from pancreatitis by appropriate laboratory and clinical studies.	
NOTE 2: Following total or partial pancreatectomy, rate under above, symptoms, min-	
imum rating 30 percent. 7348 Vagotomy with pyloroplasty or gastro-	
enterostomy: Followed by demonstrably confirmative post- operative complications of stricture or con-	
tinuing gastric retention	4
rhea	3
Recurrent ulcer with incomplete vagotomy  NOTE: Rate recurrent ulcer following complete vagotomy under diagnostic code 7305, minimum rating 20 percent; and rate dumping syndrome under diagnostic code 7308.	2

[29 FR 6718, May 22, 1964, as amended at 34 FR 5063, Mar. 11, 1969; 40 FR 42540, Sept. 15, 1975; 41 FR 11301, Mar. 18, 1976]

#### THE GENITOURINARY SYSTEM

## §4.115 Nephritis.

Albuminuria alone is not nephritis, nor will the presence of transient albumin and casts following acute febrile illness be taken as nephritis. The glomerular type of nephritis is usually preceded by or associated with severe infectious disease; the onset is sudden, and the course marked by red blood cells, salt retention, and edema; it may clear up entirely or progress to a chronic condition. The nephrosclerotic type, originating in hypertension or arteriosclerosis, develops slowly, with minimum laboratory findings, and is associated with natural progress. Separate ratings are not to be assigned for disability from disease of the heart and any form of nephritis, on account of the close interrelationships of cardiovascular disabilities. If, however, absence of a kidney is the sole renal disability, even if removal was required because of nephritis, the absent kidney and any hypertension or heart disease will be separately rated. Also, in the event that chronic renal disease has progressed to the point where regular

dialysis is required, any coexisting hypertension or heart disease will be separately rated.

[41 FR 34258, Aug. 13, 1976, as amended at 59 FR 2527, Jan. 18, 1994]

#### §4.115a Ratings of the genitourinary system-dysfunctions.

Diseases of the genitourinary system generally result in disabilities related to renal or voiding dysfunctions, infections, or a combination of these. The following section provides descriptions of various levels of disability in each of these symptom areas. Where diagnostic codes refer the decisionmaker to these specific areas dysfunction, only the predominant area of dysfunction shall be considered for rating purposes. Since the areas of dysfunction described below do not cover all symptoms resulting from genitourinary diseases, specific diagnoses may include a description of symptoms assigned to that diagnosis.

Renal dysfunction: Requiring regular dialysis, or precluding more than sedentary activity from one of the following: persistent edema and albuminuria; or, BUN more than 80mg%; or, creatinine more than 80mg%; or, markedly decreased function of kidney or other organ systems, estpecially cardiovascular  Persistent edema and albuminuria with BUN 40 to 80mg%; or, creatinine 4 to 8mg%; or, generalized poor health characterized by lethargy, weakness, anorexia, weight loss, or limitation of exertion	
day	

	Rat- ing
Daytime voiding interval between one and two hours, or; awakening to void three to four times	
per night	20
Daytime voiding interval between two and three hours, or; awakening to void two times per night	10
Obstructed voiding:	
Urinary retention requiring intermittent or contin- uous catheterization	30
Marked obstructive symptomatology (hesitancy, slow or weak stream, decreased force of stream) with any one or combination of the following:  1. Post void residuals greater than 150 cc.  2. Uroflowmetry; markedly diminished peak flow rate (less than 10 cc/sec).  3. Recurrent urinary tract infections secondary to	
obstruction.	
Stricture disease requiring periodic dilatation every 2 to 3 months	10
Obstructive symptomatology with or without stric- ture disease requiring dilatation 1 to 2 times per	
year	0
Urninary tract infection:	
Poor renal function: Rate as renal dysfunction.	
Recurrent symptomatic infection requiring drain- age/frequent hospitalization (greater than two times/year), and/or requiring continuous inten- sive management	30
Long-term drug therapy, 1–2 hospitalizations per year and/or requiring intermittent intensive man-	

[59 FR 2527, Jan. 18, 1994; 59 FR 10676, Mar. 7, 1994]

#### §4.115b Ratings of the genitourinary system—diagnoses.

		Rat- ing
	Note: When evaluating any claim involving loss or loss of use of one or more creative organs, refer to	
	§ 3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensa-	
	tion. Footnotes in the schedule indi- cate conditions which potentially establish entitlement to special	
	monthly compensation; however, there are other conditions in this section which under certain circumstances also establish entitlement to special monthly compensation.	
7500	Kidney, removal of one: Minimum evaluation	20
	Or rate as renal dysfunction if there is	30
	nephritis, infection, or pathology of the other.	
7501	Kidney, abscess of: Rate as urinary tract infection	

ing

100

# § 4.115b

7502 Nephritis, chronic: Rate as renal dysfunction. 7504 Pyelonephritis, chronic: Rate as renal dysfunction or urinary tract infection, whichever is predominant. 7505 Kidney, tuberculosis of: Rate in accordance with §§4.88b or 4.89, whichever is appropriate. 7507 Nephrosclerosis, arteriolar: Rate according to predominant symptoms as renal dysfunction, hypertension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation. 7508 Nephrolithiasis: Rate as voiding dysfunction. Multiple urethroperineal fistulae		Rat-		Rat-
Rate as renal dysfunction. 7504 Pyelonephritis, chronic: Rate as renal dysfunction or urinary tract infection, whichever is predominant. 7505 Kidney, tuberculosis of: Rate in accordance with §§ 4.88b or 4.89, whichever is appropriate. 7507 Nephrosclerosis, arteriolar: Rate according to predominant symptoms as renal dysfunction, hypertension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation. 7508 Nephrolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. died therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year charled and not requiring catheter drainage.  Only an occasional attack of colic, requiring one or more of the following: 1. died therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year charled and not requiring catheter drainage and the following: 1. died therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year charled and not requiring catheter drainage and the following: 20 Il therent stone formation requiring one or more of the following: 21 died therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year charled the developed the therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year charled the drained processing the developed thereof the service-connected testicular loss. Testis, underscended, or congenitally undeveloped is not a rational developed testis, with the absence or nonfunctioning of the other testis unrelated to service, an evaluation of 30 percent will be assigned for the service-connected testicular loss. Testis, underscended, or congenitally undeveloped is not a rational development of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year procedures more than two times/year procedures more		ing		ing
Rate as renal dysfunction or urinary tract infection, whichever is predominant.  7505 Kidney, tuberculosis of: Rate in accordance with §§4.88b or 4.89, whichever is appropriate.  7507 Nephrosclerosis, arteriolar: Rate according to predominant symptoms as renal dysfunction, hypertension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation.  7508 Nephrosithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1 diet therapy 2 drug therapy 3. Invasive or non-invasive procedures more than two times/year Only an occasional attack of colic, requiring cahelet drainage Only an occasional attack of colic, requiring one or more of the following: 1. Identerolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. Identerolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. Identerolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. Identerolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. Identerolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. Identerolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. Identerolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. Identerolithiasis: Rate as violiting dysfunction.  7525 Epidigymo-orchitis, chronic only: Rate as volding dysfunction.  7526 Epidigymo-orchitis, chronic only: Rate as volding dysfunction.  7527 Prostate gland injuries, infections, hypertrophy, postoperative residuals: Rate as violing dysfunction.  7528 Malignant neoplasms of the genitary viact infection, whiche				
Rate as renal dysfunction or urinary tract infection, whichever is predominant.  7508 Kidney, tuberculosis of: Rate in accordance with §\$4.88b or 4.89, whichever is appropriate.  7507 Nephrosclerosis, arteriolar: Rate accordance predominant symptoms as renal dysfunction, hypertension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation.  7508 Nephrolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year eter drainage  7510 Ureterolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year drainage  7510 Ureterolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year drainage  7511 Ureter, stricture of: Rate as voiding dysfunction.  7522 Penis, removal of glans  7522 Penis, deformity, with loss of erectile power—201  7523 Testis, attorphy complete:.  80th—201  7524 Testis, attorphy complete:.  80th—201  7525 Testis, attorphy complete:.  80th—301  7526 Neigramoval.  80th—301  7527 Penis removal of glans  7528 Alegramoval.  80th—201  7528 Testis, attorphy complete:.  80th—301  7529 Penis, removal of glans  7520 Penis, removal of glans  7521 Penis removal of glans  7522 Penis, deformity, with loss of erectile power—201  7523 Testis, attorphy complete:.  80th—301  7524 Testis, attorphy complete:.  80th—301  7525 Testis, undersomed of the removal of one testis as the result of a service-incurred injury or disease, other than an undescended or congenitally undeveloped is not a ratable disability.  80th—301  80th—302  80th—303  80th—303  80th—303  80th—304			9 ,	
tract infection, whichever is pre- dominant.  Rate in accordance with § \$4.88b or 4.89, whichever is appropriate.  7507 Nephrosclerosis, arteriolar: Rate according to predominant symp- toms as renal dysfunction, hyper- tension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation.  7508 Nephrolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive proce- dures more than two times/year Only an occasional attack of colic, requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive proce- dures more than two times/year Only an occasional attack of colic, requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive proce- dures more than two times/year Only an occasional attack of colic, requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive proce- dures more than two times/year Total Ureter, fisticule of: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive proce- dures more than two times/year Total Ureter, stricture of: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive proce- dures more than two times/year Total Ureter, stricture of: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive proce- dures more than two times/year Total Ureter, fisticular of: Rate as voiding dysfunction.  7522 Penis, defermity, with loss of erectile power—201.  Note: In cases of the removal of one testis, satiophy complete: Both—201  Note: In cases of the removal			•	
dominant.  dominant whereulosis of: Rate in accordance with § 4.88b or 4.89 whichever is appropriate.  Rate any hichever is appropriate.  Rate accordance yeth § 4.88b or 4.89 whichever is appropriate.  Rate accordance predominant symptoms as renal dysfunction, hypertension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation.  Robert of recurrent stone formation requiring one or more of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year eter drainage  Only an occasional attack of colic, requiring one or more of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year eter drainage  Only an occasional attack of colic, requiring one or more of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year eter drainage  Total Creation of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year eter drainage  Total Creation of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year eter drainage  Total Creation of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year eter drainage  Total Creation of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year eter drainage  Total Creation of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year eter drainage  Total Creation of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year eter drainage  Total Creation of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year eter drainage  Total Creation of			<b>3</b> ,	
Rate in accordance with §§ 4.88b or 4.89, whichever is appropriate.  Roby in consideration of the state of th			,	
4.89, whichever is appropriate. 7507 Nephroscious, arteriolar: Rate according to predominant symptoms as renal dysfunction, hypertension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation. 7508 Nephrolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year Hydronephrosis; except for recurrent stone formation in paired more or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year frequent attacks of colic, requiring catheter drainage for evaluation of 30 percent will be assigned for the service-connected testicular loss. Testis, undersecended, or congenitally undeveloped is not a ratable disability.  7510 Ureterolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or mor	7505 Kidney, tuberculosis of:			
7507 Nephrosolerosis, arteriolar: Rate according to predominant symptoms as renal dysfunction, hypertension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation.  7508 Nephrolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year of physione or more of the following: Frequent attacks of colic with infection (pyonephrosis), kidney function impaired ————————————————————————————————————			·	
Rate according to predominant symptoms as renal dysfunction, hypertension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation.  7508 Nephrolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procesurer attacks of colic, requiring catheter drainage.  7510 Ureterolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for re			,	30
toms as renal dysfunction, hypertensision or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation.  7508 Nephrolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year Severe; Rate as renal dysfunction. Frequent attacks of colic with infection (pyonephrosis), kidney function impaired Severe requiring catheter drainage			9 ,	
tension or heart disease. If rated under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation.  7508 Nephrolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year one phydronephrosis: Severe; Rate as renal dysfunction. Frequent attacks of colic with infection (pyonephrosis), kindey function impaired			9	20
under the cardiovascular schedule, however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation.  7508 Nephrolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year or notinected and not requiring cather drainage			· ,	
however, the percentage rating which would otherwise be assigned will be elevated to the next higher evaluation.  7508 Nephrolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year hydronephrosis; Severe; Rate as renal dysfunction. Frequent attacks of colic with infection (pyonephrosis), kidney function infected and not requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year resurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year resurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year resurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year resurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year resurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year resurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year resurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures mor				
which would otherwise be assigned will be elevated to the next higher evaluation.  7508 Nephrolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent attacks of colic with infection (pyonephrosis), kidney function impaired paired paired as a serial attack of colic, requiring catheter drainage one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy			·	
will be elevated to the next higher evaluation.  Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year Severe; Rate as renal dysfunction.  Frequent attacks of colic, requiring catheter drainage Only an occasional attack of colic, not infected and not requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 3. invasive or non-invasive procedures more than two times/year for recurrent stone formation requiring one or more of the following: 3. invasive or non-invasive procedures more than two times/year for the vertice testis unrelated to service, an evaluation of 30 percent will be assigned for the service-connected testis. with east or result inverticions; for the vertice formation requiring and invertice, infe				
7508 Nephrolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year only an occasional attack of colic, requiring catheter drainage Only an occasional attack of colic, requiring one or more of the following: 1. diet therapy 2. drug therapy 30	will be elevated to the next higher			
Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year paired undersomer than two times/year paired undersomer than two times/year one or more of the following: 1. diet therapy 2. drug therapy 30  Note: In cases of the removal of one testis as the result of a service-incurred injury or disease, other than an undescended or congenitally undeveloped testis, with the absence or nonfunctioning of the other testis unrelated to service-connected testicular loss. Testis, underscended, or congenitally undeveloped testis, with the absence or nonfunctioning of the other testis unrelated to service-connected testicular loss. Testis, underscended, or congenitally undeveloped is not a ratable disability.  7525  Epididymo-orchitis, chronic only: Rate as uninary tract infections. Rate in accordance with §§4.88b or 4.89, whichever is appropriate.  7510 Ureterolithiasis:  Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year residuals:  Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year residuals:  Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year residuals:  Rate as voiding dysfunction or urinary tract infections, includes interstitial and all etiologies, infectious and non-infectious:  Rate as voiding dysfunction:  Rate as				
recurrent stone formation requiring one or more of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year hydronephrosis: Severe; Rate as renal dysfunction. Frequent attacks of colic with infection (pyonephrosis), kidney function impaired maintenance or nonfunctioning of the other testis unrelated to service, an evaluation of 30 percent will be assigned for the service-connected testicular loss. Testis, underscended, or congenitally undeveloped is not a ratable disability.  Total Ureterolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year one or more of the following: 1. diet dreapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year one or more of the following: 1. diet dreapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year one or more of the following: 1. diet dreapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year one or more of the following: 1.			-	
one or more of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7509 Hydronephrosis: Severe; Rate as renal dysfunction. Frequent attacks of colic with infection (pyonephrosis), kidney function impaired				
1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7509 Hydronephrosis: Severe; Rate as renal dysfunction. Frequent attacks of colic with infection (pyonephrosis), kidney function impaired			One—o	
2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7509 Hydronephrosis: Severe; Rate as renal dysfunction, Frequent attacks of colic with infection (pyonephrosis), kidney function impaired				
3. invasive or non-invasive procedures more than two times/year Hydronephrosis:  Severe; Rate as renal dysfunction. Frequent attacks of colic with infection (pyonephrosis), kidney function impaired				
7509 Hydronephrosis: Severe; Rate as renal dysfunction. Frequent attacks of colic with infection (pyonephrosis), kidney function impaired				
Severe; Rate as renal dysfunction. Frequent attacks of colic with infection (pyonephrosis), kidney function impaired		30		
Frequent attacks of colic with infection (pyonephrosis), kidney function impaired				
(pyonephrosis), kidney function impaired (propose), kidney function of 30 percent will be assigned for the service-connected testicular loss. Testis, underscended, or congenitally undeveloped is not a ratable disability.  7525 Epididymo-orchitis, chronic only: Rate as unimary tract infections. For tubercular infections. For tubercular infections. Rate in accordance with §§ 4.88b or 4.89, whichever is appropriate.  7527 Prostate gland injuries, infections, hypertrophy, postoperative residuals: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.  7528 Malignant neoplasms of the genitourinary system (propose), whichever is predominant.  7529 Malignant neoplasms of the genitourinary system (propose), whichever is predominant.  7529 Malignant neoplasms of the genitourinary system (propose), whichever is predominant.				
Frequent attacks of colic, requiring catheter drainage				
Frequent attacks of colic, requiring catheter drainage		30	the service-connected testicular	
Only an occasional attack of colic, not infected and not requiring catheter drainage				
infected and not requiring catheter drainage		20		
drainage			able disability.	
7510 Ureterolithiasis: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7511 Ureter, stricture of: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.  7528 Malignant neoplasms of the genitourinary system		40	7525 Epididymo-orchitis, chronic only:	
Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year  7511 Ureter, stricture of: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year  7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction  7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.		10	Rate as urinary tract infection.	
recurrent stone formation requiring one or more of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year  7511 Ureter, stricture of: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.				
one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7511 Ureter, stricture of: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction or urinary tract infection, whichever is predominant. 7528 Malignant neoplasms of the genitorurinary system				
2. drug therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year  7511 Ureter, stricture of: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year  7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction  7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.  Rate as voiding dysfunction or urinary tract infection, whichever is predominant.			., .	
3. invasive or non-invasive procedures more than two times/year  7511 Ureter, stricture of: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year  7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction  7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.  7528 Malignant neoplasms of the genitorurinary system				
dures more than two times/year 7511 Ureter, stricture of: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction. 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.  30 tract infection, whichever is predominant. 7528 Malignant neoplasms of the genito-urinary system				
7511 Ureter, stricture of: Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following: 1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction. 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.		20		
Rate as hydronephrosis, except for recurrent stone formation requiring one or more of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction. 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.		30		
recurrent stone formation requiring one or more of the following:  1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction. 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.				
1. diet therapy 2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction. 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.				100
2. drug therapy 3. invasive or non-invasive procedures more than two times/year 7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction. 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.	one or more of the following:			
3. invasive or non-invasive procedures more than two times/year 7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious:  Rate as voiding dysfunction. 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.				
dures more than two times/year 7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious: Rate as voiding dysfunction. 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.				
7512 Cystitis, chronic, includes interstitial and all etiologies, infectious and non-infectious:  Rate as voiding dysfunction.  7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction  7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.		20		
and all etiologies, infectious and non-infectious:  Rate as voiding dysfunction.  7515 Bladder, calculus in, with symptoms interfering with function:  Rate as voiding dysfunction  7516 Bladder, fistula of:  Rate as voiding dysfunction or urinary tract infection, whichever is predominant.		30		
fectious: Rate as voiding dysfunction. 7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.				
7515 Bladder, calculus in, with symptoms interfering with function: Rate as voiding dysfunction 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.				
interfering with function: Rate as voiding dysfunction 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.				
Rate as voiding dysfunction 7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.				
7516 Bladder, fistula of: Rate as voiding dysfunction or urinary tract infection, whichever is predominant.				
Rate as voiding dysfunction or urinary tract infection, whichever is predominant.				
tract infection, whichever is pre- dominant.				
dominant.				
		100		

	Rat- ing		Rat- ing
Note—Following the cessation of surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure, the rating of 100 percent shall continue with a mandatory VA examination at the expiration of six months. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local reoccurrence or metastasis, rate on residuals as voiding dysfunction or renal dysfunction, whichever is predominant.  7529 Benign neoplasms of the genitourinary system: Rate as voiding dysfunction or renal dysfunction, whichever is predominant.  7530 Chronic renal disease requiring regular dialysis: Rate as renal dysfunction.  7531 Kidney transplant: Following transplant surgery	100	Or rate as renal dysfunction.  7533 Cystic diseases of the kidneys (polycystic disease, uremic medullary cystic disease, Medullary sponge kidney, and similar conditions):  Rate as renal dysfunction.  7534 Atherosclerotic renal disease (renal artery stenosis or atheroembolic renal disease):  Rate as renal dysfunction.  7535 Toxic nephropathy (antibotics, radiocontrast agents, nonsteroidal anti-inflammatory agents, heavy metals, and similar agents):  Rate as renal dysfunction.  7536 Glomerulonephritis:  Rate as renal dysfunction.  7537 Interstitial nephritis:  Rate as renal dysfunction.  7538 Papillary necrosis:  Rate as renal dysfunction.  7539 Renal amyloid disease:  Rate as renal dysfunction.  7540 Disseminated intravascular coagulation with renal cortical necrosis:  Rate as renal dysfunction.  7541 Renal involvement in diabetes mellitus, sickle cell anemia, systemic lupus erythematosus, vasculitis, or other systemic disease processes.  Rate as renal dysfunction.  7542 Neurogenic bladder:  Rate as voiding dysfunction.  1 Review for entitlement to special monthl pensation under § 3.350 of this chapter.	Iar. 29,
nephron function, etc.):  Minimum rating for symptomatic condition	20		

# § 4.116

GYNECOLOGICAL CONDITIONS AND DISORDERS OF THE BREAST

# §4.116 Schedule of ratings—gynecological conditions and disorders of the breast.

	Rating
Note 1: Natural menopause, primary amenor- rhea, and pregnancy and childbirth are not	
disabilities for rating purposes. Chronic re- siduals of medical or surgical complications of pregnancy may be disabilities for rating purposes.	
Note 2: When evaluating any claim involving	
loss or loss of use of one or more creative organs, refer to §3.350 of this chapter to determine whether the veteran may be entitled to special monthly compensation. Footnotes in the schedule indicate conditions which po-	
tentially establish entitlement to special monthly compensation; however, almost any condition in this section might, under certain circumstances, establish entitlement to spe-	
cial monthly compensation.  7610 Vulva, disease or injury of (including vulvovaginitis).	
<ul><li>7611 Vagina, disease or injury of.</li><li>7612 Cervix, disease or injury of.</li></ul>	
<ul> <li>7613 Uterus, disease, injury, or adhesions of.</li> <li>7614 Fallopian tube, disease, injury, or adhesions of (including pelvic inflammatory disease (PID)).</li> <li>7615 Ovary, disease, injury, or adhesions of.</li> </ul>	
General Rating Formula for Disease, Injury, or Adhesions of Female Reproductive Organs (diagnostic codes 7610 through 7615):	
Symptoms not controlled by continuous treat- ment	30 10
Symptoms that do not require continuous treat- ment	0
7617 Uterus and both ovaries, removal of, complete:	
For three months after removal  Thereafter	<sup>1</sup> 100 <sup>1</sup> 50
7618 Uterus, removal of, including corpus: For three months after removal Thereafter	<sup>1</sup> 100
7619 Ovary, removal of: For three months after removal	1100
Thereafter:  Complete removal of both ovaries  Removal of one with or without partial re-	130
moval of the other	<sup>1</sup> 0 <sup>1</sup> 20
7621 Uterus, prolapse: Complete, through vagina and introitus Incomplete	50 30
7622 Uterus, displacement of: With marked displacement and frequent or con-	30
tinuous menstrual disturbances With adhesions and irregular menstruation	30 10
7623 Pregnancy, surgical complications of: With rectocele or cystocele	50
With relaxation of perineum	10
quiring wearing of pad	100
of padVaginal fecal leakage one to three times per	60
week requiring wearing of pad Vaginal fecal leakage less than once a week	30 10

	Rating
Without leakage	0
'625 Fistula, urethrovaginal:  Multiple urethrovaginal fistulae	100
Requiring the use of an appliance or the wear-	100
ing of absorbent materials which must be	
changed more than four times per day	60
Requiring the wearing of absorbent materials which must be changed two to four times per	
day	40
Requiring the wearing of absorbent materials	
which must be changed less than two times per day	20
626 Breast, surgery of:	
Following radical mastectomy:	
Both One	80
Following modified radical mastectomy:	50
Both	60
One	40
Following simple mastectomy or wide local ex-	
cision with significant alteration of size or form:	
Both	50
One	30
Following wide local excision without significant alteration of size or form:	
Both or one	0
Note: For VA purposes:	
(1) Radical mastectomy means removal of	
the entire breast, underlying pectoral muscles, and regional lymph nodes up	
to the coracoclavicular ligament.	
(2) Modified radical mastectomy means re-	
moval of the entire breast and axillary lymph nodes (in continuity with the	
breast). Pectoral muscles are left intact.	
(3) Simple (or total) mastectomy means re-	
moval of all of the breast tissue, nipple,	
and a small portion of the overlying skin, but lymph nodes and muscles are left in-	
tact.	
(4) Wide local excision (including partial	
mastectomy, lumpectomy, tylectomy, segmentectomy, and quadrantectomy)	
means removal of a portion of the breast	
tissue.	
'627 Malignant neoplasms of gynecological sys-	100
Note: A rating of 100 percent shall continue	100
beyond the cessation of any surgical, X-ray,	
antineoplastic chemotherapy or other thera-	
peutic procedure. Six months after dis- continuance of such treatment, the appro-	
priate disability rating shall be determined by	
mandatory VA examination. Any change in	
evaluation based upon that or any subsequent examination shall be subject to the	
provisions of §3.105(e) of this chapter. If	
there has been no local recurrence or metas-	
tasis, rate on residuals.  628 Benign neoplasms of the gynecological sys-	
'628 Benign neoplasms of the gynecological system or breast. Rate according to impairment in	
function of the urinary or gynecological systems,	
or skin.	
'629 Endometriosis:  Lesions involving bowel or bladder confirmed	
by laparoscopy, pelvic pain or heavy or irreg-	
ular bleeding not controlled by treatment, and	
bowel or bladder symptoms	50
Pelvic pain or heavy or irregular bleeding not controlled by treatment	30
Pelvic pain or heavy or irregular bleeding not controlled by treatment  Pelvic pain or heavy or irregular bleeding re-	30

Rating

	Rating
<b>Note:</b> Diagnosis of endometriosis must be substantiated by laparoscopy.	
1 Deview for antillary and to any deligrantity	

<sup>1</sup>Review for entitlement to special monthly compensation under § 3.350 of this chapter.

[60 FR 19855, Apr. 21, 1995]

THE HEMIC AND LYMPHATIC SYSTEMS

# §4.117 Schedule of ratings—hemic and lymphatic systems.

Rating

7700 Anemia, hypochromicmicrocytic and megaloblastic, such as iron-deficiency and pernicious anemia: Hemoglobin 5gm/100ml or less, with findings such as high output congestive heart failure or dyspnea at rest ..... 100 Hemoglobin 7gm/100ml or less, with findings such as dyspnea on mild exertion, cardiomegaly, tachycardia (100 to 120 beats per minute) or syncope (three episodes in the last six months) ..... 70 Hemoglobin 8gm/100ml or less, with findings such as weakness, easy fatigability, headaches, lightheadedness, or shortness of breath ..... Hemoglobin 10gm/100ml or less with findings such as weakness, easy fatigability or headaches ..... Hemoglobin 10gm/100ml or less, asymptomatic .....

Note: Evaluate complications of pernicious anemia, such as dementia or peripheral neuropathy, separately.

7702 Agranulocytosis, acute:

Requiring bone marrow transplant, or; requiring transfusion of platelets or red cells at least once every six weeks, or; infections recurring at least once every six weeks .....
Requiring transfusion of platelets or red cells at least once every three months, or; infections recurring at least once every three months ...............
Requiring transfusion of platelets or red cells at least once every three months or platelets or red cells at least once per year but less than once every three months or infections recurring transfusion of platelets or red cells at least once per year but less than once every three months or infections.

NOTE: The 100 percent rating for bone marrow transplant shall be assigned as of the date of hospital admission and shall continue with a mandatory VA examination six months following hospital discharge. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter.

#### 7703 Leukemia:

NOTE: The 100 percent rating shall continue beyond the cessation of any surgical, radiation, antineoplastic chemotherapy or other therapeutic procedures. Six months after discontinuance of such treatment, the appropriate disability rating shall be determined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no recurrence, rate on residuals.

## 7704 Polycythemia vera:

NOTE: Rate complications such as hypertension, gout, stroke or thrombotic disease separately.

7705 Thrombocytopenia, primary, idiopathic or immune:

Platelet count of less than 20,000, with active bleeding, requiring treatment with medication and transfusions ....... 100 Platelet count between 20,000 and 70,000, not requiring treatment, without bleeding ..... 70 Stable platelet count between 70,000 and 100,000, without bleeding ..... 30 Stable platelet count of 100,000 or more, without bleeding ...... n 7706 Splenectomy ..... 20

Note: Rate complications such as systemic infections with encapsulated bacteria separately.

10

30

30 10

Rating	R	ating
7707 Spleen, injury of, healed. Rate for any residuals. 7709 Hodgkin's disease: With active disease or during a treatment phase	Note: The 100 percent rating shall continuous beyond the cessation of any surgical, ration, antineoplastic chemotherapy other therapeutic procedures. Six more after discontinuance of such treatmenthe appropriate disability rating shall dtermined by mandatory VA examition. Any change in evaluation be upon that or any subsequent examinal shall be subject to the provisions §3.105(e) of this chapter. If there has been local recurrence or metastasis, rateresiduals.  7716 Aplastic anemia:  Requiring bone marrow transplant, or; requiring transfusion of platelets or red cells at least once every six weeks, or; infections recurring at least once every six weeks	inue radi- or nths ent, l be ina- ased tion of peen
or inactive.  Rate under §§ 4.88c or 4.89 of this part, whichever is appropriate.  7714 Sickle cell anemia:	Requiring transfusion of plate- lets or red cells at least once every three months, or; infec-	100
With repeated painful crises, occurring in skin, joints, bones or any major organs caused by hemolysis and sickling of red blood cells, with anemia, thrombosis and infarction, with symptoms precluding even light manual labor	tions recurring at least once every three months	the con- tion rge. that
NOTE: Sickle cell trait alone, without a history of directly attributable pathological findings, is not a ratable disability. Cases of symptomatic sickle cell trait will be forwarded to the Director, Compensation and Pension Service, for consideration under §3.321(b)(1) of this chapter.	this chapter.  [60 FR 49227, Sept. 22, 1995]  THE SKIN  §4.118 Schedule of ratings—skin.	,
7715 Non-Hodgkin's lymphoma: With active disease or during a treatment phase		Rat- ing
•	7800 Scars, disfiguring, head, face or neck:  Complete or exceptionally repugnant deformity of one side of face or marked or repugnant bilateral disfigurement	50 30 10

Slight .....

	Rat- ing	
Note: When in addition to tissue loss and	_	Rate so
cicatrization there is marked discoloration,		cons
color contrast, or the like, the 50 percent rat-		7819 New
ing under Code 7800 may be increased to 80		Rate as
percent, the 30 percent to 50 percent, and the		Unless
10 percent to 30 percent. The most repugnant,		throu
disfiguring conditions, including scars and dis-		locati
eases of the skin, may be submitted for cen-		disab
tral office rating, with several unretouched photographs.		Note: subm
7801 Scars, burns, third degree:		unret
Area or areas exceeding 1 square foot (0.1 m.2)	40	ings
Area or areas exceeding one-half square foot		Cent
(0.05 m.²)	30	pemp
Area or areas exceeding 12 square inches (77.4		stituti
cm.2)	20	
Area or areas exceeding 6 square inches (38.7		[29 FR 6
cm. <sup>2</sup> )	10	FR 5063,
NOTE (1): Actual third degree residual involve-		,
ment required to the extent shown under		1975; 43 F
7801.		
NOTE (2): Ratings for widely separated areas, as on two or more extremities or on anterior and		
posterior surfaces of extremities or trunk, will		64 110
be separately rated and combined.		§4.119
7802 Scars, burns, second degree:		syst
Area or areas approximating 1 square foot (0.1		-
m.²)	10	
NOTE: See NOTE (2) under diagnostic code		
7801.		7900 Hype
7803 Scars, superficial, poorly nourished, with re-		Thyroid 6
peated ulceration	10	beats
7804 Scars, superficial, tender and painful on ob-	10	weakne
jective demonstration	10	ous sy
when the requirements are met, even though		sympto Emotiona
the location may be on tip of finger or toe, and		creased
the rating may exceed the amputation value		Tachycar
for the limited involvement.		or bloo
7805 Scars, other.		Tachycar
Rate on limitation of function of part affected.		or; con
7806 Eczema:		NOTE (1):
With ulceration or extensive exfoliation or crust-		finding, (DC 70
ing, and systemic or nervous manifestations,		evaluat
or exceptionally repugnant	50	NOTE (2)
With exudation or itching constant, extensive lesions, or marked disfigurement	30	evaluat
With exfoliation, exudation or itching, if involving	30	diplopia
an exposed surface or extensive area	10	ual acu
With slight, if any, exfoliation, exudation or		7901 Thyr
itching, if on a nonexposed surface or small		Thyroid 6
area	0	beats weakne
7807 Leishmaniasis, americana (mucocutaneous,		OUS S
espundia).		sympto
7808 Leishmaniasis, old world (cutaneous, oriental		Emotiona
sore).		creased
7809 Lupus erythematosus, discoid. (Not to be combined with ratings under diag-		Tachycar
nostic code 6350.)		or bloo
7810 Pinta.		Tachycar
7811 Tuberculosis luposa (lupus vulgaris), active or		or; con
inactive:		Note (1): finding,
Active	100	(DC 70
Inactive: See §§ 4.88b and 4.89.		evaluat
7812 Verruga peruana.		NOTE (2)
7813 Dermatophytosis.		evaluat
7814 Tinea barbae.	1	diplopia
7815 Pemphigus.		ual acu
<ul><li>7815 Pemphigus.</li><li>7816 Psoriasis.</li></ul>		7902 Thyr
7815 Pemphigus.		

	Rat- ing
Rate scars, disfigurement, etc., on the extent of constitutional symptoms, physical impairment.  7819 New growths, benign, skin. Rate as scars, disfigurement, etc. Unless otherwise provided, rate codes 7807 through 7819 as for eczema, dependent upon location, extent, and repugnant or otherwise disabling character of manifestations.  NOTE: The most repugnant conditions may be submitted for central office rating with several	
unretouched photographs. Total disability ratings may be assigned without reference to	
Central Office in the most severe cases of pemphigus and dermatitis exfoliativa with con- stitutional symptoms.	

[29 FR 6718, May 22, 1964, as amended at 34 FR 5063, Mar. 11, 1969; 40 FR 42540, Sept. 15, 1975; 43 FR 45262, Oct. 2, 1978]

## THE ENDOCRINE SYSTEM

# § 4.119 Schedule of ratings—endocrine system.

	Rat ing
7900 Hyperthyroidism	
Thyroid enlargement, tachycardia (more than 100	
beats per minute), eye involvement, muscular	
weakness, loss of weight, and sympathetic nerv- ous system, cardiovascular, or astrointestinal	
symptoms	10
Emotional instability, tachycardia, fatigability, and in-	10
creased pulse pressure or blood pressure	6
Tachycardia, tremor, and increased pulse pressure	"
or blood pressure	3
Tachycardia, which may be intermittent, and tremor,	
or; continuous medication required for control	1
NOTE (1): If disease of the heart is the predominant	
finding, evaluate as hyperthyroid heart disease	
(DC 7008) if doing so would result in a higher	
evaluation than using the criteria above.  NOTE (2): If ophthalmopathy is the sole finding,	
evaluate as field vision, impairment of (DC 6080);	
diplopia (DC 6090); or impairment of (DC 6000),	
ual acuity (DC 6061–6079).	
7901 Thyroid gland, toxic adenoma of	
Thyroid enlargement, tachycardia (more than 100	
beats per minute), eye involvement, muscular	
weakness, loss of weight, and sympathetic nerv-	
ous system, cardiovascular, or gastrointestinal	
symptoms Emotional instability, tachycardia, fatigability, and in-	10
creased pulse pressure or blood pressure	6
Tachycardia, tremor, and increased pulse pressure	
or blood pressure	3
Tachycardia, which may be intermittent, and tremor,	`
or; continuous medication required for control	1
NOTE (1): If disease of the heart is the predominant	
finding, evaluate as hyperthyroid heart disease	
(DC 7008) if doing so would result in a higher	
evaluation than using the criteria above.	
NOTE (2): If ophthalmopathy is the sole finding,	
evaluate as field vision, impairment of (DC 6080); diplopia (DC 6090); or impairment of central vis-	
ual acuity (DC 6061–6079).	
7902 Thyroid gland, nontoxic adenoma of	
	1
With disfigurement of the head or neck	2

NOTE: If there are symptoms due to pressure on adjacent organs such as the trachea, larynx, or esophagus, evaluate under the diagnostic code  Rating  One or two crises during the past year, or; two to four episodes during the past year, or; weakness and fatigability, or; corticosteroid therapy required	ing 20
jacent organs such as the trachea, larynx, or esophagus, evaluate under the diagnostic code four episodes during the past year, or; weakness and fatigability, or; corticosteroid therapy required	20
for disability of that organ, if doing so would result in a higher evaluation than using this diagnostic code.  NOTE (1): An Addisonian "crisis" consists of the rapid onset of peripheral vascular collapse (with	
7903 Hypothyroidism acute hypotension and shock), with findings that	
Cold intolerance, muscular weakness, cardio- may include: anorexia; nausea; vomiting; dehy-	
vascular involvement, mental disturbance (demen-	
tia, slowing of thought, depression), bradycardia legs, and back; fever; apathy, and depressed (less than 60 beats per minute), and sleepiness 100 mentation with possible progression to coma,	
Muscular weakness, mental disturbance, and weight renal shutdown, and death.	
gain	
Fatigability, constipation, and mental sluggishness   30 poses, is a less acute and less severe event than Fatigability, or; continuous medication required for an Addisonian crisis and may consist of anorexia,	
control	
7904 Hyperparathyroidism ness, malaise, orthostatic hypotension, or hypo-	
Generalized decalcification of bones, kidney stones, glycemia, but no peripheral vascular collapse.	
gastrointestinal symptoms (nausea, vomiting, ano- rexia, constipation, weight loss, or peptic ulcer), NOTE (3): Tuberculous Addison's disease will be evaluated as active or inactive tuberculosis. If in-	
and weakness	
Gastrointestinal symptoms and weakness	
Continuous medication required for control	
digestive, skeletal, renal, or cardiovascular residu-	
als or as endocrine dysfunction. Evaluate according to major manifestations.	
7905 Hypoparathyroidism  Marked neuromuscular excitability (such as convul-  Marked neuromuscular excitability (such as convul-	
sions, muscular spasms (tetany), or laryngeal restricted diet, and regulation of activities (avoid-	
stridor) plus either cataract or evidence of in-	
creased intracranial pressure (such as activities) with episodes of ketoacidosis or hypo-	
papilledema)	
(of arms, legs, or circumoral area) plus either cat-	
aract or evidence of increased intracranial pres-	400
sure	100
7907 Cushing's syndrome activities with episodes of ketoacidosis or hypo-	
As active, progressive disease including loss of glycemic reactions requiring one or two hos-	
muscle strength, areas of osteoporosis, hyper- tension, weakness, and enlargement of pituitary pitalizations per year or twice a month visits to a diabetic care provider, plus complications that	
or adrenal gland	60
Loss of muscle strength and enlargement of pitui-	
tary or adrenal gland	40
With strae, obesity, moon face, glucose intolerance, and vascular fragility	20
Note: With recovery or control, evaluate as residu- Manageable by restricted diet only	10
als of adrenal insufficiency or cardiovascular, psylochiatric, skin, or skeletal complications under ap-	
chiatric, skin, or skeletal complications under appropriate diagnostic code.  diabetes separately unless they are part of the criteria used to support a 100 percent evaluation.	
7908 Acromegaly Noncompensable complications are considered	
Evidence of increased intracranial pressure (such as visual field defect), arthropathy, glucose intoler- 7913.	
visual field defect), arthropathy, glucose intoler- ance, and either hypertension or cardiomegaly 100 Note (2): When diabetes mellitus has been conclu-	
Arthropathy, glucose intolerance, and hypertension 60 sively diagnosed, do not request a glucose toler-	
Enlargement of acral parts or overgrowth of long bones, and enlarged sella turcica	
bones, and enlarged sella turcica	100
Polyuria with near-continuous thirst, and more than NOTE: A rating of 100 percent shall continue beyond	
two documented episodes of dehydration requir-	
ing parenteral hydration in the past year	
documented episodes of dehydration requiring the appropriate disability rating shall be deter-	
parenteral hydration in the past year	
Polyuria with near-continuous thirst, and one or more episodes of dehydration in the past year not examination shall be subject to the provisions of	
requiring parenteral hydration	
Polyuria with near-continuous thirst	
7911 Addison's disease (Adrenal Cortical 7915 Neoplasm, benign, any specified part of the en-	
Hypofunction) docrine system rate as residuals of endocrine dys- Four or more crises during the past year 60 function.	
Three crises during the past year, or, five or more 7916 Hyperpituitarism (prolactin secreting pituitary	
episodes during the past year	

	Rat- ing
7917 Hyperaldosteronism (benign or malignant) 7918 Pheochromocytoma (benign or malignant) NOTE: Evaluate diagnostic codes 7916, 7917, and 7918 as malignant or benign neoplasm as appro- priate. 7919 C-cell hyperplasia of the thyroid NOTE: A rating of 100 percent shall continue beyond the cessation of any surgical, X-ray, antineoplastic chemotherapy or other therapeutic procedure. Six months after discontinuance of such treatment, the appropriate disability rating shall be deter- mined by mandatory VA examination. Any change in evaluation based upon that or any subsequent examination shall be subject to the provisions of §3.105(e) of this chapter. If there has been no local recurrence or metastasis, rate on residuals.	100

[61 FR 20446, May 7, 1996]

NEUROLOGICAL CONDITIONS AND CONVULSIVE DISORDERS

#### §4.120 Evaluations by comparison.

Disability in this field is ordinarily to be rated in proportion to the impairment of motor, sensory or mental function. Consider especially psychotic manifestations, complete or partial loss of use of one or more extremities, speech disturbances, impairment of vision, disturbances of gait, tremors, visceral manifestations, injury to the skull, etc. In rating disability from the conditions in the preceding sentence refer to the appropriate schedule. In rating peripheral nerve injuries and their residuals, attention should be given to the site and character of the injury, the relative impairment in motor function, trophic changes, or sensory disturbances.

#### §4.121 Identification of epilepsy.

When there is doubt as to the true nature of epileptiform attacks, neurological observation in a hospital adequate to make such a study is necessary. To warrant a rating for epilepsy, the seizures must be witnessed or verified at some time by a physician. As to frequency, competent, consistent lay testimony emphasizing convulsive and immediate post-convulsive characteristics may be accepted. The frequency of seizures should be ascertained under the ordinary conditions of life (while not hospitalized).

#### §4.122 Psychomotor epilepsy.

The term psychomotor epilepsy refers to a condition that is characterized by seizures and not uncommonly by a chronic psychiatric disturbance as well.

(a) Psychomotor seizures consist of episodic alterations in conscious control that may be associated with automatic states, generalized convulsions, random motor movements (chewing, lip smacking, fumbling), hallucinatory phenomena (involving taste, smell, sound, vision), perceptual illusions (deja vu, feelings of loneliness, strangeness, macropsia, micropsia, dreamy states), alterations in thinking (not open to reason), alterations in memory, abnormalities of mood or affect (fear, alarm, terror, anger, dread, wellbeing), and autonomic disturbances (sweating, pallor, flushing of the face, visceral phenomena such as nausea, vomiting, defecation, a rising feeling of warmth in the abdomen). Automatic states or automatisms are characterized by episodes of irrational, irrelevant, disjointed, unconventional, asocial, purposeless though seemingly coordinated and purposeful, confused or inappropriate activity of one to several minutes (or, infrequently, hours) duration with subsequent amnesia for the seizure. Examples: A person of high social standing remained seated, muttered angrily, and rubbed the arms of his chair while the National Anthem was being played; an apparently normal person suddenly disrobed in public; a man traded an expensive automobile for an antiquated automobile in poor mechanical condition and after regaining conscious control, discovered that he had signed an agreement to pay an additional sum of money in the trade. The seizure manifestations of psychomotor epilepsy vary from patient to patient and in the same patient from seizure to seizure.

(b) A chronic mental disorder is not uncommon as an interseizure manifestation of psychomotor epilepsy and may include psychiatric disturbances extending from minimal anxiety to severe personality disorder (as distinguished from developmental) or almost complete personality disintegration (psychosis). The manifestations of a chronic mental disorder associated

#### § 4.123

with psychomotor epilepsy, like those of the seizures, are protean in character.

#### §4.123 Neuritis, cranial or peripheral.

Neuritis, cranial or peripheral, characterized by loss of reflexes, muscle atrophy, sensory disturbances, and constant pain, at times excruciating, is to be rated on the scale provided for injury of the nerve involved, with a maximum equal to severe, incomplete, paralysis. See nerve involved for diagnostic code number and rating. The maximum rating which may be assigned for neuritis not characterized by organic changes referred to in this section will be that for moderate, or with sciatic nerve involvement, for moderately severe, incomplete paralysis.

### § 4.124 Neuralgia, cranial or peripheral.

Neuralgia, cranial or peripheral, characterized usually by a dull and intermittent pain, of typical distribution so as to identify the nerve, is to be rated on the same scale, with a maximum equal to moderate incomplete paralysis. See nerve involved for diagnostic code number and rating. Tic douloureux, or trifacial neuralgia, may be rated up to complete paralysis of the affected nerve.

#### §4.124a Schedule of ratings—neurological conditions and convulsive disorders.

[With the exceptions noted, disability from the following diseases and their residuals may be rated from 10 percent to 100 percent in proportion to the impairment of motor, sensory, or mental function. Consider especially psychotic manifestations, complete or partial loss of use of one or more extremities, speech disturbances, impairment of vision, disturbances of gait, tremors, visceral manifestations, etc., referring to the appropriate bodily system of the schedule. With partial loss of use of one or more extremities from neurological lesions, rate by comparison with the mild, moderate, severe, or complete paralysis of peripheral nerves]

ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM

	Rat- ing
8000 Encephalitis, epidemic, chronic: As active febrile disease	100

### ORGANIC DISEASES OF THE CENTRAL NERVOUS SYSTEM—Continued

SYSTEM—Continued	
	Rat- ing
Rate residuals, minimum	10
Brain, new growth of:	100
8002 Malignant	100
for 2 years following cessation of surgical,	
chemotherapeutic or other treatment modality.	
At this point, if the residuals have stabilized,	
the rating will be made on neurological residuals according to symptomatology.	
Minimum rating	30
8003 Benign, minimum	60
Rate residuals, minimum	10
8004 Paralysis agitans:	
Minimum rating	30
8005 Bulbar palsy	100
8007 Brain, vessels, embolism of.	
8008 Brain, vessels, thrombosis of.	
8009 Brain, vessels, hemorrhage from: Rate the vascular conditions under Codes 8007	
through 8009, for 6 months	100
Rate residuals, thereafter, minimum	10
8010 Myelitis:	
Minimum rating	10
8011 Poliomyelitis, anterior:	
As active febrile disease	100
Rate residuals, minimum	10
8012 Hematomyelia:	400
For 6 months	100 10
Rate residuals, minimum	10
8014 Syphilis, meningovascular.	
8015 Tabes dorsalis.	
NOTE: Rate upon the severity of convulsions, pa-	
ralysis, visual impairment or psychotic involvement, etc.	
8017 Amyotrophic lateral sclerosis:	
Minimum rating	30
8018 Multiple sclerosis:	30
Minimum rating	30
As active febrile disease	100
Rate residuals, minimum	10
8020 Brain, abscess of:	
As active disease	100
Rate residuals, minimum	10
Spinal cord, new growths of:.	
8021 Malignant	100
NOTE: The rating in code 8021 will be continued for 2 years following cessation of surgical,	
chemotherapeutic or other treatment modality.	
At this point, if the residuals have stabilized,	
the rating will be made on neurological residu-	
als according to symptomatology.	
Minimum rating	30
8022 Benign, minimum rating	60
Rate residuals, minimum	10
8023 Progressive muscular atrophy: Minimum rating	30
	30
8024 Syringomyelia: Minimum rating	30
8025 Myasthenia gravis:	00
Minimum rating	30
Ÿ	

SYSTEM—Continued	
	Rat- ing
Note: It is required for the minimum ratings for residuals under diagnostic codes 8000–8025, that there be ascertainable residuals. Determinations as to the presence of residuals not capable of objective verification, i.e., headaches, dizziness, fatigability, must be approached on the basis of the diagnosis recorded; subjective residuals will be accepted when consistent with the disease and not more likely attributable to other disease or no disease. It is of exceptional importance that when ratings in excess of the prescribed minimum ratings are assigned, the diagnostic codes utilized as bases of evaluation be cited, in addition to the codes identifying the diagnoses.  8045 Brain disease due to trauma:  Purely neurological disabilities, such as hemiplegia, epileptiform seizures, facial nerve paralysis, etc., following trauma to the brain, will be rated under the diagnostic codes specifi-	
cally dealing with such disabilities, with citation of a hyphenated diagnostic code (e.g., 8045–8207).	
Purely subjective complaints such as headache, dizziness, insomnia, etc., recognized as symptomatic of brain trauma, will be rated 10 percent and no more under diagnostic code 9304. This 10 percent rating will not be combined with any other rating for a disability due to brain trauma. Ratings in excess of 10 percent for brain disease due to trauma under diagnostic code 9304 are not assignable in the absence of a diagnosis of multi-infarct dementia associated with brain trauma.	
8046 Cerebral arteriosclerosis:  Purely neurological disabilities, such as hemiplegia, cranial nerve paralysis, etc., due to cerebral arteriosclerosis will be rated under the diagnostic codes dealing with such specific disabilities, with citation of a hyphenated diagnostic code, or, 8046, 8207.	

disabilities, with citation of a hyphenated diagnostic code (e.g., 8046–8207).

Purely subjective complaints such as headache, dizziness, tinnitus, insomnia and irritability, recognized as symptomatic of a properly diagnosed cerebral arteriosclerosis, will be rated 10 percent and no more under diagnostic code 9305. This 10 percent rating will not be combined with any other rating for a disability due to cerebral or generalized arteriosclerosis. Ratings in excess of 10 percent for cerebral arteriosclerosis under diagnostic code 9305 are not assignable in the absence of a diagnosis of multi-infarct dementia with cerebral arteriosclerosis.

NOTE: The ratings under code 8046 apply only

NOTE: The ratings under code 8046 apply only when the diagnosis of cerebral arteriosclerosis is substantiated by the entire clinical picture and not solely on findings of retinal arterio-

#### MISCELLANEOUS DISEASES

		Rat- ing
8100	Migraine:	

#### MISCELLANEOUS DISEASES—Continued

	Rat- ing
With very frequent completely prostrating and prolonged attacks productive of severe economic inadaptability	50
months	30
one in 2 months over last several months With less frequent attacks	10 0
8103 Tic, convulsive: Severe Moderate Mild NOTE: Depending upon frequency, severity, mus-	30 10 0
cle groups involved. 8104 Paramyoclonus multiplex (convulsive state, myoclonic type):	
Rate as tic; convulsive; severe cases	60
Pronounced, progressive grave types	100 80 50 30 10
Rate as for epilepsy, petit mal.	

DISEASES OF THE CRANIAL NERVES	
	Rat- ing
Disability from lesions of peripheral portions of first, second, third, fourth, sixth, and eighth nerves will be rated under the Organs of Special Sense. The ratings for the cranial nerves are for unilateral involvement; when bilateral, combine but without the bilateral factor.  Fifth (trigeminal) cranial nerve  8205 Paralysis of:	
Complete	50 30 10
Note: Tic douloureux may be rated in accordance with severity, up to complete paralysis.  Seventh (facial) cranial nerve  8207 Paralysis of:	00
Complete Incomplete, severe Incomplete, moderate NOTE: Dependent upon relative loss of innervation of facial muscles. 8307 Neuritis. 8407 Neuralgia. Ninth (glossopharyngeal) cranial nerve. 8209 Paralysis of:	30 20 10
Complete	30 20 10

### 38 CFR Ch. I (7-1-99 Edition)

#### § 4.124a

### DISEASES OF THE CRANIAL NERVES—Continued

	Rat- ing
NOTE: Dependent upon relative loss of ordinary sensation in mucous membrane of the pharynx, fauces, and tonsils.  8309 Neuritis.	
8409 Neuralgia.	
Tenth (pneumogastric, vagus) cranial nerve.	
8210 Paralysis of:	
Complete	5
Incomplete, severe	3
Incomplete, moderate	1
NOTE: Dependent upon extent of sensory and motor loss to organs of voice, respiration, pharynx, stomach and heart.	
8310 Neuritis.	
8410 Neuralgia.	
Eleventh (spinal accessory, external branch) cra- nial nerve.	
8211 Paralysis of:	
Complete	3
Incomplete, severe	2
Incomplete, moderate	1
Note: Dependent upon loss of motor function of	
sternomastoid and trapezius muscles.	
8311 Neuritis.	
8411 Neuralgia.	
Twelfth (hypoglossal) cranial nerve.	
8212 Paralysis of:	
Complete	5
Incomplete, severe	3
Incomplete, moderate	1
NOTE: Dependent upon loss of motor function of	
tongue.	
8312 Neuritis.	
8412 Neuralgia.	

#### DISEASES OF THE PERIPHERAL NERVES

Cabadula of votings	Rat	ing
Schedule of ratings	Major	Minor
The term "incomplete paralysis," with this and other peripheral nerve injuries, indicates a degree of lost or impaired function substantially less than the type picture for complete paralysis given with each nerve, whether due to varied level of the nerve lesion or to partial regeneration. When the involvement is wholly sensory, the rating should be for the mild, or at most, the moderate degree. The ratings for the peripheral nerves are for unilateral involvement; when bilateral, combine with application of the bilateral factor.  Upper radicular group (fifth and sixth cervicals)		
8510 Paralysis of:  Complete; all shoulder and elbow movements lost or severely affected, hand and wrist movements not affected Incomplete:	70	60
Severe	50	40
Moderate	40	30
Mild	20	20

#### DISEASES OF THE PERIPHERAL NERVES— Continued

Schedule of ratings	Rati	ing
Concadio of fallings	Major	Minor
8610 Neuritis.		
8710 Neuralgia.		
Middle radicular group		
8511 Paralysis of:		
Complete; adduction, abduction and ro-		
tation of arm, flexion of elbow, and ex- tension of wrist lost or severely af-		
fected	70	6
Incomplete:		
Severe	50	4
Moderate	40	3
Mild8611 Neuritis.	20	2
8711 Neuralgia.		
Lower radicular group		
8512 Paralysis of:		
Complete; all intrinsic muscles of hand,		
and some or all of flexors of wrist and		
fingers, paralyzed (substantial loss of	70	6
use of hand) Incomplete:	70	0
Severe	50	4
Moderate	40	3
Mild	20	2
8612 Neuritis.		
8712 Neuralgia.		
All radicular groups		
8513 Paralysis of:  Complete	90	8
Incomplete:	00	Ū
Severe	70	6
Moderate	40	3
Mild	20	2
8613 Neuritis. 8713 Neuralgia.		
ŭ		
The musculospiral nerve (radial nerve) 8514 Paralysis of:		
Complete; drop of hand and fingers,		
wrist and fingers perpetually flexed,		
the thumb adducted falling within the		
line of the outer border of the index finger; can not extend hand at wrist,		
extend proximal phalanges of fingers,		
extend thumb, or make lateral move-		
ment of wrist; supination of hand, extension and flexion of elbow weak-		
ened, the loss of synergic motion of		
extensors impairs the hand grip seri-		
ously; total paralysis of the triceps oc-	70	^
curs only as the greatest rarity	70	6
Severe	50	4
Moderate	30	2
	20	2

#### DISEASES OF THE PERIPHERAL NERVES— Continued

0.1.1.7.5	Rat	ing
Schedule of ratings	Major	Minor
8614 Neuritis. 8714 Neuralgia.		
NOTE: Lesions involving only "dissociat communis digitorum" and "paralysis bel communis digitorum," will not exceed the ing under code 8514.	ow the e	xtensor
The median nerve		
8515 Paralysis of: Complete; the hand inclined to the ulnar side, the index and middle fingers more extended than normally, considerable atrophy of the muscles of the thenar eminence, the thumb in the plane of the hand (ape hand); pronation incomplete and defective, absence of flexion of index finger and feeble flexion of middle finger, cannot make a fist, index and middle fingers remain extended; cannot flex distal phalanx of thumb, defective opposition and abduction of the thumb, at right angles to palm; flexion of wrist weakened; pain with trophic disturbances Incomplete: Severe	70 50 30 10	60 40 20 10
8715 Neuralgia.		
The ulnar nerve  8516 Paralysis of: Complete; the "griffin claw" deformity, due to flexor contraction of ring and lit- tle fingers, atrophy very marked in dor- sal interspace and thenar and hypothenar eminences; loss of exten- sion of ring and little fingers cannot spread the fingers (or reverse), cannot adduct the thumb; flexion of wrist		
weakenedIncomplete:	60	50
Severe	40 30 10	30 20 10
Musculocutaneous nerve		
8517 Paralysis of: Complete; weakness but not loss of flex- ion of elbow and supination of forearm Incomplete:	30	20
Severe	20	20
Moderate	10 0	10 0
Circumflex nerve 8518 Paralysis of:		
Complete; abduction of arm is impos- sible, outward rotation is weakened; muscles supplied are deltoid and teres minor	50	40
Incomplete: Severe	30	20
Moderate	10	10

## DISEASES OF THE PERIPHERAL NERVES—Continued

0 30 20	Minor (
30	20
20	
20	
	20
10 0	10
·	houlder
	by ref- t in ex-

60		Rating	
40	Sciatic nerve		
20 10	8520 Paralysis of: Complete; the foot dangles and drops, no active movement possible of mus- cles below the knee, flexion of knee		
	weakened or (very rarely) lost	80	
	Severe, with marked muscular atrophy Moderately severe Moderate Mild 8620 Neuritis. 8720 Neuralgia.	60 40 20 10	
50	External popliteal nerve (common peroneal)		
30 20 10	8521 Paralysis of:  Complete; foot drop and slight droop of first phalanges of all toes, cannot dorsiflex the foot, extension (dorsal flexion) of proximal phalanges of toes lost; abduction of foot lost, adduction weakened; anesthesia covers entire dorsum of foot and toes	40	
20	Incomplete:		
20 10 0	Severe Moderate Mild 8621 Neuritis. 8721 Neuralgia.	30 20 10	
Musculocutaneous nerve (superficial peroneal)			
	8522 Paralysis of: Complete; eversion of foot weakened	30	
40	Incomplete: Severe	20	
20 10	ModerateMild	10 0	

#### § 4.124a

	Rating		Rating
8622 Neuritis. 8722 Neuralgia.		8628 Neuritis. 8728 Neuralgia.	
Anterior tibial nerve (deep peroneal)		External cutaneous nerve of thigh	
8523 Paralysis of:		8529 Paralysis of:	
Complete; dorsal flexion of foot lost	30	Severe to complete	10
Incomplete:		Mild or moderate	0
Severe	20	8629 Neuritis.	
Moderate	10	8729 Neuralgia.	
Mild	0	Ilio-inguinal nerve	
8723 Neuralgia.		8530 Paralysis of:	
9		Severe to complete	10
Internal popliteal nerve (tibial)		Mild or moderate	0
8524 Paralysis of:		8730 Neuralgia.	
Complete; plantar flexion lost, frank adduction of foot impossible, flexion		8540 Soft-tissue sarcoma (of neurogenic	
and separation of toes abolished; no		origin)	100
muscle in sole can move; in lesions of		NOTE: The 100 percent rating will be con	tinued
the nerve high in popliteal fossa, plan-		for 6 months following the cessation of su	
tar flexion of foot is lost	40	X-ray, antineoplastic chemotherapy or	
Incomplete:	20	therapeutic procedure. At this point, if the been no local recurrence or metastase	
Severe	30 20	rating will be made on residuals.	.o, tilo
Mild	10		
8624 Neuritis.		THE EPILEPSIES	
8724 Neuralgia.			
Posterior tibial nerve			Rat- ing
8525 Paralysis of:			+
Complete; paralysis of all muscles of		A thorough study of all material in §§ 4.121 and 4.122 of the preface and under the ratings for	
sole of foot, frequently with painful pa-		epilepsy is necessary prior to any rating ac-	
ralysis of a causalgic nature; toes can- not be flexed; adduction is weakened;		tion.	
plantar flexion is impaired	30	8910 Epilepsy, grand mal.  Rate under the general rating formula for major	.
Incomplete:		seizures.	
Severe	20	8911 Epilepsy, petit mal.	
Moderate	10	Rate under the general rating formula for minor seizures.	
Mild	10	NOTE (1): A major seizure is characterized by	
8625 Neuritis. 8725 Neuralgia.		the generalized tonic-clonic convulsion with unconsciousness.	
ŭ		NOTE (2): A minor seizure consists of a brief	:
Anterior crural nerve (femoral)		interruption in consciousness or conscious	
8526 Paralysis of:		control associated with staring or rhythmic blinking of the eyes or nodding of the head	
Complete; paralysis of quadriceps extensor muscles	40	("pure" petit mal), or sudden jerking move-	
Incomplete:	40	ments of the arms, trunk, or head (myoclonic	
Severe	30	type) or sudden loss of postural control (akinetic type).	
Moderate	20	General Rating Formula for Major and Minor Ep-	.
Mild	10	ileptic Seizures:  Averaging at least 1 major seizure per	
8626 Neuritis.		month over the last year	
8726 Neuralgia.		Averaging at least 1 major seizure in 3	
Internal saphenous nerve		months over the last year; or more than 10 minor seizures weekly	
8527 Paralysis of:		Averaging at least 1 major seizure in 4	
Severe to complete	10	months over the last year; or 9-10 minor	
Mild to moderate	0	seizures per week	
8627 Neuritis. 8727 Neuralgia.		or 2 in the last year; or averaging at least	
9		5 to 8 minor seizures weekly	
Obturator nerve		At least 1 major seizure in the last 2 years; or at least 2 minor seizures in the last 6	
8528 Paralysis of:		months	20
Severe to complete	10	A confirmed diagnosis of epilepsy with a	
Mild or moderate	0	history of seizures	1 10

#### THE EPILEPSIES—Continued

Rat-

NOTE (1): When continuous medication is shown necessary for the control of epilepsy, the minimum evaluation will be 10 percent. This rating will not be combined with any other rating for

epilepsy. NOTE (2): In the presence of major and minor seizures, rate the predominating type.

NOTE (3): There will be no distinction between diurnal and nocturnal major seizures

8912 Epilepsy, Jacksonian and focal motor or sen-

sory. 8913 Epilepsy, diencephalic.

Rate as minor seizures, except in the presence of major and minor seizures, rate the predominating type.

8914 Epilepsy, psychomotor.

Major seizures:

Psychomotor seizures will be rated as major seizures under the general rating formula when characterized by automatic states and/or generalized convulsions with unconsciousness.

Minor seizures:

Psychomotor seizures will be rated as minor seizures under the general rating formula when characterized by brief transient episodes of random motor movements, hallucinations, perceptual illusions, abnormalities of thinking, memory or mood, or autonomic disturbances.

Mental Disorders in Epilepsies: A nonpsychotic organic brain syndrome will be rated separately under the appropriate diagnostic code (e.g., 9304 or 9307). In the absence of a diagnosis of non-psychotic organic psychiatric disturbance (psychotic, psychoneurotic or personality disorder) if diagnosed and shown to be secondary to or directly associated with epilepsy will be rated separately. The psychotic or psychroneurotic disorder will be rated under the appropriate diagnostic code. The personality disorder will be rated as a dementia (e.g., diagnostic code 9304 or 9307). Epilepsy and Unemplovability. (1) Rating specialists must

Epilepsy and Unemployability: (1) Rating specialists must bear in mind that the epileptic, although his or her seizures are controlled, may find employment and rehabilitation difficult of attainment due to employer reluctance to the hiring of the orbitotist.

of attainment due to employer reluctance to the hiring of the epileptic.

(2) Where a case is encountered with a definite history of unemployment, full and complete development should be undertaken to ascertain whether the epilepsy is the determining factor in his or her inability to obtain employment.

(3) The assent of the claimant should first be obtained for permission to conduct this economic and social survey. The purpose of this survey is to secure all the relevant facts and data necessary to permit of a true judgment as to the reason for his or her unemployment and should include information as to:

for his or her unemployment and should include information as to:

(a) Education;
(b) Occupations prior and subsequent to service;
(c) Places of employment and reasons for termination;
(d) Wages received;
(e) Number of seizures.

(4) Upon completion of this survey and current examination, the case should have rating board toonsideration. Where in the judgment of the rating board the veteran's unemployability is due to epilense and jurisdiction is not vested in that body by due to epilepsy and jurisdiction is not vested in that body by reason of schedular evaluations, the case should be submitted to the Director, Compensation and Pension Service.

(Authority: 38 U.S.C. 1155)

[29 FR 6718, May 22, 1964, as amended at 40 FR 42540, Sept. 15, 1975; 41 FR 11302, Mar. 18, 1976; 43 FR 45362, Oct. 2, 1978; 54 FR 4282, Jan. 30, 1989; 54 FR 49755, Dec. 1, 1989; 55 FR 154, Jan. 3, 1990; 56 FR 51653, Oct. 15, 1991; 57 FR 24364. June 9, 19921

#### MENTAL DISORDERS

#### §4.125 Diagnosis of mental disorders.

- (a) If the diagnosis of a mental disorder does not conform to DSM-IV or is not supported by the findings on the examination report, the rating agency shall return the report to the examiner to substantiate the diagnosis.
- (b) If the diagnosis of a mental disorder is changed, the rating agency shall determine whether the new diagnosis represents progression of the prior diagnosis, correction of an error in the prior diagnosis, or development of a new and separate condition. If it is not clear from the available records what the change of diagnosis represents, the rating agency shall return the report to the examiner for a determination.

(Authority: 38 U.S.C. 1155) [61 FR 52700, Oct. 8, 1996]

#### §4.126 Evaluation of disability from mental disorders.

- (a) When evaluating a mental disorder, the rating agency shall consider the frequency, severity, and duration of psychiatric symptoms, the length of remissions, and the veteran's capacity for adjustment during periods of remission. The rating agency shall assign an evaluation based on all the evidence of record that bears on occupational and social impairment rather than solely on the examiner's assessment of the level of disability at the moment of the examination.
- (b) When evaluating the level of disability from a mental disorder, the rating agency will consider the extent of social impairment, but shall not assign an evaluation solely on the basis of social impairment.
- (c) Delirium, dementia, and amnestic and other cognitive disorders shall be evaluated under the general rating formula for mental disorders; neurologic deficits or other impairments stemming from the same etiology (e.g., a head injury) shall be evaluated separately and combined with the evaluation for delirium, dementia, or amnestic or other cognitive disorder (see §4.25).
- (d) When a single disability has been diagnosed both as a physical condition

#### § 4.127

and as a mental disorder, the rating agency shall evaluate it using a diagnostic code which represents the dominant (more disabling) aspect of the condition (see § 4.14).

(Authority: 38 U.S.C. 1155) [61 FR 52700, Oct. 8, 1996]

### §4.127 Mental retardation and personality disorders.

Mental retardation and personality disorders are not diseases or injuries for compensation purposes, and, except as provided in §3.310(a) of this chapter, disability resulting from them may not be service-connected. However, disability resulting from a mental disorder that is superimposed upon mental retardation or a personality disorder may be service-connected.

(Authority: 38 U.S.C. 1155) [61 FR 52700, Oct. 8, 1996]

### §4.128 Convalescence ratings following extended hospitalization.

If a mental disorder has been assigned a total evaluation due to a continuous period of hospitalization lasting six months or more, the rating agency shall continue the total evaluation indefinitely and schedule a mandatory examination six months after the veteran is discharged or released to nonbed care. A change in evaluation based on that or any subsequent exam-

ination shall be subject to the provisions of §3.105(e) of this chapter.

(Authority: 38 U.S.C. 1155) [61 FR 52700, Oct. 8, 1996]

### §4.129 Mental disorders due to traumatic stress.

When a mental disorder that develops in service as a result of a highly stressful event is severe enough to bring about the veteran's release from active military service, the rating agency shall assign an evaluation of not less than 50 percent and schedule an examination within the six month period following the veteran's discharge to determine whether a change in evaluation is warranted.

(Authority: 38 U.S.C. 1155) [61 FR 52700, Oct. 8, 1996]

### § 4.130 Schedule of ratings—mental disorders.

The nomenclature employed in this portion of the rating schedule is based upon the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, of the American Psychiatric Association (DSM-IV). Rating agencies must be thoroughly familiar with this manual to properly implement the directives in §4.125 through §4.129 and to apply the general rating formula for mental disorders in §4.130. The schedule for rating for mental disorders is set forth as follows:

		Ratino
	Schizophrenia and Other Psychotic Disorders	
9201	Schizophrenia, disorganized type	
9202	Schizophrenia, catatonic type	
9203	Schizophrenia, paranoid type	
9204	Schizophrenia, undifferentiated type	
9205	Schizophrenia, residual type; other and unspecified types	
9208	Delusional disorder	
9210	Psychotic disorder, not otherwise specified (atypical psychosis)	
9211	Schizoaffective disorder	
	Delirium, Dementia, and Amnestic and Other Cognitive Disorders	
9300	Delirium	
9301	Dementia due to infection (HIV infection, syphilis, or other systemic or intracranial infections)	
9304	Dementia due to head trauma	
9305	Vascular dementia	
9310	Dementia of unknown etiology	
9312	Dementia of the Alzheimer's type	
9326	Dementia due to other neurologic or general medical conditions (endocrine disorders, metabolic disorders,	
Pick	c's disease, brain tumors, etc.) or that are substance-induced (drugs, alcohol, poisons)	
9327	Organic mental disorder, other (including personality change due to a general medical condition)	

§ 4.130

		Rating
	Anxiety Disorders	
9400	Generalized anxiety disorder	
9403	Specific (simple) phobia; social phobia	
9404		
9410		
9411	Post-traumatic stress disorder Panic disorder and/or agoraphobia	
	Anxiety disorder, not otherwise specified	
	Dissociative Disorders	
9416 9417		
	Somatoform Disorders	
9421	Somatization disorder	
9422		
9423		
9424 9425	Conversion disorder Hypochondriasis	
	Mood Disorders	
9431	Cyclothymic disorder	
9432		
9433		
9434		
9435	Mood disorder, not otherwise specified	
	Chronic Adjustment Disorder	
	Chronic adjustment disorder	
G	eneral Rating Formula for Mental Disorders:	
	Total occupational and social impairment, due to such symptoms as: gross impairment in thought proc-	
	esses or communication; persistent delusions or hallucinations; grossly inappropriate behavior; per- sistent danger of hurting self or others; intermittent inability to perform activities of daily living (including	
	maintenance of minimal personal hygiene); disorientation to time or place; memory loss for names of	
	close relatives, own occupation, or own name	10
	Occupational and social impairment, with deficiencies in most areas, such as work, school, family rela-	
	tions, judgment, thinking, or mood, due to such symptoms as: suicidal ideation; obsessional rituals which interfere with routine activities; speech intermittently illogical, obscure, or irrelevant; near-contin-	
	uous panic or depression affecting the ability to function independently, appropriately and effectively;	
	impaired impulse control (such as unprovoked irritability with periods of violence); spatial disorientation;	
	neglect of personal appearance and hygiene; difficulty in adapting to stressful circumstances (including	
	work or a worklike setting); inability to establish and maintain effective relationships	7
	flattened affect; circumstantial, circumlocutory, or stereotyped speech; panic attacks more than once a	
	week; difficulty in understanding complex commands; impairment of short- and long-term memory	
	(e.g., retention of only highly learned material, forgetting to complete tasks); impaired judgment; im-	
	paired abstract thinking; disturbances of motivation and mood; difficulty in establishing and maintaining	
	effective work and social relationships	
	Occupational and social impairment with occasional decrease in work efficiency and intermittent periods of inability to perform occupational tasks (although generally functioning satisfactorily, with routine be-	
	havior, self-care, and conversation normal), due to such symptoms as: depressed mood, anxiety, sus-	
	piciousness, panic attacks (weekly or less often), chronic sleep impairment, mild memory loss (such as	
	forgetting names, directions, recent events)	;
	Occupational and social impairment due to mild or transient symptoms which decrease work efficiency and ability to perform occupational tasks only during periods of significant stress, or; symptoms con-	
	trolled by continuous medication	
	A mental condition has been formally diagnosed, but symptoms are not severe enough either to interfere	
	with occupational and social functioning or to require continuous medication	

9520 Anorexia nervosa 9521 Bulimia nervosa

#### § 4.149

	Rating
Rating Formula for Eating Disorders:	
Self-induced weight loss to less than 80 percent of expected minimum weight, with incapacitating epi-	
sodes of at least six weeks total duration per year, and requiring hospitalization more than twice a year for parenteral nutrition or tube feeding	10
Self-induced weight loss to less than 85 percent of expected minimum weight with incapacitating epi- sodes of six or more weeks total duration per year	6
Self-induced weight loss to less than 85 percent of expected minimum weight with incapacitating epi- sodes of more than two but less than six weeks total duration per year	3
Binge eating followed by self-induced vomiting or other measures to prevent weight gain, or resistance to weight gain even when below expected minimum weight, with diagnosis of an eating disorder and incapacitating episodes of up to two weeks total duration per year	1
Binge eating followed by self-induced vomiting or other measures to prevent weight gain, or resistance to weight gain even when below expected minimum weight, with diagnosis of an eating disorder but with-	
out incapacitating episodes	

Note: An incapacitating episode is a period during which bed rest and treatment by a physician are required.

(Authority: 38 U.S.C. 1155)

[61 FR 52700, Oct. 8, 1996]

#### DENTAL AND ORAL CONDITIONS

#### §4.149 [Reserved]

### § 4.150 Schedule of ratings—dental and oral conditions.

	Rat- ing
9900 Maxilla or mandible, chronic osteomyelitis or osteoradionecrosis of: Rate as osteomyelitis, chronic under diagnostic	
code 5000.	
9901 Mandible, loss of, complete, between angles	100
9902 Mandible, loss of approximately one-half:	
Involving temporomandibular articulation	50
Not involving temporomandibular articulation	30
9903 Mandible, nonunion of:	
Severe	30
Moderate	10
NOTE—Dependent upon degree of motion and	
relative loss of masticatory function.	
9904 Mandible, malunion of:	_
Severe displacement	2
Moderate displacement	10
Slight displacement	(
relative loss of masticatory function.	
9905 Temporomandibular articulation, limited mo-	
tion of:	
Inter-incisal range:	
0 to 10 mm	4
11 to 20 mm	30
21 to 30 mm	20
31 to 40 mm	10
Range of lateral excursion:	
0 to 4 mm	1
Note—Ratings for limited inter-incisal movement	
shall not be combined with ratings for limited	
lateral excursion.	
9906 Ramus, loss of whole or part of:	
Involving loss of temporomandibular articulation Bilateral	5
Unilateral	3
Offinal Crail	3

	Rat- ing
Not involving loss of temporomandibular articula-	
tion	
Bilateral	3
Unilateral	2
9907 Ramus, loss of less than one-half the sub-	
stance of, not involving loss of continuity:	
Bilateral	2
Unilateral	1
9908 Condyloid process, loss of, one or both sides	3
9909 Coronoid process, loss of:	
Bilateral	2
Unilateral	1
9911 Hard palate, loss of half or more:	
Not replaceable by prosthesis	3
Replaceable by prosthesis	1
9912 Hard palate, loss of less than half of:	
Not replaceable by prosthesis	2
Replaceable by prosthesis	
9913 Teeth, loss of, due to loss of substance of	
body of maxilla or mandible without loss of con-	
tinuity:	
Where the lost masticatory surface cannot be re-	
stored by suitable prosthesis:  Loss of all teeth	4
Loss of all upper teeth	3
Loss of all lower teeth	3
All upper and lower posterior teeth missing	2
All upper and lower posterior teeth missing	2
All upper anterior teeth missing	1
All lower anterior teeth missing	1
All upper and lower teeth on one side miss-	'
ing	1
Where the loss of masticatory surface can be re-	
stored by suitable prosthesis	
Note—These ratings apply only to bone loss	
through trauma or disease such as osteomy-	
elitis, and not to the loss of the alveolar proc-	
ess as a result of periodontal disease, since	
such loss is not considered disabling.	
9914 Maxilla, loss of more than half:	
Not replaceable by prosthesis	10
Replaceable by prosthesis	5
9915 Maxilla, loss of half or less:	•

#### Pt. 4, App. A

	Rat- ing
Loss of 25 to 50 percent:	
Not replaceable by prosthesis	40
Replaceable by prosthesis	30
Loss of less than 25 percent:	
Not replaceable by prosthesis	
Replaceable by prosthesis	0
9916 Maxilla, malunion or nonunion of:	
Severe displacement	
Moderate displacement	
Slight displacement	

#### [59 FR 2530, Jan. 18, 1994]

#### APPENDIX A TO PART 4—TABLE OF AMEND-MENTS AND EFFECTIVE DATES SINCE 1946

#### 4.16 Last sentence; March 1, 1963. 4.17 October 7, 1948. March 1, 1963. 4.17a March 1, 1993. March 1, 1963. Paragraph (a) "first day of continuous hospitalization"; April 8, 1959. Paragraph (a) "terminated last day of month"; December 1, 1962. 4.29 Paragraph (a) penultimate sentence; November 13, 1970. Paragraph (b); April 8, 1959. Paragraph (c); August 16, 1948. Paragraph (d); August 16, 1948. Paragraph (e); March 1, 1963. Paragraph (f); August 9, 1976. NOTE: Application of this section to psychoneurotic and psychophysiologic disorders effective October 1, 1961. 4.30 Introductory portion of paragraph (a) preceding subparagraph (1); July 6, 1950. Paragraph (a)(1); June 9, 1952. Paragraph (a)(2); June 9, 1952. Paragraph (a)(3); June 9, 1952. Effective as to outpatient treatment March 10, 1976. Paragraph (b)(1); March 1, 1963. Paragraph (b)(2); August 9, 1976. Paragraph (b) first sentence; March 1, 1963. 4.55 4.63 June 17, 1948. October 1, 1956. 4.64 Diagnostic Code 5000—60 percent; February 1, 1962. Diagnostic Code 5000 NOTE (2): First three sentences; July 10, 1956. Last sentence; July 6, 1950. Diagnostic Code 5002—100 percent, 60 percent, 40 percent, 20 percent; March 1, 1963. Diagnostic Code 5003; July 6, 1950. Diagnostic Code 5012—Note; March 10, 1976.

In sentence following DC 5024: "except gout which will be rated under 5002"; March 1,

Diagnostic Code 5054; September 9, 1975.
Diagnostic Code 5055; September 9, 1975.
Diagnostic Code 5056; September 9, 1975.
Diagnostic Code 5056;
Diagnostic Code 5164—60 percent; June 9,

Diagnostic Code 5172; July 6, 1950. Diagnostic Code 5173; June 9, 1952. Diagnostic Code 5255 "or hip"; July 6, 1950.

1963. Diagnostic Code 5051; Diagnostic Code 5052; Diagnostic Code 5053;

1952.

#### APPENDIX A TO PART 4—TABLE OF AMEND-MENTS AND EFFECTIVE DATES SINCE 1946— Continued

Contir	
Sec.	
	Diagnostic Code 5257—Evaluation; July 6,
	1950.
	Diagnostic Code 5297—(Removal of one rib)
	"or resection of 2 or more"; August 23, 1948. Diagnostic Code 5297—Note (2): Reference to
	lobectomy; pneumonectomy and graduated
	ratings; February 1, 1962.
	Diagnostic Code 5298; August 23, 1948. Diagnostic Code 5324; February 1, 1962.
4.73	Diagnostic Code 5324; February 1, 1962.
	Diagnostic Code 5327; March 10, 1976.
	Diagnostic Code 5328; March 10, 1976.
4.78	Last sentence; December 1, 1963.
4.84a	Diagnostic Code 6029—Note; August 23, 1948.
	Diagnostic Code 6035; September 9, 1975.
	Diagnostic Code 6076—60%: Vision 1 eye 15/
	200 and other eye 20/100; August 23, 1948.
	Diagnostic Code 6080—Note—"as to 38 U.S.C.
	1114(L)"; July 6, 1950.
	Diagnostic Code 6081-Words "unilateral",
	"minimal" and all of Note; March 10, 1976.
4.84b	Removed-December 18, 1987 (text redesig-
	nated § 4.871, December 18, 1987)
4.85	March 23, 1956. December 18, 1987.
4.86	March 23, 1956. December 18, 1987.
4.86a	March 23, 1956. December 18, 1987.
4.87	Tables VI and VII replaced by new Tables VI
4.07	
4.070	VIa and VII December 18, 1987.
4.87a	Diagnostic Codes 6277 through 6297; March
	23, 1956; removed December 18, 1987. (Text
	from §4.84b redesignated §4.87a, December
	18, 1987).
4.88a	Diagnostic Code 6304—Notes (1) and (2); Au-
	gust 23, 1948.
	Diagnostic Code 6309; March 1, 1963.
	Diagnostic Code 6350; 80% Evaluation and Cri-
	Diagnostic Code 6350; 80% Evaluation and Criterion for 60% and 30% Evaluations; March
	10, 1976. Other Evaluations and Note; March
	1, 1963.
4.89	Ratings for nonpulmonary TB; December 1,
	1949.
4.97	Diagnostic Code 6600—100% Evaluations and
	Criteria for 60%; September 9, 1975.
	Diagnostic Code 6602—Criteria for all Evalua-
	tions and Note; September 9, 1975.
	Diagnostic Code 6603; September 9, 1975.
	Second note following Diagnostic Code 6724;
	December 1, 1949.
	Diagnostic Code 6802—Criteria for all Evalua-
	tions; September 9, 1975.
	Diagnostic Code 6819—Note; March 10, 1976.
	Diagnostic Code 6821—Evaluations and Note;
	August 23, 1948.
4.104	Diagnostic Code 7000-30 percent; July 6,
	1950.
	Diagnostic Code 7000—100 percent inactive
	"with signs of congestive failure upon any ex-
	ertion beyond rest in bed" revoked;
	Diagnostic Code 7005—80 percent revoked;
	Diagnostic Code 7005—80 percent revoked; Diagnostic Code 7007—80 percent revoked; Diagnostic Code 7015—100 percent Evaluation.
	Diagnostic Code 7015—100 percent Evaluation
	Criteria for All Evaluations and Notes (1) and
	(2); September 9, 1975.
	Diagnostic Code 7016; September 9, 1975.
	Diagnostic Code 7017;
	Diagnostic Code 7100—20 percent; July 6,
	1950.
	Diagnostic Code 7101 "or more"; September 1,
	1960.
	Diagnostic Code 7101—Note (2); September 9,
	1975

#### Pt. 4, App. B

APPENDIX A TO PART 4—TABLE OF AMEND-MENTS AND EFFECTIVE DATES SINCE 1946— Continued

Sec.	
	Diagnostic Code 7110—Criteria for 100 percent, NOTE and 60 percent and 20 percent Evalua- tions; September 9, 1975. Diagnostic Code 7111—NOTE; September 9,
	1975. Diagnostic Codes 7114, 7115, 7116, and NOTE; June 9, 1952.
	Diagnostic Code 7117 and NoTE; June 9, 1952.  NoTE following Diagnostic Code 7120; July 6, 1950.
	Diagnostic Code 7121—100 percent Criterion and Evaluation and 60 percent Criterion; March 10, 1976. Criteria for 30 percent and 10 percent and NoTE; July 6, 1950. Last sentence of NoTE following Diagnostic
4.114	Code 7122; July 6, 1950. Diagnostic Codes 7304 and 7305—Evaluations; November 1, 1962.
	Diagnostic Code 7308—Evaluations; April 8, 1959.
	Diagnostic Code 7312—70% Evaluation and 50% Evaluation and Criterion; March 10, 1976.
	Diagnostic Code 7313—20% Evaluation; March 10, 1976.
	Diagnostic Code 7319—Evaluations; November 1, 1962.
	Diagnostic Code 7321—Evaluations and Note; July 6, 1950.
	Diagnostic Code 7328—Evaluations and Note;
	November 1, 1962.  Diagnostic Code 7329—Evaluations and Note;
	November 1, 1962. Diagnostic Code 7330—60% Evaluation; No-
	vember 1, 1962. Diagnostic Code 7332—60% Evaluation; No-
	vember 1, 1962. Diagnostic Code 7334—50% and 30% Evaluations; July 6, 1950.
	Diagnostic Code 7334—10% Evaluation; No-
	vember 1, 1962. Diagnostic Code 7339—Criterion for 20% Eval-
	uation; March 10, 1976. Diagnostic Code 7343—Note; March 10, 1976. Diagnostic Code 7345—100%, 60% and 30%
	Evaluations; August 23, 1948.
	ruary 17, 1955. Diagnostic Code 7345—10% Evaluation; Feb-
	ruary 17, 1955. Diagnostic Code 7346—Evaluations; February 1, 1962.
	Diagnostic Code 7347; September 9, 1975.
4.115a	Diagnostic Code 7348; March 10, 1976. Diagnostic Code 7500—Note; July 6, 1950. Diagnostic Code 7519—20%, 40% and 60%
	Evaluations; March 10, 1976. Diagnostic Code 7524—Note; July 6, 1950. Diagnostic Code 7528—Note; March 10, 1976.
	Diagnostic Code 7528—Note; March 10, 1976. Diagnostic Code 7530; September 9, 1975.
4.116a	Diagnostic Code 7531; September 9, 1975.
4.117	Diagnostic Code 7627—Note; March 10, 1976. Diagnostic Code 7703—Evaluations; August 23, 1948.
	Diagnostic Code 7709—Note; March 10, 1976. Evaluations; June 9, 1952.
4.440	Diagnostic Code 7714; September 9, 1975.
4.118	Diagnostic Code 7801—Note (2); July 6, 1950. Diagnostic Code 7804—Note; July 6, 1950.

APPENDIX A TO PART 4—TABLE OF AMEND-MENTS AND EFFECTIVE DATES SINCE 1946— Continued

Sec.	
4.119	Diagnostic Code 7900—10% Evaluation; and Notes (2) and (3); August 13, 1981.
	Diagnostic Code 7902—20% Evaluation; August 13, 1981.
	Diagnostic Code 7903—10% Evaluation; August
	Diagnostic Code 7905—10% Evaluation; August 13. 1981.
	Diagnostic Code 7907—60% Evaluation; August 13, 1981.
	Diagnostic Code 7909—40% and 20% Evaluation; August 13, 1981.
	Diagnostic Code 7911—Evaluations and Note; March 1, 1963; 40% and 20% Evaluations; August 13, 1981.
	Diagnostic Code 7913—Note; September 9, 1975.
4.122	Diagnostic Code 7914—Note; March 10, 1976. October 1, 1961.
4.124a	Diagnostic Code 8002, NOTE;
	Diagnostic Code 8021, NOTE;
	Diagnostic Code 8045; October 1, 1961.
	Diagnostic Code 8046; October 1, 1961. Diagnostic Code 8100—Evaluations; June 9, 1953.
	Diagnostic Codes 8910 through 8914; October 1, 1961.
	Diagnostic Codes 8910 through 8914 General Rating Formula—Criteria and Evaluations; September 9, 1975.
4.125– 4.132	All Diagnostic Codes under Mental Disorders; October 1, 1961, except as to evaluation for Diagnostic Codes 9500 through 9511; Sep- tember 9, 1975.

[29 FR 6718, May 22, 1964, as amended at 34 FR 5064, Mar. 11, 1969; 40 FR 42541, Sept. 15, 1975; 41 FR 11291, Mar. 18, 1976; 41 FR 34258, Aug. 13, 1976; 43 FR 45362, Oct. 2, 1978; 46 FR 43666, Aug. 31, 1981; 52 FR 44122, Nov. 18, 1987; 52 FR 46439, Dec. 7, 1987]

### APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

Diag- nos- tic Code No.	
5000 5001	Osteomyelitis, acute, subacute, or chronic. Bones and Joints, tuberculosis of.
5001	
	Arthritis, rheumatoid (atrophic).
5003	Arthritis, degenerative, hypertrophic, or osteoarthritis.
5004	Arthritis, gonorrheal.
5005	Arthritis, pneumococcic.
5006	Arthritis, typhoid.
5007	Arthritis, syphilitic.
5008	Arthritis, streptococcic.
5009	Arthritis, other types.
5010	Arthritis, due to trauma.
5011	Bones, caisson disease of.
5012	Bones, new growths of, malignant.
5013	Osteoporosis, with joint manifestations.
5014	Osteomalacia.
5015	Bones, new growths of, benign.
	- · · · · · · · · · · · · · · · · · · ·

Pt. 4, App. B

APPENDIX B TO PART 4—NUMERICAL INDEX	OF
DISABILITIES—Continued	

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

Diag- nos- tic Code No.	
5016	Osteitis deformans.
5017	Gout.
5018	Hydrarthrosis, intermittent.
5019	Bursitis.
5020	Synovitis.
5021	Myositis.
5022	Periostitis.
5023	Myositis ossificans.
5024	Tenosynovitis.
	COMBINATIONS OF DISABILITIES

5100	Anatomical loss of both hands and both feet.
5101	Loss of use of both hands and both feet.
5102	Anatomical loss of both hands and one foot.
5103	Anatomical loss of both feet and one hand.
5104	Loss of use of both hands and one foot.
5105	Loss of use of both feet and one hand.
5106	Anatomical loss of both hands.
5107	Anatomical loss of both feet.
5108	Anatomical loss of one hand and one foot.
5109	Loss of use of both hands.
5110	Loss of use of both feet.
5111	Loss of use of one hand and one foot.

#### AMPUTATIONS: UPPER EXTREMITY

```
Arm, amputation of:
                     Disarticulation.
                     Above insertion of deltoid.
Below insertion of deltoid.
5121
5122
             Forearm, amputation of:
Above insertion of pronator teres.
Below insertion of pronator teres.
5123
5124
             Hand, loss of use of.
Five digits of one hand, amputation of:
Four digits of one hand, amputation of:
5125
5126
                     Thumb, index, middle and ring.
Thumb, index, middle and little.
Thumb, index, ring and little.
5127
5128
5129
             Thumb, middle, ring and little.
Index, middle, ring and little.
Three digits of one hand, amputation of:
5130
5131
5132
5133
                     Thumb, index and middle.
Thumb, index and ring.
Thumb, index and little.
5134
5135
5136
                     Thumb, middle and ring.
Thumb, middle and little.
5137
                     Thumb, ring and little.
5138
5139
                     Index, middle and ring.
Index, middle and little.
5140
                     Index, ring and little.
5141
              Middle, ring and little.
Two digits of one hand, amputation of:
                     Thumb and index.
Thumb and middle.
Thumb and ring.
5142
5143
5144
5145
                     Thumb and little.
5146
5147
                     Index and middle.
Index and ring.
5148
                     Index and little.
                     Middle and ring.
Middle and little.
5149
5150
5151
                     Ring and little.
             Thumb, amputation of.
5152
```

### APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES—Continued

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

Diag- nos- tic Code No.	
5153	Index finger, amputation of.
5154	Middle finger, amputation of.
5155	Ring finger, amputation of.
5156	Little finger, amputation of.

#### AMPUTATIONS: LOWER EXTREMITY

-	
	Thigh, amputation of:
5160	Disarticulation.
5161	Upper third.
5162	Middle or lower thirds.
	Leg, amputation of:
5163	With defective stump.
5164	With loss of natural knee action.
5165	At a lower level.
5166	Forefoot, amputation proximal to metatarsal bones.
5167	Foot, loss of use of.
5170	Toes, all, amputation of, without metatarsal loss.
5171	Toe, great, amputation of.
5172	Toe, other, amputation of.
5173	Toes, three or more, amputation of, not including
	great toe.

#### THE SHOULDER AND ARM

5200 5201	Scapulohumeral articulation, ankylosis of. Arm, limitation of motion of.
5202	Humerus, other impairment of.
5203	Clavicle or scapula, impairment of.

#### THE ELBOW AND FOREARM

5205	Elbow, ankylosis of.
5206	Forearm, limitation of flexion of.
5207	Forearm, limitation of extension of
5208	Forearm, flexion limited to 100° and extension to 45°.
5209	Elbow, other impairment of.
5210	Radius and ulna, nonunion of, with flail false joint.
5211	Ulna, impairment of.
5212	Radius, impairment of.
5213	Supination and pronation, impairment of.
	THE WRIST AND HAND

5214	Wrist, ankylosis.
5215	Wrist, limitation of motion of.
5216	Five digits of one hand, unfavorable ankylosis of.
5217	Four digits of one hand, unfavorable ankylosis of.
5218	Three digits of one hand, unfavorable ankylosis of.
5219	Two digits of one hand, unfavorable ankylosis of.
5220	Five digits of one hand, favorable ankylosis of.
5221	Four digits of one hand, favorable ankylosis of.
5222	Three digits of one hand, favorable ankylosis of.
5223	Two digits of one hand, favorable ankylosis of.
5224	Thumb, ankylosis of.
5225	Index finger, ankylosis of.
5226	Middle finger, ankylosis of.
5227	Finger, any other, ankylosis of.
	THE HIR AND THICH

#### THE HIP AND THIGH

5250	Hip, ankylosis of.
5251	Thigh, limitation of extension of
5252	Thigh, limitation of flexion of.

#### 38 CFR Ch. I (7-1-99 Edition)

#### Pt. 4, App. B

# APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES—Continued

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

Spine, complete bony fixation (ankylosis) of. Spine, ankylosis of, cervical. Spine, ankylosis of, dorsal. Spine, ankylosis of, lumbar. Spine, limitation of motion of, cervical. Spine, limitation of motion of, dorsal. Spine, limitation of motion of, lumbar. Intervertebral disc syndrome. Sacrolliac injury and weakness. Lumbosacral strain.  THE SKULL  Skull, loss of part of, both inner and outer tables.
Spine, ankylosis of, cervical. Spine, ankylosis of, dorsal. Spine, ankylosis of, durbar. Spine, limitation of motion of, cervical. Spine, limitation of motion of, dorsal. Spine, limitation of motion of, durbar. Intervertebral disc syndrome. Sacrolilac injury and weakness. Lumbosacral strain.
Spine, ankylosis of, cervical. Spine, ankylosis of, dorsal. Spine, ankylosis of, lumbar. Spine, limitation of motion of, cervical. Spine, limitation of motion of, dorsal. Spine, limitation of motion of, lumbar. Intervertebral disc syndrome. Sacroiliac injury and weakness.
Spine, ankylosis of, cervical. Spine, ankylosis of, dorsal. Spine, ankylosis of, lumbar. Spine, limitation of motion of, cervical. Spine, limitation of motion of, dorsal. Spine, limitation of motion of, lumbar. Intervertebral disc syndrome.
Spine, ankylosis of, cervical. Spine, ankylosis of, dorsal. Spine, imitation of motion of, cervical. Spine, limitation of motion of, dorsal. Spine, limitation of motion of, lumbar.
Spine, ankylosis of, cervical. Spine, ankylosis of, dorsal. Spine, ankylosis of, lumbar. Spine, limitation of motion of, cervical. Spine, limitation of motion of, dorsal.
Spine, ankylosis of, cervical. Spine, ankylosis of, dorsal. Spine, ankylosis of, lumbar.
Spine, ankylosis of, cervical. Spine, ankylosis of, dorsal.
Spine, ankylosis of, cervical.
Spine, complete bony fixation (ankylosis) of.
Vertebra, fracture of, residuals.
THE SPINE
Foot injuries, other.
of.
Hammer toe.  Tarsal, or metatarsal bones, malunion of, or nonunion
Hallux rigidus.
Hallux valgus.
Claw foot (pes cavus), acquired.  Metatarsalgia, anterior (Morton's disease).
Weak foot, bilateral.
Flatfoot, acquired.
THE FOOT
Bones, of the lower extremity, shortening of.
SHORTENING OF THE LOWER EXTREMITY
Astragalectomy.
Os calcis or astragalus, malunion of.
Subastragalar or tarsal joint, ankylosis of.
Ankle, limited motion of.
Ankle, ankylosis of.
THE ANKLE
Genu recurvatum.
Tibia and fibula, impairment of.
Leg, limitation of flexion of. Leg, limitation of extension of.
Cartilage, semilunar, removal of.
Cartilage, semilunar, dislocated.
Knee, other impairment of.
Knee, ankylosis of.
THE KNEE AND LEG
Hip, flail joint. Femur, impairment of.
Thigh, impairment of.

5297 Ribs, removal of.

## APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES—Continued

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

Diag- nos- tic Code No.	
	THE COCCYX
5298	Coccyx, removal of.
	MUSCLE INJURIES
5301	Group I—Extrinsic muscles of shoulder girdle.
5302	Group II—Extrinsic muscles of shoulder girdle.
5303	Group III—Intrinsic muscles of shoulder girdle.
5304	Group IV—Intrinsic muscles of shoulder girdle.
5305	Group V—Flexor muscles of the elbow.
5306	Group VI—Extensor muscles of the elbow.
5307	Group VII—Muscles arising from internal condyle of humerus.
5308	Group VIII—Muscles arising mainly from external
	condyle of humerus.
5309	Group IX—Intrinsic muscles of the hand.
5310	Group X—Intrinsic muscles of the foot.
5311	Group XI—Posterior and lateral muscles of the leg.
5312	Group XII—Anterior muscles of the leg.
5313	Group XIII—Posterior thigh group.
5314	Group XIV—Anterior thigh group.
5315	Group XV—Mesial thigh group.
5316	Group XVI—Pelvic girdle group 1.
5317 5318	Group XVII—Pelvic girdle group 2. Group XVIII—Pelvic girdle group 3.
5319	Group XVII—Pelvic girdle group 3.  Group XIX—Muscles of the abdominal wall.
5320	Group XX—Spinal muscles.
5321	Group XXI—Muscles of respiration.
5322	Group XXII—Lateral, supra and infrahyoid group.
5323	Group XXIII—Lateral and posterior muscles of the neck.
5324	Diaphragm, rupture of.
5325	Muscle injury, facial muscles.
5326	Muscle hernia.

#### DISEASES OF THE EYE

6000 Uveitis.

6001	Keratitis.
6002	Scleritis.
6003	Iritis.
6004	Cyclitis.
6005	Choroiditis.
6006	Retinitis.
6007	Hemorrhage, intra-ocular, recent.
6008	Retina, detachment of.
6009	Eye, injury of, unhealed.
6010	Eye, tuberculosis of.
6011	Retina, localized scars.
6012	Glaucoma, congestive or inflammatory.
6013	Glaucoma, simple, primary, noncongestive.
6014	New growths, malignant, eyeball.
6015	New growths, benign, eyeball and adnexa.
6016	Nystagmus, central.
6017	Conjunctivitis, trachomatous, chronic.
6018	Conjunctivitis, other, chronic.
6019	Ptosis, eyelids.
6020	Ectropion.
6021	Entropion.
6022	Lagophthalmos.
6023	Eyebrows, loss of.
6024	Eyelashes, loss of.
6025	Epiphora.
6026	Neuritis, optic.

Pt. 4, App. B

APPENDIX B TO PART 4—NUMERICAL INDEX C	)F
DISABILITIES—Continued	

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

Diag- nos- tic Code No.	
6027	Cataract, traumatic.
6028 6029	Cataract, senile, and others.  Aphakia.
6030	Accommodation, paralysis of.
6031	Dacryocystitis.
6032	Eyelids, loss of portion of.
6033 6034	Lens, crystalline, dislocation of. Pterygium.
	COMBINATIONS OF DISABILITIES
6050	Blindness in both eyes having only light perception and anatomical loss of both hands and both feet.
6051	Blindness in both eyes having only light perception and loss of use of both hands and both feet.
6052	Blindness in both eyes having only light perception and anatomical loss of both hands.
6053	Blindness in both eyes having only light perception and anatomical loss of both feet.
6054	Blindness in both eyes having only light perception and anatomical loss of one hand and one foot.
6055	Blindness in both eyes having only light perception and loss of use of both hands.
6056	Blindness in both eyes having only light perception and loss of use of both feet.
6057	Blindness in both eyes having only light perception and loss of use of one hand and one foot.
6058	Blindness in both eyes having only light perception and anatomical loss of one hand.
6059	Blindness in both eyes having only light perception and anatomical loss of one foot.
6060	Blindness in both eyes having only light perception and loss of use of one hand.
6061	Blindness in both eyes having only light perception and loss of use of one foot.
6062	Blindness in both eyes having only light perception.

#### IMPAIRMENT OF CENTRAL VISUAL ACUITY

```
Blindness, anatomical loss, one eye:
6063
6064
                     Other blind (5/200 or less).
Other impaired (20/200 or less).
6065
                     Other impaired.
             Other Impaired.
Other normal.
Blindness, light perception only one eye:
Other blind (5/200 or less).
Other impaired (20/200 or less).
Other impaired.
6066
6067
6068
6069
             Other normal.

Blindness, total (5/200 or less):
Both eyes.
6070
6071
             Blindness, total one eye (5/200 or less):
Other impaired (20/200 or less).
Other impaired.
6072
6073
             Other normal.

Blindness, partial (20/200 or less):
Both eyes.
6074
6075
                     One eye:
Other impaired.
Other normal.
6076
6077
             Blindness, partial:
6078
                     Both eyes.
One eye only.
6079
             Field vision, impairment of. Scotoma, pathological.
6080
6081
```

### APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES—Continued

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

	[AGGTE, GOBAGGTE, OK GINGING BIGEAGEG]
Diag-	
nos-	
tic	
Code No.	
INU.	
6090	Muscle function, ocular, impairment of.
6091	Symblepharon.
6092	Diplopia, due to limited muscle function.
	IMPAIRMENT OF AUDITORY ACUITY
	IMPAIRMENT OF AUDITORY ACUITY
6100	0% evaluation based on Table VII
6101	10% evaluation based on Table VII
6102	20% evaluation based on Table VII
6103	30% evaluation based on Table VII
6104	40% evaluation based on Table VII
6105	50% evaluation based on Table VII
6106	60% evaluation based on Table VII
6107 6108	70% evaluation based on Table VII
6109	80% evaluation based on Table VII 90% evaluation based on Table VII
6110	100% evaluation based on Table VII.
	DISEASES OF THE EAR
6200	Otitio modio suppurativo abrania
6201	Otitis media, suppurative, chronic. Otitis media, catarrhal, chronic.
6202	Otosclerosis.
6203	Otitis interna.
6204	Labyrinthitis.
6205	Meniere's syndrome.
6206	Mastoiditis.
6207	Auricle, loss or deformity.
6208	New growths, malignant, ear.
6209	New growths, benign, ear.
6210	Auditory canal, disease of.
6211	Tympanic membrane, perforation of.
6260	Tinnitus.
	OTHER SENSE ORGANS
6275	Smell, loss of sense of.
6276	Taste, loss of sense of.
	SYSTEMIC DISEASES
6300	Choloro Asiatio
6301	Cholera, Asiatic. Kala-azar (visceral leishmaniasis).
6302	Leprosy.
6304	Malaria.
6305	Filariasis.
6306	Oroya fever.
6307	Plague.
6308	Relapsing fever.
6309	Rheumatic fever.
6310	Syphilis, unspecified.
6311	Tuberculosis, military.
6313	Avitaminosis.
6314	Beriberi.
6315	Pellagra.
6316	Brucellosis (Malta or undulant fever).
6317	Typhus, scrub.
6350	Lupus erythematosus, systemic.
	RESPIRATORY SYSTEM
	THE NOSE AND THROAT

6501 Rhinitis, atrophic, chronic.

#### 38 CFR Ch. I (7-1-99 Edition)

#### Pt. 4, App. B

## APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES—Continued

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

	[ACUTE, SUBACUTE, OR CHRONIC DISEASES]
Diag-	
nos-	
tic	
Code	
No.	
6502	Septum, nasal, deflection of.
6504	Nose, loss of part of, or scars.
6510	Sinusitis, pansinusitis, chronic.
6511	Sinusitis, ethmoid, chronic.
6512	Sinusitis, frontal, chronic.
6513	Sinusitis, maxillary, chronic.
6514	Sinusitis, sphenoid, chronic.
6515	Laryngitis, tuberculous.
6516	Laryngitis, chronic.
6517	Larynx, injuries of, healed.
6518	Laryngectomy.
6519	Aphonia, organic.
6520	Larynx, stenosis of.
	THE TRACHEA AND BRONCHI
6600	Bronchitis, chronic.
6601	Bronchiectasis.
6602	Asthma, bronchial.
	THE LUNGS AND PLEURA
6701	Tuberculosis, pulmonary, chronic, far advanced, active.
6702	Tuberculosis, pulmonary, chronic, moderately advanced, active.
6703	Tuberculosis, pulmonary, chronic, minimal, active.
6704	Tuberculosis, pulmonary, chronic, active, advance-
	ment unspecified.
6707	Tuberculosis, pulmonary, chronic, far advanced, active.
6708	Tuberculosis, pulmonary, chronic, moderately ad-
0.00	vanced, active.
6709	Tuberculosis, pulmonary, chronic, minimal, active.
6710	Tuberculosis, pulmonary, chronic, active, advance-
	ment unspecified.
6721	Tuberculosis, pulmonary, chronic, far advanced, inac-
	_ tive.
6722	Tuberculosis, pulmonary, chronic, moderately ad-
076-	vanced, inactive.
6723	Tuberculosis, pulmonary, chronic, minimal, inactive.
6724	Tuberculosis, pulmonary, chronic, inactive, advance-
C70E	ment unspecified.
6725	Tuberculosis, pulmonary, chronic, far advanced, inactive.
6726	Tuberculosis, pulmonary, chronic, moderately ad-
JU	vanced, inactive.
6727	Tuberculosis, pulmonary, chronic, minimal, inactive.
6728	Tuberculosis, pulmonary, chronic, inactive, advance-
	ment unspecified.
6732	Pleurisy, tuberculous.
6800	Anthracosis.
6801	Silicosis.
6802	Pneumoconiosis, unspecified.
6803	Actinomycosis of lung.
6804	Streptotrichosis of lung.
6805	Blastomycosis of lung.
6806	Sporotrichosis of lung.
6807	Aspergillosis of lung.
6808	Mycosis of lung, unspecified.
6809	Lung, abscess of.
6810 6811	Pleurisy, serofibrinous.
6812	Pleurisy, purulent (empyema). Fistula, bronchocutaneous, or bronchopleural.
6813	Lung, permanent collapse of.
3010	

## APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES—Continued

	DISABILITIES—Continued
	[ACUTE, SUBACUTE, OR CHRONIC DISEASES]
Diag-	
nos-	
tic	
Code	
No.	
6814	Pneumothorax, spontaneous.
6815	Pneumonectomy.
6816	Lobectomy.
6817	Lung, chronic passive congestion of.
6818	Pleural cavity, injuries, residuals of, including gunshot wounds.
6819	New growths, malignant, any specified part of res- piratory system.
6820	New growths, benign, any specified part of respiratory system.
6821	Coccidioidomycosis.
	THE CARDIOVASCULAR SYSTEM
	THE HEART
7000	Rheumatic heart disease.
7001	Endocarditis, bacterial, subacute.
7002	Pericarditis, bacterial or rheumatic, acute.
7003	Adhesions, pericardial.
7004	Syphilitic heart disease.
7005	Arteriosclerotic heart disease.
7006	Myocardium, infarction of, due to thrombosis or embolism.
7007	Hypertensive heart disease.
7008	Hyperthyroid heart disease.
7010	Auricular flutter, paroxysmal.  Auricular fibrillation, paroxysmal.
7011 7012	Auricular librillation, paroxysmai.  Auricular fibrillation, permanent.
7012	Tachycardia, paroxysmal.
7013	Sinus tachycardia.
7015	Auriculoventricular block.
	THE ARTERIES AND VEINS
7100	Arteriosclerosis, general.
7101	Hypertensive vascular disease (essential arterial hy-
	pertension).
7110	Aorta or branches, aneurysm of.
7111	Artery, any large artery, aneurysm of.
7112	Artery, small aneurysmal dilatation.
7113	Arteriovenous aneurysm, traumatic.
7114	Arteriosclerosis obliterans.
7115	Thrombo-angiitis obliterans (Buerger's disease).
7116	Claudication, intermittent.
7117	Raynaud's disease.
7118	Angioneurotic edema.
7119 7120	Erythromelalgia.
7121	Varicose veins. Phlebitis.
7122	Frozen feet, residuals of (Immersion foot).
	THE DIGESTIVE SYSTEM
7200	Mouth, injuries of.
7201	Lips, injuries of.
7202	Tongue, loss of, whole or part.
7203	Esophagus, stricture of.
7204	Esophagus, spasm of (cardiospasm).
7205	Esophagus, diverticulum of, acquired.
7301	Peritoneum, adhesions of.
7304	Ulcer, gastric.
7305 7306	Ulcer, duodenal. Ulcer, marginal (gastrojejunal).
7300	Gastritis, hypertrophic.

Pt. 4, App. B

# APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES—Continued

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

Diag- nos-	
tic	
Code	
No.	
7308	Postgastrectomy syndromes.
7309	Stomach, stenosis of.
7310	Stomach, injury of, residuals.
7311	Liver, injury of.
7312	Liver, cirrhosis of.
7313	Liver, abscess of, residuals.
7314	Cholecystitis, chronic.
7315	Cholelithiasis, chronic.
7316	Cholangitis, chronic.
7317	Gall bladder, injury of.
7318	Gall bladder, removal of.
7319	Irritable colon syndrome (spastic colitis, mucous coli tis, etc.).
7321	Amebiasis.
7322	Dysentery, bacillary.
7323	Colitis, ulcerative.
7324	Distomiasis, intestinal or hepatic.
7325	Enteritis, chronic.
7326	Enterocolitis, chronic.
7327	Diverticulitis.
7328	Intestine, small, resection of.
7329	Intestine, large, resection of.
7330	Intestine, fistula of.
7331	Peritonitis, tuberculous, active.
7332	Rectum and anus, impairment of sphincter control.
7333	Rectum and anus, stricture of.
7334	Rectum, persistent prolapse of.
7335	Ano, fistula in.
7336	Hemorrhoids, external or internal.
7337	Pruritus ani.
7338	Hernia, inguinal.
7339	
7340	Hernia, femoral.
7341	Wounds, incised, healed, abdominal wall.
7342	Visceroptosis.
7343	New growths, malignant, any specified part of diges tive system.
7344	New growths, benign, any specified part of digestive system.
7345	Hepatitis, infectious.
7346	Hernia, hiatal.

#### THE GENITOURINARY SYSTEM

7500	Kidney, removal of.
7501	Kidney, abscess of.
7502	Nephritis, chronic.
7503	Pyelitis.
7504	Pyelonephritis, chronic.
7505	Kidney, tuberculosis of, active.
7507	Nephrosclerosis, arteriolar.
7508	Nephrolithiasis.
7509	Hydronephrosis.
7510	Ureterolithiasis.
7511	Ureter, stricture of.
7512	Cystitis, chronic.
7513	Cystitis, interstitial (Hunner), submucous or elusive ulcer.
7514	Bladder, tuberculosis of.
7515	Bladder, calculus in.
7516	Bladder, fistula of.
7517	Bladder, injury of.
7518	Urethra, stricture of.
7519	Urethra, fistula of.
7520	Penis, removal of half or more.
7521	Penis, removal of glans.

# APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES—Continued

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

Diag- nos- tic Code No.	
7522	Penis, deformity, with loss of erectile power.
7523	Testis, atrophy, complete.
7524	Testis, removal of.
7525	Epididymo-orchitis (tuberculous).
7526	Prostate gland, resection or removal.
7527	Prostate gland injuries, infectious hypertrophy, post- operative residuals.
7528	New growths, malignant, any specified part of genito- urinary system.
7529	New growths, benign, any specified part of genito- urinary system.
	GYNECOLOGICAL CONDITIONS

7610	Vulvovaginitis.
7611	Vaginitis.
7612	Cervicitis.
7613	Metritis.
7614	Salpingitis.
7615	Oophoritis.
7617	Uterus and ovaries, removal of, complete.
7618	Uterus, removal of, including corpus.
7619	Ovaries, removal of.
7620	Ovaries, atrophy of both.
7621	Uterus, prolapse.
7622	Uterus, displacement of.
7623	Pregnancy, surgical complications of.
7624	Fistula, rectovaginal.
7625	Fistula, urethrovaginal.
7626	Mammary glands, removal of.
7627	New growth, malignant, gynecological system, or
. 02.	mammary glands.
	THE HEMIC AND LYMPHATIC SYSTEMS

7700	Anemia, pernicious.
7701	Anemia, secondary.
7702	Agranulocytosis, acute.
7703	Leukemia.
7704	Polycythemia, primary.
7705	Purpura hemorrhagica.
7706	Splenectomy.
7707	Spleen, injury of, healed.
7709	Lymphogranulomatosis (Hodgkin's disease).
7710	Adenitis, cervical, tuberculous.
7711	Adenitis, axillary, tuberculous.
7712	Adenitis, inguinal, tuberculous.
7713	Adenitis, secondary.
	THE SKIN
7800	Scars, disfiguring, head, face or neck.
7801	Scars, burns, third degree.
7802	Scars, burns, second degree.
7803	Scars, superficial, poorly nourished.
7804	Scars, superficial, tender and painful.
7805	Scars, others.
7806	Eczema.
7807	Leishmaniasis, americana (mucocutaneous, espundia).
7808	Leishmaniasis, old world (cutaneous, oriental sore).
7809	Lupus erythematosus, discoid.
	Pinta.
7810	Pinta.
7810 7811	Tuberculosis luposa (lupus vulgaris).

#### 38 CFR Ch. I (7-1-99 Edition)

#### Pt. 4, App. B

## APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES—Continued

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

	• • •
Diag- nos- tic Code No.	
7813	Dermatophytosis.
7814	Tinea barbae.
7815	Pemphigus.
7816	Psoriasis.
7817	Dermatitis exfoliativa.
7818	New growths, malignant, skin.
7819	New growths, benign, skin.
	THE ENDOCRINE SYSTEM
7900	Hyperthyroidism

7900	Hyperthyroidism.
7901	Thyroid gland, toxic adenoma of.
7902	Thyroid gland, non-toxic adenoma of.
7903	Hypothyroidism.
7904	Hyperparathyroidism (osteitis fibrosa cystica).
7905	Hypoparathyroidism.
7907	Hyperpituitarism (pituitary basophilism, Cushing's syn-
	drome).
7908	Hyperpituitarism (acromegaly or gigantism).
7909	Hypopituitarism (diabetes inspidus).
7910	Hyperadrenia (adrenogenital syndrome).
7911	Addison's disease.
7912	Pluriglandular syndromes.
7913	Diabetes mellitus.
7914	New growths, malignant, endocrine system.
7915	New growths, benign, endocrine system.

#### NEUROLOGICAL CONDITIONS AND CONVULSIVE DISORDERS

8000	Encephalitis, epidemic, chronic.
	Brain, new growth of:
8002	Malignant.
8003	Benign.
8004	Paralysis agitans.
8005	Bulbar palsy.
8007	Brain, vessels, embolism of.
8008	Brain, vessels, thrombosis of.
8009	Brain, vessels, hemorrhage from.
8010	Myelitis.
8011	Poliomyelitis, anterior.
8012	Hematomyelia.
8013	Syphilis, cerebrospinal.
8014	
8015	Tabes dorsalis.
8017	Amyotrophic lateral sclerosis.
8018	Multiple sclerosis.
8019	
8020	
8021	Malignant.
8022	
8023	
8024	
8025	,
8045	
	Cerebral arteriosclerosis.
	Migraine.
	Tic, convulsive.
8104	Paramyoclonus multiplex (convulsive state, myoclonic type).
8105	Chorea, Sydenham's.
8106	Chorea, Huntington's.
8107	Athetosis, acquired.
8108	Narcolepsy.

## APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES—Continued

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

Diag- nos- tic Code No.	
	THE CRANIAL NERVES
8205	Fifth (trigeminal) cranial nerve, paralysis of.
8207	Seventh (facial) cranial nerve, paralysis of.
8209	Ninth (glossopharyngeal) cranial nerve, paralysis of.
8210	Tenth (pneumogastric, vagus) cranial nerve, paralysis of.
8211	Eleventh (spinal accessory, external branch) cranial nerve, paralysis of.
8212	Twelfth (hypoglossal) cranial nerve, paralysis of.
8305	Fifth (trigeminal) cranial nerve, neuritis.
8307	Seventh (facial) cranial nerve, neuritis.
8309	Ninth (glossopharyngeal) cranial nerve, neuritis.
8310	Tenth (pneumogastric, vagus) cranial nerve, neuritis.
8311	Eleventh (spinal accessory, external branch) cranial nerve, neuritis.
8312	Twelfth (hypoglossal) cranial nerve, neuritis.
8407	Seventh (facial) cranial nerve, neuralgia.
8409	Ninth (glossopharyngeal) cranial nerve, neuralgia.
8410	Tenth (pneumogastric, vagus) cranial nerve, neuralgia.
8411	Eleventh (spinal accessory, external branch) cranial nerve, neuralgia.
8412	Twelfth (hypoglossal) cranial nerve, neuralgia.
	PERIPHERAL NERVES: PARALYSIS

8510	Upper radicular group (fifth and sixth cervicals), paral- vsis of.
8511	Middle radicular group, paralysis of.
8512	Lower radicular group, paralysis of.
8513	All radicular groups, paralysis of.
8514	The musculospiral nerve (radial nerve), paralysis of.
8515	The median nerve, paralysis of.
8516	The ulnar nerve, paralysis of.
8517	Musculocutaneous nerve, paralysis of.
8518	Circumflex nerve, paralysis of.
8519	Long thoracic nerve, paralysis of.
8520	The sciatic nerve, paralysis of.
8521	External popliteal nerve (common peroneal), paralysis
	of.
8522	Musculocutaneous nerve (superficial peroneal), paral- ysis of.
8523	Anterior tibial nerve (deep peroneal), paralysis of.
8524	Internal popliteal nerve (tibial), paralysis of.
8525	Posterior tibial nerve, paralysis of.
8526	Anterior crural nerve (femoral), paralysis of.
8527	Internal saphenous nerve, paralysis of.
8528	Obturator nerve, paralysis of.
8529	External cutaneous nerve of thigh, paralysis of.
8530	Ilio-inguinal nerve, paralysis of.

	PERIPHERAL NERVES: NEURITIS
8610	Upper radicular group (fifth and sixth cervicals), ne ritis.
8611	Middle radicular group, neuritis.
8612	Lower radicular group, neuritis.
8613	All radicular groups, neuritis.
8614	The musculospiral nerve (radial nerve), neuritis.
8615	The median nerve, neuritis.
8616	The ulnar nerve, neuritis.
8617	Musculocutaneous nerve, neuritis.
8618	Circumflex nerve, neuritis.
8619	Long thoracic nerve neuritis

### Pt. 4, App. B

#### **Department of Veterans Affairs**

## APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES—Continued

Diag- nos- tic	
Code No.	
8620	The sciatic nerve, neuritis.
8621 8622	External popliteal nerve (common peroneal), neuritis.  Musculocutaneous nerve (superficial peroneal), neuritis.
8623	Anterior tibial nerve (deep peroneal), neuritis.
8624	Internal popliteal nerve (tibial) neuritis.
8625	Posterior tibial nerve, neuritis.
8626	Anterior crural nerve (femoral), neuritis.
8627	Internal saphenous nerve, neuritis.
8628	Obturator nerve, neuritis.
8629	External cutaneous nerve of thigh, neuritis.
8630	Ilio-inguinal nerve, neuritis.
	PERIPHERAL NERVES: NEURALGIA
8710	Upper radicular group (fifth and sixth cervicals), neu ralgia.
8711	Middle radicular group, neuralgia.
8712	Lower radicular group, neuralgia.
8713	All radicular groups, neuralgia.
8714	The musculospiral nerve (radial nerve), neuralgia.
8715	The median nerve, neuralgia.
8716	The ulnar nerve, neuralgia.
8717	Musculocutaneous nerve, neuralgia.
8718	Circumflex nerve, neuralgia.
8719	Long thoracic nerve, neuralgia.
8720	The sciatic nerve, neuralgia.
8721	External popliteal nerve (common peroneal), neu ralgia.
8722	Musculocutaneous nerve (superficial peroneal), neu ralgia.
8723	Anterior tibial nerve (deep peroneal), neuralgia.
8724	Internal popliteal nerve (tibial), neuralgia.
8725	Posterior tibial nerve, neuralgia.
8726	Anterior crural nerve (femoral), neuralgia.
8727	Internal saphenous nerve, neuralgia.
8728	Obturator nerve, neuralgia.
8729	External cutaneous nerve of thigh neuralgia.
8730	Ilio-inguinal nerve, neuralgia.

#### THE EPILEPSIES

8910	Epilepsy, grand mal.
8911	Epilepsy, petit mal.
8912	Jacksonian type.
8913	Epilepsy, diencephalic.
8914	Epilepsy, psychomotor.
	PSYCHOTIC DISORDERS
9200	Schizophrenic reaction, simple type.
9201	Schizophrenic reaction, hebephrenic type.
9202	Schizophrenic reaction, catatonic type.
9203	Schizophrenic reaction, paranoid type.
9204	Schizophrenic reaction, chronic undifferentiated type.
9205	Schizophrenic reaction, other.
9206	Manic depressive reaction.
9207	Psychotic depressive reaction.
9208	Paranoid reaction (specify).
9209	Involutional psychotic reaction.
9210	Psychotic reaction, other.

# APPENDIX B TO PART 4—NUMERICAL INDEX OF DISABILITIES—Continued

[ACUTE, SUBACUTE, OR CHRONIC DISEASES]

Diag- nos- tic Code No.	
	ORGANIC BRAIN DISORDERS
9300	Acute brain syndrome (associated with infection, trauma, circulatory disturbance, etc.).
9301	Chronic brain syndrome associated with central nervous system syphilis (all forms).
9302	Chronic brain syndrome associated with intracranial infections other than syphilis.
9303	Chronic brain syndrome associated with intoxication.
9304	Chronic brain syndrome associated with brain trauma.
9305	Chronic brain syndrome associated with cerebral arteriosclerosis.
9306	Chronic brain syndrome associated with circulatory disturbance other than cerebral arteriosclerosis.
9307	Chronic brain syndrome associated with convulsive disorder (idiopathic epilepsy).
9308	Chronic brain syndrome associated with disturbance of metabolism, growth or nutrition.
9309	Chronic brain syndrome associated with intracranial neoplasm.
9310	Chronic brain syndrome associated with diseases of unknown or uncertain cause.
9311	Chronic brain syndrome of unknown cause.
	PSYCHONEUROTIC DISORDERS
9400	Anxiety reaction.
9401	Dissociative reaction.
9402	Conversion reaction.

9401	Dissociative reaction.
9402	Conversion reaction.
9403	Phobic reaction.
9404	Obsessive compulsive reaction.
9405	Depressive reaction.
9406	Psychoneurotic reaction, other.
	DEVCHORHASIOI OCIC DISORDERS

#### PSYCHOPHYSIOLOGIC DISORDERS

9500 9501 9502 9503 9504	Psychophysiologic skin reaction. Psychophysiologic cardiovascular reaction. Psychophysiologic gastrointestinal reaction. Psychophysiologic nervous system reaction. Psychophysiologic reaction, other.
	DENTAL AND ORAL CONDITIONS

	DENTAL AND ORAL CONDITIONS
9900	Maxilla or mandible, osteomyelitis of.
9901	Mandible, loss of, complete, between angles.
9902	Mandible, loss of approximately one-half.
9903	Mandible, nonunion of.
9904	Mandible, malunion of.
9905	Temporomandibular articulation, limited motion of.
9906	Ramus, loss of whole or part of.
9907	Ramus, loss of less than one-half the substance of,
	not involving loss of continuity.
9908	Condyloid process, loss of, one or both sides.
9909	Coronoid process, loss of.
9910	Maxilla, loss of whole or part of substance of, non- union of, or malunion of.
9911	Hard palate, loss of half or more.
9912	Hard palate, loss of less than half of.
9913	Teeth, loss of, due to loss of substance of body of maxilla or mandible.

#### Pt. 4, App. C

[29 FR 6718, May 22, 1964, as amended at 34 FR 5064, Mar. 11, 1969, 52 FR 44122, Nov. 18, 1987; 53 FR 24938, July 1, 1988]

## APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES

	Diag- nostic Code No.
Abscess: Brain Kidney	8020 7501
Liver Lung	7313 6809
Actinomycosis, lung Addison's disease	6803 7911
Adenitis, secondary	7713
Adenoma, thyroid:	
Nontoxic Toxic	7902 7901
Adhesions:	
Pericardial	7003 7301
PeritoneumAgranulocytosis	7702
Amebiasis	7321
Amputation: Arm:	
Disarticulation	5120
Above deltoid	5121
Below deltoid Feet, both, and hand, one	5122 5103
Feet, both	5107
Finger (digit) individual:	
ThumbIndex	5152 5153
Middle	5154
Ring	5155
Little Fingers (digits) of one hand:	5156
Five	5126
Four, thumb, index, middle, ring	5127
Four, thumb, index, middle, little Four, thumb, index, ring, little	5128 5129
Four, thumb, middle, ring, little	5130
Four, index, middle, ring, little	5131
Three, thumb, index, middle Three, thumb, index, ring	5132 5133
Three, thumb, index, little	5134
Three, thumb, middle, ring	5135
Three, thumb, middle, little Three, thumb, ring, little	5136 5137
Three, index, middle, ring	5138
Three, index, middle, little	5139
Three, index, ring, little Three, middle, ring, little	5140 5141
Two, thumb, index	5142
Two, thumb, middle	5143
Two, thumb, ring	5144 5145
Two, thumb, little Two, index, middle	5145
Two, index, ring	5147
Two, index, little	5148
Two, middle, ring Two, middle, little	5149 5150
Two, ring, little	5151
Forearm:	5400
Above pronator teres  Below pronator teres	5123 5124
Forefoot	5166
Hand, one, and foot, one	5108
Hands, both, and feet, both	5100
Hands, both, and foot, one Hands, both	5102 5106

### APPENDIX C TO PART 4—ALPHABETICAL INDEX

OF DISABILITIES—Continued		
	Diag- nostic Code No.	
Leg:		
With defective stump With loss of natural knee action	5163 5164	
At lower level	5165	
Thigh:	=100	
Disarticulation Upper third	5160 5161	
Middle or lower thirds	5162	
Toe, great	5171	
Toe, other, with removal metatarsal head Toes, all	5172 5170	
Toes, three or more	5173	
Anemia:		
Pernicious	7700 7701	
Secondary	7701	
Aorta or branches	7110	
Arteriovenous, traumatic	7113	
Artery Angioneurotic edema	7111 7118	
Ankylosis:	7110	
Ankle	5270	
Elbow Finger (digit) individual:	5205	
Thumb	5224	
Index	5225	
Middle	5226	
OtherFingers (digits) of one hand, unfavorable:	5227	
Five	5216	
Four	5217	
Three	5218	
Two Hip	5219 5250	
Knee	5256	
Scapulohumeral	5200	
Spine: Complete	5286	
Cervical	5287	
Dorsal	5288	
Lumbar	5289	
Subastragular or Tarsal Wrist	5272 5214	
Anthracosis	6800	
Aphakia	6029	
Aphonia, organic	6519	
Cerebral	8046	
General	7100	
Obliterans	7114	
Arteriosclerotic heart disease	7005	
Atrophic (rheumatoid)	5002	
Gonorrheal	5004	
Hypertrophic (degenerative) Other types	5003 5009	
Pneumococcic	5009	
Streptococcic	5008	
Syphilitic	5007	
Traumatic Typhoid	5010 5006	
Aspergillosis, lung	6807	
Asthma, bronchial	6602	
Astragalectomy	5274 8107	
Athetosis	8107	
Muscular, progressive	8023	
Ovaries, both	7620	

## APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES—Continued

OF DISABILITIES—Continued		OF DISABILITIES—Continued			
	Diag- nostic Code No.		Diag- nostic Code No.		
Testis, both	7523	Other	6018		
Auditory canal, disease	6210	Coccyx	5298		
Avitaminosis	6313	Cushing's syndrome	7907		
Beriberi	6314	Cyclitis	6004		
Blastomycosis, lung	6805	Cystitis:			
Blindness, anatomical loss, one eye:	0000	Chronic	7512		
Other blind (5/200 or less)	6063 6064	Interstitial (Hunner)	7513 6031		
Other impaired (20/200 or less) Other impaired	6065	Dacryocystitis  Deafness	0031		
Other normal	6066	0% evaluation based on Table VII	6100		
Blindness, light perception only:	0000	10% evaluation based on Table VII	6101		
Both eyes	6062	20% evaluation based on Table VII	6102		
One eye:		30% evaluation based on Table VII	6103		
Other blind, 5/200 or less	6067	40% evaluation based on Table VII	6104		
Other impaired, 20/200 or less	6068	50% evaluation based on Table VII	6105		
Other impaired	6069	60% evaluation based on Table VII	6106		
Other normal	6070	70% evaluation based on Table VII	6107		
Blindness, light perception only and loss or loss of	6050-	80% evaluation based on Table VII	6108 6109		
use of hands and/or feet		100% evaluation based on Table VII	6110		
Blindness, total (5/200 or less):	6061	Deflection, nasal septum	6502		
Both eyes	6071	Dermatitis, exfoliativa	7817		
One eye:	0071	Dermatophytosis	7813		
Other impaired, (20/200 or less)	6072	Diabetes mellitus	7913		
Other impaired	6073	Diabetes insipidus	7909		
Other normal	6074	Diaphragm, rupture	5324		
Blindness, partial (20/200 or less):		Dilation, aneurysmal artery	7112		
Both eyes	6075	Diplopia	6090		
One eye:		Disease:			
Other impaired	6076	Addison's	7911		
Other normal	6077	Hodgkin's	7709 5279		
Blindness, partial:	6078	Morton's Raynaud's	7117		
Both eyes One eye only	6078	Dislocation:	7117		
Block, auricular ventricular	7015	Cartilage, semilunar	5258		
Bones, Caisson disease of	5011	Lens, crystalline	6033		
Bones and joints, tuberculosis of	5001	Disorders, mental:			
Bronchiectasis	6601	Organic brain disorders:			
Bronchitis	6600	Acute brain syndrome	9300		
Buerger's disease	7115	Chronic brain syndrome associated with:			
Brucellosis	6316	Central nervous system syphilis	9301		
Bursitis	5019	Intracranial infections other than	0202		
Calsson disease	5011 7515	syphilis Intoxication	9302 9303		
Calculus, bladder Cataract:	7515	Brain trauma	9303		
Senile and others	6028	Cerebral arteriosclerosis	9305		
Traumatic	6027	Circulatory disturbance other than			
Cervicitis	7612	cerebral arteriosclerosis	9306		
Cholangitis	7316	Convulsive disorder (idiopathic epi-			
Cholecystitis	7314	lepsy)	9307		
Cholelithiasis	7315	Disturbance of metabolism, growth or			
Cholera, Asiatic	6300	nutrition	9308		
Chorea:	0400	Intracranial neoplasm	9309		
Huntington's	8106	Diseases of unknown or uncertain	9310		
Sydenham's	8105 6005	causeUnknown cause	9310		
Choroiditis	5278	Psychoneurotic disorders:	9311		
Cirrhosis of liver	7312	Anxiety reaction	9400		
Claudication, intermittent	7116	Dissociative reaction	9401		
Coccidioidomycosis	6821	Conversion reaction	9402		
Colitis:		Phobic reaction	9403		
Mucous (See Colon syndrome, irritable)	7319	Obsessive compulsive reaction	9404		
Spastic (See Colon syndrome, irritable)	7319	Depressive reaction	9405		
Ulcerative	7323	Psychoneurotic reaction, other	9406		
Collapse, lung, permanent	6813	Psychophysiologic disorders:			
Colon syndrome, irritable	7319	Psychophysiologic skin reaction	9500		
Congestion, lung, passive Conjunctivitis:	6817	Psychophysiologic cardiovascular reaction	9501		
Trachomatous	6017	Psychophysiologic gastrointestinal reac- tion	9502		
11aU1U11aUU3	0017	uOII	3002		

#### Pt. 4, App. C

## APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES—Continued

OI DISABILITIES CONTINUCA		or Bioribicines Continuou	
	Diag- nostic Code No.		Diag- nostic Code No.
Psychophysiologic nervous system reac-		Noncongestive	601
tion	9503	Gout	501
Psychophysiologic reaction, other	9504	Growths, new benign:	001
Psychotic disorders:		Bones, joints and muscles	501
Schizophrenic reaction:		Brain	800
Simple type	9200	Digestive system	734
Hebephrenic type	9201	Ear	620
Catatonic type	9202	Endocrine system	791
Paranoid type Chronic undifferentiated type	9203 9204	Eyeball and adnexa	601
Other	9204	Genitourinary system	752 682
Manic depressive reaction	9206	RespiratorySkin	781
Psychotic depressive reaction	9207	Spinal cord	802
Paranoid reaction	9208	Growths, new, malignant:	002
Involutional psychotic reaction	9209	Bones	501
Psychotic reaction, other	9210	Brain	800
Distomiasis, intestinal	7324	Digestive system	734
Diverticulitis, intestinal	7327	Ear	620
Diverticulum of esophagus	7205	Endocrine system	791
Dupuytren's contracture—see Ankylosis, fingers.  Dysentery, bacillary	7322	Eyeball only	601
Ectropion	6020	Genitourinary system	752
Eczema	7806	Gynecological system or mammary glands	762
Edema, angioneurotic	7118	Respiratory	681
Embolism, brain	8007	Skin	781
Emphysema (No DC; follows DC 6602).		Spinal cord	802 528
Encephalitis	8000	Hallux rigidusHallux valgus	528 528
Endocarditis, bacterial, subacute	7001	Hammer toe	528
Enteritis	7325	Hematomyelia	801
Enterocolitis	7326	Hemorrhage:	001
Entropion	6021	Brain	800
Enucleation, eye, see Blindness.		Intraocular	600
Epilepsy:	0040	Hemorrhoids	733
Grand mal Petit mal	8910 8911	Hepatitis, infectious	734
Jacksonian	8912	Hernia:	
Diencephalic	8913	Femoral	734
Psychomotor	8914	Hiatal	734
Epiphora (lacrymal duct)	6025	Inguinal	733
Erythromelalgia	7119	Muscle	532 733
Eyelids, loss of portion of	6032	Ventral Hodgkin's disease	770
Fever:		Hydrarthrosis, intermittent	501
Hemoglobinuric, see Malaria.	2042	Hydronephrosis	750
Malta	6316	Hyperadrenia	791
Oroya	6306	Hyperparathyroidism	790
Relapsing	6308 6309	Hyperpituitarism:	
Undulant	6316	Acromegaly or gigantism	790
Fibrillation, auricular:	0010	Cushing's syndrome	790
Paroxysmal	7011	Hypertensive heart disease	700
Permanent	7012	Hypertensive vascular disease	710
Filariasis	6305	Hyperthyroid heart disease	700
Fistula:		Hyperthyroidism	790
Ano	7335	Hypoadrenia	791
Bladder	7516	Hypoparathyroidism	790
Bronchocutaneous or bronchopleural	6812	Hypopituitarism	790
Intestine	7330	Hypothyroidism	790
Rectovaginal	7624	Immersion footImpairment:	712
Urethra	7625 5254	Auditory acuity, see Deafness.	
Flail hip	5254 5276	Clavicle	520
Flatfoot (pes planus) acquiredFlutter, auricular	7010	Elbow	520 520
Fracture, vertebra, residuals of	5285	Eye (field vision)	608
Frozen feet	7122	Eye (muscle function)	609
Gastritis, atrophic (see DC 7307).	1122	Femur	525
Gastritis, hypertrophic	7307	Humerus	520
Genu, recurvatum	5263	Knee	525
Glaucoma:		Radius	521
Congestive	6012	Sphincter control	733

APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES—Continued

OF DISABILITIES—Continued		OF DISABILITIES—Continued			
	Diag- nostic Code No.		Diag- nostic Code No.		
Supination and pronation	5213	Thigh	525		
Thigh, motion Tibia and fibula	5253 5262	Limitation of flexion and extension: Forearm	520		
Ulna	5211	Limitation of motion:	320		
Visual acuity, see Blindness.	02	Ankle	527		
Infarction of myocardium	7006	Arm	520		
Injury:		Cervical	529		
Bladder	7517	Dorsal Lumbar	529 529		
Gall bladder Eye, unhealed	7317 6009	Temporomandibular articulation	990		
Foot	5284	Wrist	521		
Larynx	6517	Limitation, pronation	521		
Lips	7201	Limitation, supination	521		
Liver	7311	Limitation of muscle function, eye	609		
Mouth	7200	Loss:	681		
Muscle:	E22E	Auricle or deformity	620		
FacialGroup I	5325 5301	Condyloid process	990		
Group II	5301	Coronoid process	990		
Group III	5303	Eyebrows	602		
Group IV	5304	Eyelashes	602		
Group V	5305	Mandible: Complete	990		
Group VI	5306	One-half	990		
Group VIII	5307	Maxilla	991		
Group VIIIGroup IX	5308 5309	Teeth	991		
Group X	5310	Nose, loss of part, or scars	650		
Group XI	5311	Palate, hard:			
Group XII	5312	Half or more	991		
Group XIII	5313	Less than half Ramus:	991		
Group XIV	5314	Less than one-half substance	990		
Group XV	5315	Whole or part	990		
Group XVIII	5316 5317	Skull, part	529		
Group XVIIGroup XVIII	5318	Smell, sense of	627		
Group XIX	5319	Taste, sense of	627		
Group XX	5320	Tongue or partOthers, see Amputation, removal, etc.	720		
Group XXI	5321	Loss of use:			
Group XXII	5322	Feet, both	511		
Group XXIII	5323	Feet, both, and hand, one	510		
Pleural cavity Prostate	6818 7527	Foot, one	516		
Sacroiliac	5294	Hand, one	512		
Spleen	7707	Hand, one, and foot, one Hands, both, and feet, both	511 510		
Stomach, residuals	7310	Hands, both, and foot, one	510		
Tongue, whole or part	7202	Hands, both	510		
Intervertebral disc	5293	Lupus, erythematosus, discoid	780		
Iritis Kala-azar	6003 6301	Lupus, erythematosus systemic (disseminated)	635		
Keratitis	6001	Lupus, vulgaris	781		
Labyrinthitis	6204	Lymphogranulomatosis Malaria	770 630		
Lagophthalmos	6022	Malunion:	030		
Laryngectomy	6518	Clavicle	520		
Laryngitis	6516	Os calcis (or astragalus)	527		
Leishmaniasis:	7007	Mandible	990		
Americana	7807 7808	Maxilla (or nonunion)	991		
Old World Lens, crystalline, dislocation of	6033	Scapula Tarsal or metatarsal (or nonunion)	520 528		
Leprosy	6302	Others, see Impairment.	320		
Leukemia	7703	Mastoiditis	620		
Limitation of extension:		Meniere's disease	620		
Forearm	5207	Meningitis, cerebrospinal	801		
Leg	5261	Mental disorders—see Disorders, mental.	F07		
Thigh Limitation of field vision	5251 6080	Metatarsalgia	527		
Limitation of flexion:	0000	Metritis Migraine	761 810		
Forearm	5206	Muscle injury, see Injury, muscle.	010		
Leg		Myasthenia gravis	802		

#### Pt. 4, App. C

## APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES—Continued

OF DISABILITIES—Continued		OF DISABILITIES—Continued	
	Diag- nostic Code No.		Diag- nostic Code No.
Mycosis, lung, unspecified	6808	Obturator	8628
Myelitis	8010	External cutaneous, thigh	8629
Myositis	5021	Ilio-inguinal	8630
Myositis ossificans	5023 8108	Non-union of bones:  Mandible	9903
Nephritis, chronic	7502	Radius and Ulna	5210
Nephrolithiasis	7508	Tibia and fibula	5262
Nephrosclerosis, arteriolar	7507	Others, see Impairment.	
Neuralgia:		Nystagmus, central	6016
Cranial nerves: Fifth (trigeminal)	8405	Oophoritis Oroya fever	7615 6306
Seventh (facial)	8407	Osteitis deformans	5016
Ninth (glossopharyngeal)	8409	Osteomalacia	5014
Tenth (pneumogastric, vagus)	8410	Osteomyelitis, jaw	9900
Eleventh (spinal accessory, external		Osteomyelitis	5000
branch)	8411	Osteoporosis	5013
Twelfth (hypoglossal) Peripheral nerves:	8412	Otitis externa	6210 6203
Upper radicular group	8710	Otitis interna	6203
Middle radicular group	8711	Catarrhal	6201
Lower radicular group	8712	Suppurative	6200
All radicular groups	8713	Otosclerosis	6202
Musculospiral	8714	Palsy, bulbar	8005
Median	8715	Paralysis:	
Ulnar	8716	Accommodation	6030 8004
Musculocutaneous Circumflex	8717 8718	Agitans Paralysis, nerve:	8004
Long thoracic	8719	Cranial:	
Sciatic	8720	Fifth (trigeminal)	8205
External popliteal	8721	Seventh (facial)	8207
Musculocutaneous (superficial peroneal)	8722	Ninth (glossopharyngeal)	8209
Anterior tibial	8723	Tenth (pneumogastric, vagus)	8210
Internal popliteal Posterior tibial	8724 8725	Eleventh (spiral accessory, external branch)	8211
Anterior crural	8726	Twelfth (hypoglossal)	8212
Internal saphenous	8727	Peripheral:	0212
Obturator	8728	Upper radicular group	8510
External cutaneous, thigh	8729	Middle radicular group	8511
Ilio-inguinal	8730	Lower radicular group	8512
Neuritis, optic	6026	All radicular groups  Musculospiral	8513 8514
Cranial nerves:		Median	8515
Fifth (trigeminal)	8305	Ulnar	8516
Seventh (facial)	8307	Musculocutaneous	8517
Ninth (glossopharyngeal)	8309	Circumflex	8518
Tenth (pneumogastric, vagus)	8310	Long thoracic	8519
Eleventh (spinal accessory, external	8311	Sciatic	8520
branch) Twelfth (hypoglossal)	8312	External popliteal	8521 8522
Peripheral:	0012	Anterior tibial	8523
Upper radicular group	8610	Internal popliteal	8524
Middle radicular group	8611	Posterior tibial	8525
Lower radicular group	8612	Anterior crural	8526
All radicular groups	8613	Internal saphenous	8527
Musculospiral Median	8614	Obturator thigh	8528
Ulnar	8615 8616	External cutaneous, thighIlio-inguinal	8529 8530
Musculocutaneous	8617	Paramyoclonus multiplex	8104
Circumflex	8618	Pellagra	6315
Long thoracic	8619	Pemphigus	7815
Sciatic	8620	Penis, deformity of	7522
External popliteal	8621	Perforation: Tympanic membrane	6211
Musculocutaneous (superficial peroneal) Anterior tibial	8622 8623	Pericarditis Periostitis	7002 5022
Internal popliteal	8624	Periositis	5022
Posterior tibial	8625	Pes planus	5276
Anterior crural	8626	Phlebitis	7121
Internal saphenous	8627	Pinta	7810

APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES—Continued

OF DISABILITIES—Continued		OF DISABILITIES—Continued		
	Diag- nostic Code No.		Diag- nostic Code No.	
Plague	6307	Frontal	6512	
Pleurisy:		Maxillary	6513	
Purulent (empyema)	6811	Pansinusitis	6510	
Serofibrinous	6810	Sphenoid	6514	
Pluriglandular syndrome	7912	Spasm, esophagus	7204	
Pneumoconiosis	6802	Splenectomy	7706	
Pneumonectomy	6815	Sporotrichosis, lung	6806	
Pneumothorax, spontaneous	6814	Stenosis:		
Poliomyelitis, anterior	8011	Larynx	6520	
Polycythemia	7704	Stomach	7309	
Pregnancy, surgical complications of	7623	Strain, lumbosacral	5295	
Prolapse:		Streptotrichosis, lung	6804	
Rectum	7334	Stricture:		
Uterus	7621	Esophagus	7203	
Pronation, limitation of	5213	Rectum, anus	7333	
Pruritis, ani	7337	Ureter	7511	
Psoriasis	7816	Urethra	7518	
Psychiatric disorders, see Disorders, mental.		Supination, limitation of	5213	
Pterygium	6034	Symblepharon	6091	
Ptosis, eyelid	6019	Syndrome:		
Purpura, hemorrhagica	7705	Cushing's	7907	
Pyelitis	7503	Intervertebral disc	5293	
Pyelonephritis, chronic	7504	Meniere's	6205	
Raynaud's disease	7117	Pluriglandular	7912	
Removal:	/ / / /	Postgastrectomy	7308	
	6207	Synovitis	5020	
Auricle or deformity Cartilage, semilunar	5259	Syphilis:	3020	
Coccyx	5298	Cerebrospinal	8013	
			8014	
Gall bladder	7318	Meningovascular		
Kidney	7500	Unspecified	6310	
Mammary glands	7626	Syphilitic heart disease	7004	
Ovaries, both	7619	Syringomyelia	8024	
Penis, half or more	7520	Tabes dorsalis	8015	
Penis, glans	7521	Tachycardia:		
Prostate, or resection	7526	Paroxysmal	7013	
Ribs	5297	Sinus	7014	
Testis	7524	Tenosynovitis	5024	
Uterus	7618	Thrombo-anglitis obliterans	7115	
Uterus and ovaries	7617	Thrombophlebitis	7121	
Others, see Amputation, loss, etc.		Thrombosis, brain	8008	
Resection:		Tic, convulsive	8103	
Intestine:		Tinea barbae	7814	
Large	7329	Tinnitus	6260	
Small	7328	Tuberculosis:		
Stomach	7308	Adenitis, tuberculous:		
Retina, detachment of	6008	Axillary	7711	
Retinitis	6006	Cervical	7710	
Rheumatic fever	6309	Inguinal	7712	
Rheumatic heart disease	7000	Bladder	7514	
Rhinitis: Atrophic	6501	Bones and joints	5001	
Rupture, diaphragm	5324	Epididymo-orchitis, tuberculous	7525	
Salpingitis	7614	Eye	6010	
Scars:	7014	Kidney	7505	
Burns, second degree	7802	Laryngitis, tuberculous	6515	
	7801	Luposa	7811	
Burns, third degree	7800	Miliary	6311	
Head, etc., disfiguring			0311	
Retina	6011	Nonpulmonary, inactive (see § 4.89)	7331	
Superficial, tender	7804	Peritonitis, tuberculous		
Superficial, with ulceration	7803	Pleurisy, tuberculous	6732	
Others	7805	Pulmonary:		
Scleritis	6002	Active:	0704.5	
Sclerosis:		Far advanced	6701 &	
Amyotrophic, lateral	8017		6707	
Multiple	8018	Moderately advanced	6702 &	
Scotoma, pathological	6081		6708	
Shortening, leg	5275	Minimal	6703 &	
Silicosis	6801		6709	
Sinusitis:		Advancement unspecified	6704 &	
Ethmoid	6511		6710	

#### Pt. 6

### APPENDIX C TO PART 4—ALPHABETICAL INDEX OF DISABILITIES—Continued

	Diag- nostic Code No.
Inactive:	
Far advanced	6721 &
	6725
Moderately advanced	6722 &
	6726
Minimal	6723 &
	6727
Advancement unspecified	6724 &
	6728
Tympanic membrane, perforation of	6211
Typhus, scrubUlcer:	6317
Duodenal	7305
Gastric	7304
Marginal	7306
Undescended testis (see Note under DC 7524).	
Uterus, displacement of	7622
Ureterolithiasis	7510
Uveitis	6000
Vaginitis	7611 7120
Varicose veins	7120 7812
Verruga peruana Vertebra, fracture	5285
Visceroptosis	7342
Vision, impairment of, see Blindness.	1342
Vulvovaginitis	7610
Weak foot	5277
Wound, incised, abdominal wall	7341

[29 FR 6718, May 22, 1964, as amended at 34 FR 5064, Mar. 11, 1969; 52 FR 44122, Nov. 18, 1987; 53 FR 24938, July 1, 1988]

#### PART 5—[RESERVED]

### PART 6—UNITED STATES GOVERNMENT LIFE INSURANCE

AGE

Sec.

6.1 Misstatement of age.

PREMIUMS

6.2 Premium rate.

POLICIES

6.3 Incontestability of United States Government life insurance.

BENEFICIARY OF UNITED STATES GOVERNMENT LIFE INSURANCE

- 6.4 Proof of age, relationship and marriage.
- 6.5 Conditional designation of beneficiary.
- 6.6 Change of beneficiary.
- 6.7 Claims of creditors, taxation.

#### OPTIONAL SETTLEMENT

- 6.8 Selection, revocation and election.
- 6.9 Election of optional settlement by beneficiary.

6.10 Options.

#### DIVIDENDS

6.11 How dividends are paid.

6.12 Special dividends.

LOANS

6.13 Policy loans.

#### CASH VALUE

- 6.14 Cash value; other than special endowment at age 96 plan policy.
- 6.15 Cash value; special endowment at age 96 plan policy.
- 6.16 Payment of cash value in monthly installments.

#### INDEBTEDNESS

6.17 Collection of any indebtedness.

TOTAL PERMANENT DISABILITY BENEFITS

6.18 Other disabilities deemed to be total and permanent.

#### DEATH BENEFITS

6.19 Evidence to establish death of the insured.

DETERMINATION OF LIABILITY UNDER SECTIONS 302 AND 313, WORLD WAR VETERANS' ACT, 1924, SECTIONS 607 AND 602(V)(2), NATIONAL SERVICE LIFE INSURANCE ACT, 1940, AS AMENDED, AND SECTIONS 1921 AND 1957 OF TITLE 38, UNITED STATES CODE

6.20 Jurisdiction.

#### APPEALS

6.21 Guardian: definition and authority.

AUTHORITY: 38 U.S.C. 501, 1940-1963, 1981-1988, unless otherwise noted.

#### AGE

#### § 6.1 Misstatement of age.

If the age of the insured under a United States Government life insurance policy has been understated, the amount of the insurance payable under the policy shall be such exact amount as the premium paid would have purchased at the correct age; if overstated, the excess of premiums paid shall be refunded without interest. Guaranteed surrender and loan values will be modified accordingly. The age of the insured will be admitted by the Department of Veterans Affairs at any time upon satisfactory proof.

[13 FR 7089, Nov. 27, 1948. Redesignated at 61 FR 29024, June 7, 1996]